Sem iclassical tim e-evolution of the reduced density matrix and dynam ically assisted generation of entanglem ent for bipartite quantum systems

Ph. Jacquod¹

¹D epartem ent de Physique Theorique, Universite de Geneve, CH-1211 Geneve 4, Switzerland (D ated: February 16, 2004)

Two particles, initially in a product state, become entangled when they come together and start to interact. Using sem iclassical methods, we calculate the time evolution of the corresponding reduced density matrix 1, obtained by integrating out the degrees of freedom of one of the particles. To quantify the generation of entanglement, we calculate the purity $P(t) = Tr[1(t)^2]$. We not that entanglement generation sensitively depends (i) on the interaction potential, especially on its strength and range, and (ii) on the nature of the underlying classical dynamics. Under general statistical assumptions, and for short-ranged interaction potentials, we not that P(t) decays exponentially fast if the two particles are required to interact in a chaotic environment, whereas it decays only algebraically in a regular system. In the chaotic case, the decay rate is given by the golden rule spreading of one-particle states due to the two-particle coupling, but cannot exceed the system's Lyapunov exponent.

PACS num bers: 05.45 M t,03.65.U d,05.70 Ln,03.67.-a

When two systems (...) enter into temporary interaction (...), and when after a time of mutual in uence the systems separate again, then they can no longer be described in the same way as before, viz. by endowing each of them with a representative of its own. This is how entanglement was characterized by Schrodinger almost seventy years ago [1]. Entanglement is arguably the most puzzling property of multipartite quantum systems, and often leads to counterintuitive predictions due to, in E instein's words, spooky action at a distance. Entanglement has received a renewed, intense interest in recent years in the context of quantum information theory [2].

In the spirit of Schrodinger's above form ulation, one is naturally led to ask the following question: W hat determ ines the rate of entanglem ent production in a dynam ical system ?". Is this rate mostly determ ined by the interaction between two, initially unentangled particles, or does it depend on the underlying classical dynamics? This is the question we address in this paper. Previous attempts to answer this question have mostly focused on num erical investigations [3, 4, 5, 6, 7, 8], with claims that entanglem ent is favored by classical chaos, both in the rate it is generated [3, 4, 7] and in the maxim alam ount it can reach [5]. In particular, strong num erical evidences have been given by M iller and Sarkar for an entanglem ent production rate given by the system 's Lyapunov exponents [4]. These ndings have how ever been recently challenged by Tanaka et al. [6], whose num erical ndings show no increase of the entanglem ent production rate upon increase of the Lyapunov exponents in the strongly chaotic but weakly coupled regime, in agreement with their analytical calculations relating the rate of entanglem ent production to classical time correlators. Ref. [6] is seen ingly in a paradoxical disagreem ent with the alm ost identical analytical approach of R ef. [7], where entanglem ent production was found to be faster in chaotic system s than in regular ones [9]. This controversy thus calls for a better analytical understanding of the problem .

We present a sem iclassical calculation of the time-

evolved density matrix (t) for two interacting, distinguishable particles. In the usual way, entanglem ent is quanti ed by the properties of the reduced density ma- $T_{\mathfrak{P}}[(t)]$, obtained from the two-particle trix 1 (t) density matrix by tracing over the degrees of freedom of one (say, the second) particle. At t = 0, the two particles are in a product state of two narrow wavepackets. W e will quantify entanglem ent by calculating the purity $Tr[_1(t)^2]$ which varies from 0 for fully entangled P (t) to 1 for factorizable two-particle states [10] (the discussion of how this could translate into the violation of a Bell. inequality [11] is postponed to later works). Com pared to other m easures of entanglem ent such as the von Neum ann entropy or the concurrence, the purity presents the advantage of being analytically tractable. For the weak coupling situation we are interested in here, num erical works have moreover shown that von Neum ann and lin-1 P (t) behave very similarly [6]. ear entropy S_{lin} W e thus expect the purity to give a faithful and generic m easure of entanglem ent. W e note that our sem iclassical approach is straightforwardly extended to the case of undistinguishable particles, provided the nonfactorization of the reduced density matrix due to particle statistics is properly taken care of [12].

O ur approach is rem iniscent of the sem iclassicalm ethods developed by Jalabert and Pastawski [13, 14] (see also [15]) in the context of the Loschm idt E cho. W e will show how the o -diagonalm atrix elements of $_1$ are related to classical action correlators, in a similar way as in Ref. [6, 7]. Under nonrestrictive statistical assum ptions, we nd that, following an initial transient where $_1$ relaxes but remains almost exactly pure, entanglem ent production is exponential in chaotic system s, while it is algebraic in regular system s. The asym ptotic rate of entanglem ent production in chaotic system s depends on the strength of the interaction between the two particles, and is explicitly given by a classical time-correlator. W e note that, as is the case for the Loschm idt E cho [16, 17], this regime is also adequately captured by an approach based on R andom M atrix T heory (RM T) [22] { the timecorrelator is then replaced by the golden rule spreading of one-particle states due to the interaction. For stronger coupling how ever, the dom inant stationary phase solution becom es interaction independent, and is determ ined only by the classical dynam ics, the system 's Lyapunov exponents giving an upper bound for the rate of entanglem ent production. The crossover between the two regim es occurs once the golden rule width becom es com parable to the system 's Lyapunov exponent. Still, one has to keep in m ind that long-ranged interaction potentials can lead to signi cant m odi cations of this picture, especially at short tim es, due to an anom alously slow vanishing of o – diagonal m atrix elem ents of $_1$ within a bandwidth set by the interaction range.

We start with an initial two-particle product state j_i j_1 ; 2i. The state of each particle is a j_i Gaussian wavepacket $_{1;2}(y) = (^{2})^{d=4} \exp[ip_{0}$ (y $r_{1,2}$) jy $r_{2,2} = 2^{2}$. We write the two-particle H am iltonian as H = H I + I H + U, i.e. the two particles are subjected to the same Ham iltonian dynamics. At this point, we only specify that the interaction potential U is smooth, depends only on the distance between the particles, and is characterized by a typical length scale (which can be its range, or the scale over which it uctuates). Setting h 1, the two-particle density matrix evolves according to $(t) = \exp[iH t]_0 \exp[iH t]$ starting initially w_Rith $_0 = j_1; _2$ ih $_1; _2$ j. The elements drhx;rj (t) jy;ri of the reduced density $_{1}(x;y;t) =$ m atrix read

$$Z Z$$

$$I (x;y;t) = (2)^{d} dr = \frac{4}{i=1} dy_{i} \exp[f(y_{1} - y_{1})^{2} + (y_{2} - y_{2})^{2} + (y_{3} - y_{1})^{2} + (y_{4} - y_{2})^{2}g=2^{2}]$$

$$\exp[ip_{0}(y_{1} + y_{2} - y_{3} - y_{4})]hx;rjexp[iHt]jy_{1};y_{2}ihy_{3};y_{4}jexp[iHt]jy;ri: (1)$$

W e next introduce the sem iclassical two-particle propagator

$$hx;rjexp[iHt]_{y_1};y_2i = (i)^{d} C_{s;s^0}^{1=2} exp[ifS_s(y_1;x;t) + S_{s^0}(y_2;r;t) + S_{s;s^0}(y_1;x;y_2;r;t) - \frac{1}{2}(s + s^0)g];$$

which is expressed as a sum over pairs of classical trajectories, labelled s and s⁰, respectively connecting y_1 to x and y_2 to r in the time t. Each such pair of paths gives a contribution containing one- (denoted by S_s and S_{s^0}) and two-particle (denoted by $S_{s;s^0} = \frac{R_t}{0} dt_1 U$ ($q_s(t_1); q_{s^0}(t_1)$) [23]) action integrals accumulated along s and s⁰, a pair of M aslov indices s and s^0 , and the determ inant $C_{s;s^0}$ of the stability matrix corresponding to the two-particle dynamics in the (2d) dimensional space. (W if the above denition, $C_{s;s^0}$ is real and positive.) We consider su ciently smooth integrates varying over a distance much larger than . We thus set $S_{s;s^0}(y_1;x;y_2;r;t) ' S_{s;s^0}(r_1;x;r_2;r;t)$ (still we keep in m ind that r_1 and r_2 , taken as arguments of the two-particle actions in y_i r_j (i = 1;:::4; j = 1;2). We consider the weak coupling regime, where the one-particle actions vary faster than their two-particle counterpart. We thus perform a stationary phase approximation on the S's to get the sem iclassical reduced density matrix as

$$I_{1}(x;y;t) = (4^{2})^{d} \exp[ifS_{s}(r_{1};x;t) S_{1}(r_{1};y;t)g]$$

$$Z_{1} X_{0}^{s;1} X_{0}^{s;1} X_{1;s^{0}} \exp[ifS_{s;s^{0}}(r_{1};x;r_{2};r;t) S_{1;s^{0}}(r_{1};y;r_{2};r;t)g]; \qquad (2)$$

$$M_{1} X_{s;s^{0}} = C_{1}^{1=2} X_{1;s^{0}} \exp[\frac{i}{2}(s_{1} s_{1} s_{1})] \exp[\frac{2}{2}f(p_{1} s_{1} p_{1})^{2} + (p_{1} s_{1} p_{1})^{2}g]; \qquad (2)$$

It is straightforward to see that $Tr[_1(t)] = 1$ and $_1(x;y;t) = [_1(y;x;t)]$, as required. Enforcing a further stationary phase condition on Eq. (2) amounts to perform ing an average over di erent initial conditions $r_{1;2}$. It results in $s = s^0$, x = y, and thus $h_1(x;y;t)i = _{x,y} = ($ is the system 's volum e), i.e. only diagonal elements of the reduced density m atrix have a nonvanishing average (the ergodicity of $h_1(x;x;t)i$ is due to the average over initial conditions). For each initial condition, $_1$ has how ever nonvanishing o -diagonal matrix elements, with a zero-centered distribution whose variance is given by $h_1(x;y;t)_1(y;x;t)i$. Squaring Eq. (2), averaging over $r_{1;2}$ and enforcing a stationary phase approximation on the S's, one gets

$$h_{1}(x;y;t)_{1}(y;x;t)i = (4^{2})^{2d} drdr^{0} M_{s;s^{0}} M_{l,m} M_{l;s^{0}}^{y} M_{s,m}^{y} hFi;$$
(3)

$$F = \exp[ifS_{s;s^{0}}(r_{1};x;r_{2};r;t) \quad S_{l;s^{0}}(r_{1};y;r_{2};r;t)g] \exp[ifS_{l,m}(r_{1};y;r_{2};r^{0};t) \quad S_{s,m}(r_{1};x;r_{2};r^{0};t)g]: (4)$$

O ur analysis of Eqs. (3-4) starts by noting that hj $_1 f$ is given by the sum of two positive contributions. First, those particular paths for which $r = r^0$ and $s^0 = m$, accumulate no phase (F = 1) and thus have to be considered separately. On average, their contribution does not depend on x nor y, and decays in time only because of their decreasing measure with respect to all the paths with $r \in r^0$. Their average contribution is given by

$${}^{2}_{0}(t) = (4 \ {}^{2})^{2d} \ {}^{2} dr_{1} dr_{2} dr \frac{1}{s;lm} f M_{lm} f M_{lm} f / \frac{2 \exp[((1+2)t]; \text{ chaotic;} (5)]{}_{2} \frac{t_{0}}{t} \frac{2}{t} \frac{1}{t} \frac{1$$

This is obtained by taking half of the C's in the M's as Jacobians for the coordinate transform ation from r's to p's, and replacing the other half by the large time limit $[18]C_{s,m}$ $C_s^{(1)}C_m^{(2)}$ / $(t_0=t)^2$ for regular, and $\exp[((_1+_2)t]]$, for chaotic systems (In Eq. (5), we explicitly wrote particle indices; in our case where the two particles are subjected to the same H am iltonian, the Lyapunov exponents are equal, $_1 = _2$). Two facts are noticeable here: these contributions do not depend on the interaction strength, moreover, Eq. (5) gives a lower bound for the decay of hj $_1j$ i.

The second, generic contributions to hj₁² i decay in time with hF i. From Eq. (4), it is natural to expect that hF i is a decreasing function of j_x y j and t only. Sum s and integrals in Eq. (3) can then be performed separately to get h₁(x;y;t)₁(y;x;t)i = $\frac{2}{0}$ (t) + hF (x;y;t)i = 2 , with

$$F (x;y;t)i = \exp 2 dt_1 dt_2 hU (q_s (t_1);q_{s^0} (t_1)) U (q_s (t_2);q_{s^0} (t_2))i$$

$$Z^0_t$$

$$exp + 2 dt_1 dt_2 hU (q_s (t_1);q_{s^0} (t_1)) U (q_1 (t_2);q_{s^0} (t_2))i : (6)$$

The behavior of F(x;y;t) is sensitively depends on the distance \dot{x} y jbetween the endpoints of the classical trajectories $q_s(t)$ and $q_1(t)$ connecting r_1 to x and y respectively. For small \dot{x} y j the second line on the right-hand side of Eq. (6) almost compensates the rst one [20], and a Taylor expansion of the di erence of the two-particle action integrals in Eq.(4) gives in low est order

The reduced density matrix has a Gaussian decay away from the diagonal, which we expect to hold for \dot{x} yj (the distance over which the two terms in Eq. (6) almost cancel each other), and for short enough times t < (after which very di erent, uncorrelated classical paths l and s m ay still reach two neighboring endpoints). An estimate for

is given by the time it takes for two initial conditions within a distance (this is the uncertainty in r_1) to move away a distance / from each other. In a chaotic system, this is the Ehrenfest time = $\frac{1}{1}$ ln(=), while in a regular system, is much longer, = 0 ([=]) [19]. At larger distances, j_x yj, hF i is given by the rst term on the right-hand side of Eq. (6). Note that, because the four classical paths in that term come in two pairs, the dependence on j_x y j vanishes.

This concludes our sem iclassical calculation of the reduced density matrix for interacting two-particle system s. We have learned that the variance of o -diagonal matrix elements of $_1$ is determined by classical correlators, with the important caveat that they are bound downward by the expressions given in Eq. (5). For the rest of the discussion, we note that, provided the correlators in Eqs. (6) and (7) decay faster than $/ j_1 \pm j^1$, the o -diagonal matrix elements exhibit a dominant exponential decay in time. This condition is rather nonrestrictive and is surely satis ed in a chaotic system . We therefore assume from now on a fast decay of the correlations,

Г

 $h U (q_s (t_1); q_{s^0} (t_1)) U (q_s (t_2); q_{s^0} (t_2)) i = (t_1 \ \xi);$ (8)

$$h \mathfrak{d}^{(s)} U (q_s (t_1); q_{s^0} (t_1)) \mathfrak{d}^{(s)} U (q_s (t_2); q_{s^0} (t_2)) \mathfrak{i} = ; (t_1 \mathfrak{b}):$$
(9)

Entanglement is quantied by the purity P (t) = dxdyh_1(x;y;t)_1(y;x;t)i, which is straightforward to

compute from Eqs. (3-9). We get three distinct regimes of decay: (i) an initial regime of classical relaxation for t < . During that time, 1 evolves from a pure, but localized 1 (0) = $jr_1 ihr_1 j$ to a less localized, but still almost pure 1 (t), with an algebraic purity decay obtained from Eqs. (7) and (9) as P (t <) ' 1 (=2 t) $^{d=2}$ (even in the case of a correlator (9) saturating at a – nite value for j_{1} j_{2} ! 1, which m ay occur in regular systems, this initial decay will still be algebraic / t d); (ii) a regime where quantum coherence develops between the two particles so that $_{1}$ becomes a mixture. From

Eq. (5) and the rst line of Eq. (6) with Eq. (8) one gets P (t) ' exp [m in (2 ; $_1 + _2$)t] in a chaotic system ; in a regular system one has P (t) / (t₀=t) ², since $_0^2$ (t) dom – inates the decay of P (t) independently of the correlators in Eqs.(8-9); and (iii) a saturation regime where the purity reaches its m inim al value. In the chaotic limit, this saturation value can be estimated using a RM T approach as P (1) = 2 (^d=) + 0 (²) [22], which is in qualitative agreement with the results obtained in Ref.[5] for the von N eum ann entropy. There is no reason to expect a universal saturation value in the regular regime.

Analyzing these results, we note that Eqs. (6) and (7) are rem iniscent of the results obtained for P (t) by perturbative treatments in Refs. [6, 7], but they apply well beyond the linear response regime. Our weak coupling condition that the one-particle actions S vary faster than the two-particle actions S roughly gives an upper bound U E for the typical interaction strength U (E is the one-particle energy). The linear response regime is how ever restricted by a much more stringent condition IJ E (is the mean level spacing) [16]. The decay regime (ii) of P (t) reconciles the a priori contradicting claim sofR efs. [3, 4, 7] and R ef. [6]. Forweak coupling, the decay of P (t) is given by classical correlators, and thus depends on the interaction strength, in agreement with Ref. [6]. However, P (t) cannot decay faster than the bound given in Eq. (5), so that at stronger coupling, and in the chaotic regime, one recovers the results of Ref. [4]. Simultaneously, regime (ii) also explains the data in Fig. 2 and 4 of Ref. [7], showing an exponential decay of P (t) in the chaotic regime, and a power-law decay with an exponent close to 2 in the regular regime (this power-law decay was left unexplained by the authors of Ref. [7]). Our sem iclassical treatment thus presents a uni ed picture of the problem.

Three more remarks are in order here. First the stationary phase solutions leading to the above results still. hold in the case when the two particles are subjected to di erent one-particle Ham iltonians. Second, the powerlaw decay of P (t) predicted above for regular system s, is to be taken as an average over initial conditions $r_{1:2}$ (in that respect see Refs. [14] and [24]), but may also hold for individual initial conditions, as e.g. in [7]. Finally, there are cases when the correlators (8) and (9) decay exponentially in time with a rate related to the spectrum of Lyapunov exponents. This also may induce a dependence of P (t) on the Lyapunov exponents, which can be captured by the linear response approach of Ref. [6]. We note how ever that this is not a generic situation, as most fully chaotic, but nonhyperbolic system s have power-law decaying correlations.

As a concluding line, and noting similarities between the problem treated here and that of decoherence by an environment [10], we anticipate that a sem iclassical approach as the one presented here (see also R ef. [25]) could clarify how decoherence relates to the intrinsic dynamics of the system [10, 26].

This work has been supported by the Swiss National Science Fondation. We thank Atushi Tanaka for a clarifying discussion of Refs. [4, 6].

- E. Schrodinger, Proc. Cambridge Phil. Soc. 31, 555 (1936).
- [2] M A. Nielsen and IL. Chuang, Quantum Computation and Quantum Information, Cambridge University Press, Cambridge (2000).
- [3] K.Furuya, M L.Nem es, and G Q. Pellegrino, Phys. Rev. Lett. 80, 5524 (1998).
- [4] P A .M iller and S.Sarkar, Phys. Rev. E 60, 1542 (1999).
- [5] A. Lakshm inarayan, Phys. Rev. E 64, 036207 (2001); JN. Bandyopadhyay and A. Lakshm inarayan, quantph/0307134.
- [6] A. Tanaka, H. Fujisaki, and T. M iyadera, Phys. Rev. E 66,045201(R) (2002); H. Fujisaki, T. M iyadera, and A. Tanaka, Phys. Rev. E 67,066201 (2003).
- [7] M. Znidaric and T. Prosen, J. Phys. A 36, 2463 (2003).
- [8] A J. Scott and C M . Caves, J. Phys. A 36, 9553 (2003).
- [9] Care should however be taken when comparing fully chaotic systems with regular ones, as there is no universality to be expected in the latter case.
- [10] W H.Zurek, Rev.M od. Phys. 75, 715 (2003).
- [11] J.S. Bell, Speakable and Unspeakable in Quantum Mechanics, Cambridge University Press, Cambridge (1987).
- [12] Yu Shi, Phys. Rev. A 67, 024301 (2003).
- [13] R A. Jalabert and H M. Pastawski, Phys. Rev. Lett. 86, 2490 (2001).
- [14] Ph.Jacquod, I.A dagideli, and C W J.Beenakker, Europhys.Lett. 61, 729 (2003).

- [15] N R. Cernuti and S. Tom sovic, Phys. Rev. Lett. 88, 054103 (2002); J. Vaniœk and E.J. Heller, Phys. Rev. E 67, 016211 (2003).
- [16] Ph. Jacquod, P.G. Silvestrov, and C.W. J. Beenakker, Phys. Rev. E 64, 055203 (R) (2001).
- [17] N R. Cerruti and S. Tom sovic, J. M ath. Phys. 36, 3451 (2003).
- [18] A M. O zorio de A lm eida, H am iltonian System s: Chaos and Quantization, C am bridge (1988).
- [19] G P. Berm an and G M. Zaslavsky, Physica A 91, 450 (1978); M V. Berry and N L. Balasz, J. Phys. A 12, 625 (1979).
- [20] Because of the second line in Eq.(6), the connection proposed in Ref. [21] between decoherence and Loschmidt E cho breaks down at short times.
- [21] F.M. Cucchietti, D.A.R. Dalvit, J.P. Paz, and W.H. Zurek, Phys. Rev. Lett. 91, 210403 (2003).
- [22] Ph. Jacquod, unpublished (2003).
- [23] If U factorizes as U = V I + I W, $S_{s;s^0}(y_1;x;y_2;r;t) = S_s^{(U)}(y_1;x;t) + S_{s^0}^{(W)}(y_2;r;t)$, the two-particle action thus vanishes and no entanglem ent is generated, as should be.
- [24] T. Prosen and M. Znidaric, New Jour. Phys. 5, 109 (2003).
- [25] G A. Fiete and E J. Heller, Phys. Rev. A 68, 022112 (2003).
- [26] W H. Zurek, Physics Today 44, 36 (1991); A K. Pattanayak, Phys. Rev. Lett. 83, 4526 (1999).