

Hydrogen atom in strong magnetic field revisited

I.B. Khriplovich and G.Yu. Ruban
Budker Institute of Nuclear Physics,
630090 Novosibirsk, Russia,
and Novosibirsk University

Abstract

We derive in a straightforward way the spectrum of a hydrogen atom in a strong magnetic field.

The spectrum of hydrogen atom in a strong magnetic field was found long ago [1], and is presented now in textbooks (see, e.g., [2] (x112, Problem 3)). The approach used therein is as follows. At first, using the wave functions of electron in a magnetic field, one constructs an effective potential for the motion along the field, and then the spectrum in this potential is found. We present here a somewhat different solution of the problem, with the results coinciding in fact with those of [1, 2]. We hope however that our approach, physically straightforward and transparent, is of some interest. At least, the present note may be considered as a sort of mini-review on the subject.

Our starting point is the obvious observation that in a sufficiently strong magnetic field H (the exact criteria are discussed below) the motion of an atomic electron becomes almost one-dimensional, along the magnetic field, in the Coulomb potential $\varphi^2 = z$.

The corresponding one-dimensional wave equation is

$$u'' + \left(\frac{1}{4} + \frac{z}{a} \right) u = 0; \quad (1)$$

We have introduced in it the usual dimensionless variable:

$$\frac{2z}{a} \equiv z;$$

here $a = \hbar^2 / m_e e^2$ is the Bohr radius, m_e is the electron mass, ℓ is the effective quantum number, related to the electron energy as

$$E = - \frac{m_e e^4}{2 \hbar^2 \ell^2}; \quad (2)$$

Equation (1) coincides exactly with the radial equation for the s-wave in the three-dimensional Coulomb potential $\varphi^2 = r$, and has therefore the common hydrogen spectrum

$$E_n = - \frac{m_e e^4}{2 \hbar^2 n^2} \quad n = 1; 2; 3; \dots; \quad (3)$$

and the set of solutions

$$u_n(z) = \exp(-z/2) z F(-n; 2; z); \quad n = 1; 2; 3; \dots; \quad (4)$$

where F is the confluent hypergeometric function (here and below we are not interested in the normalization factors). These solutions vanish at the origin and are trivially continued to $z < 0$. Thus obtained solutions on the whole z axis are odd under $z \rightarrow -z$ (as reflected by the superscripts "\minus" in (3), (4)).

There is however an essential difference between the present problem and the s-wave Coulomb one. In the last case (4) is the only solution. The reason is well-known. Naively the radial wave equation for $R(r) = u(r)/r$ has two independent solutions, which behave for $r \rightarrow 0$ as $R = \text{const}$ ($u \sim r$) and $R \sim 1/r$ ($u = \text{const}$), respectively. However, in fact $R \sim 1/r$ is no solution at all for the homogeneous wave equation if the point $r = 0$ is included, since $\Delta(1/r) = -\delta(r)$. As to our problem, equation (1) does not describe really the vicinity of $z = 0$ since therein we have to consider seriously the magnetic field itself. Therefore, there are no reasons to discard those solutions of (1) which tend to a constant for small z (and of course decrease exponentially for $z \rightarrow \pm\infty$).

Such solutions are presented in a convenient form in [2] (Mathematical Appendices, x d, (d.17)). To our purpose they can be written for $z > 0$ as

$$u^+(z) = \exp(-z/2) \left[1 - z \ln z F(1; 2; z) + \sum_{k=0}^{\infty} \frac{(1+k) [(1+k)(k+2)(k+1)]^k}{(k!)^2 (k+1)!} z^k \right]; \quad (5)$$

here $\psi(x)$ denotes the logarithmic derivative of the gamma function: $\psi(x) = \Gamma'(x)/\Gamma(x)$. Being trivially continued to $z < 0$, thus obtained solutions on the whole z axis are even under $z \rightarrow -z$ (as reflected by the superscript "\plus" in (5) and in the corresponding eigenvalues below).

Under any reasonable regularization of the logarithmic singularity at $z \rightarrow 0$, the even solutions should have vanishing first derivative at the origin. In this way we obtain the following equation for the eigenvalues of (1):

$$\ln \frac{a}{a_H} = \frac{1}{2} + \psi\left(\frac{1}{2}\right); \quad (6)$$

here $a_H = \frac{q}{hc} = \frac{e}{H}$ is the typical scale for the radius of electron orbits in the magnetic field H . We are working in the logarithmic approximation, i.e. assume that

$$a = a_H e^{-1}; \quad (7)$$

This allows us to use a crude cut-off at a_H for the formal logarithmic divergence at $z \rightarrow 0$, as well as to simplify somewhat this equation.

The smallest root of equation (6) is

$$\frac{1}{2} = \frac{1}{2}; \quad (8)$$

which gives the ground state energy

$$E_0^+ = \frac{m_e e^4}{2\hbar^2} \ln^2 \frac{\hbar^3 H^2}{m_e^2 e^3 c}; \quad (9)$$

Other roots of equation (6) are

$$n^+ = n + \frac{1}{2} \quad n = 1; 2; 3; \dots; \quad (10)$$

with the corresponding energies

$$E_n = \frac{m_e e^4}{2\hbar^2 n^2} \left(1 - \frac{2}{n} \ln \left(1 + \frac{\hbar^3 H^2}{m_e^2 e^3 c} \right) \right) \quad (11)$$

Let us mention that the one-dimensional Coulomb problem was considered in [3-5] with various regularizations of the singularity at $z \rightarrow 0$, but without any relation to the problem of the hydrogen atom in a strong magnetic field.

But let us come back to our problem. The resulting spectrum of the hydrogen atom in a strong magnetic field looks as follows. Each Landau level in this field serves as an upper limit to the sequence of discrete levels of the Coulomb problem in the z direction. This discrete spectrum consists of a singlet ground state with the energy given by formula (9), and close doublets of odd and even states of energies given by formulae (3) and (11). There is also a continuous spectrum of the motion along z above each Landau level.

This picture is valid for sufficiently low Landau levels, as long as the radius of a magnetic orbit is much less than the Bohr radius. Obviously, in a strong magnetic field this description fails for large magnetic quantum numbers, i.e., in the semiclassical region. Here we can estimate the orbit radius directly from the well-known spectrum of electron in a magnetic field (see, e.g., [2] (x112, Problem 1)):

$$E = \hbar \frac{eH}{m_e c} (N + 1/2); \quad N = n + \frac{m + |m|}{2}; \quad (12)$$

here n is the radial quantum number in the xy plane, and m is the angular momentum projection onto the z axis. Now the semiclassical estimate for the magnetic radius is

$$a_H(N) \approx \frac{\hbar c}{eH} N = a_H \sqrt{N} \quad (13)$$

Thus the present picture of levels holds as long as

$$a_H \sqrt{N} \ll a \quad (13)$$

At last, let us consider the correspondence between the obtained system of levels in a strong magnetic field and the hydrogen spectrum in a vanishing field. A beautiful solution of this problem was given in [6] (and is quoted in [1]). We would like to present the solution here as well (having in mind in particular that our note can be considered as a mini-review).

The crucial observation made in [6] is as follows. While changing the magnetic field from vanishingly small to a very strong one, the number of nodal surfaces of a given wave function remains the same. A hydrogen wave function (in zero magnetic field) with quantum numbers n, l, m has $n - l - 1$ nodal spheres and $l - |m|$ nodal cones with a z axis. With the increase of the magnetic field, the nodal spheres become ellipsoids of rotation, more and more prolate, tending to cylinders in the limit of infinite field. The

correspondence between n_r and n (n being the radial quantum number in the xy plane in a strong magnetic field) gets obvious from this picture:

$$n = n_r: \quad (14)$$

The evolution of the hydrogen nodal cones is less obvious. However, due both to equation (14) and to the conservation of the total number of nodal surfaces, the number n_z of the nodes of a solution of equation (1) should coincide with the number of nodes of the corresponding spherical function,

$$n_z = 1 + j + j: \quad (15)$$

In other words, $1 + j + j$ nodal cones of a hydrogen wave function evolve into n_z planes of constant z corresponding to the nodes of an eigenfunction of equation (14).

And at last, let us note that during the whole evolution of the magnetic field, m remains constant.

Let us consider now, for instance, the ground state in the magnetic field, with $N = 0$ (see (12)). Obviously, it is degenerate, and its corresponding magnetic wave functions have $n = 0$ and $m = 0; \pm 1; \pm 2 \dots$. Let us confine further to its lowest sublevel corresponding to the ground state solution of equation (1), with $n_z = 0$. According to the above arguments, the hydrogen ancestors of those wave functions should have $n_r = 0$ and $l = j + j$. In other words, these ancestors are:

$$1s; 2p; m = \pm 1; 3d; m = \pm 2; \text{ and so on:}$$

We are grateful to V.A. Novikov, V.V. Sokolov, M.I. Vysotsky, and A.V. Zolotarev for useful discussions. We acknowledge the support by the Russian Foundation for Basic Research through grant No. 03-02-17612.

References

- [1] R.J. Elliott and R. Loudon, J. Phys. Phys. Chem. Solids. 15 (1960) 196
- [2] L.D. Landau and E.M. Lifshitz, Quantum Mechanics (1989, Nauka, Moscow)
- [3] R. Loudon, Amer. J. Phys. 27 (1959) 649
- [4] M. Andrews, Amer. J. Phys. 34 (1966) 1194
- [5] L.K. Haines and D.H. Roberts, Amer. J. Phys. 37 (1969) 1145
- [6] W.H. Kerner, Lincoln Lab. Progr. Rep. (Feb. 1960)