

Stability of Quantum Motion: Beyond Fermi-golden-rule and Lyapunov decay

Wen-ge Wang^{1,2}, G. Casati^{3,4,1}, and Baowen Li¹

¹Department of Physics, National University of Singapore, 117542 Singapore

²Department of Physics, Southeast University, Nanjing 210096, China

³Center for Nonlinear and Complex Systems, Università degli Studi dell'Insubria and Istituto Nazionale per la Fisica della Materia,

Unità di Como, Via Valleggio 11, 22100 Como, Italy

⁴Istituto Nazionale di Fisica Nucleare, Sezione di Milano, Via Celoria 16, 20133 Milano, Italy

(dated: 10 September 2003)

We study, analytically and numerically, the stability of quantum motion for a classically chaotic system. We show the existence of different regimes of delity decay. In particular, when the underlying classical dynamics is weakly chaotic, deviations from Fermi-golden-rule and Lyapunov regimes are observed and discussed.

PACS numbers: 05.45.Mt, 05.45.Pq, 03.65.Sq

The nature of correlations decay is an important subject in different fields of physics. In particular, after the discovery of the so-called dynamical chaos, a large effort has been devoted to understand their behavior in relation to dynamical properties. The main reason is to know the precise conditions under which a statistical description is legitimate and to estimate the nature of the approximations which are involved.

Another important characteristic of dynamical systems is the stability of their solutions under slight variation of the Hamiltonian. A quantitative measure of this stability is given by the so-called delity or quantum Loschmidt echo. The delity $M(t) = |\langle \psi(t) | \psi(0) \rangle|^2$, measures the overlap of two states started from the same initial state and evolved under slightly different Hamiltonians H_0 and $H = H_0 + V$, which are classically chaotic,

$$m(t) = \langle \psi_0 | \exp(iH_0 t) \exp(-iH t) | \psi_0 \rangle \quad (1)$$

Quite surprisingly, in spite of its physical relevance, the behavior of delity has been scarcely considered and only recently, in connection with quantum computation, a large number of papers appeared. Some important features of delity are now understood even though we are still far from the detailed level of knowledge we have about related quantities such as correlations functions and escape probabilities. So far, above the perturbative regime of small V , with Gaussian type decay [1, 2, 3], two main types of exponential decay of the delity have been identified: i)–The Fermi-golden-rule (FGR) decay, with the exponent given by the half-width of the corresponding local spectral density of states [3, 4, 5, 6]. This decay has been related to the decay of autocorrelation function [9]. ii)–The Lyapunov regime, above the FGR regime, with decay rate given by the Lyapunov exponent of the underlying classical dynamics [4, 7, 8, 9, 10, 11, 12, 13, 14, 15].

In this paper we show that for classically chaotic systems, in particular those with weak chaos, the behavior of delity can be much more rich and complex than




FIG. 1: Fidelity $\overline{M}(t)$ as a function of $\sigma^2 t$ for $K_0 = 0.4, 0.1, 0.2, 0.4$ (from bottom to top) where $N_0 = 4096$. ($\sigma = 0.05; 0.1; 0.2; 0.4$). The FGR decay $e^{-2\lambda^2 t}$ is shown by the dashed line. Full circles represent the semiclassical values $\overline{M}_a(t)$ at $\sigma = 0.4$, computed with expression (7). The numerically computed semiclassical values $\overline{M}_f(t)$ turn out to be negligible so that $\overline{M}(t)$ is well approximated by $\overline{M}_a(t)$, as clearly seen from the figure at $\sigma = 0.4$. Averages were performed over 400 initial point sources, with ψ_0 taken randomly in the interval $[0; 2\pi)$. (The same decaying behaviors are observed for initial Gaussian wavepackets.)

expected. In particular we study perturbation borders which separate different types of decay.

We start by displaying numerical results which strongly deviate from the expected behavior. We consider here the simple, well known, sawtooth map model [12]. The classical map writes:

$$\overline{p} = p + K_0 \left(\frac{p}{2} \right); \quad \overline{p} = \overline{p} + \overline{p} \pmod{2} \quad (2)$$

For $K_0 > 0$, the motion is completely chaotic, with Lyapunov exponent $\lambda = \ln(2 + K_0 + [(2 + K_0)^2 - 4]^{1/2}) = 2\gamma$.




FIG. 2: The exponential decay rate versus perturbation strength σ , calculated from the best fit of $\ln \overline{M}(t)$. Gaussian wavepackets are taken as initial states. The solid curve shows the rate $\gamma = 2\sigma^2$ of the FGR-decay. The dashed horizontal lines correspond to the Lyapunov exponents $\lambda = 0.62; 0.96$ and 1.32 for $K_0 = 0.4; 1$ and 2 , respectively. $N = 131072$.

The quantum evolution on one map iteration is described by

$$\hat{U} = U_0 \exp(i\hat{p}^2/(2\hbar)) \exp[ik_0(\hat{x})^2/2]; \quad (3)$$

where $\hat{p} = i\hbar\partial/\partial x$ and $k_0 = K_0/\hbar$, with the effective Planck constant $\hbar = 2/N$ and N being the dimension of the Hilbert space. For the perturbed system, $K = K_0 + \delta K$ and $k = k_0 + \delta k$, where $\delta K \ll K_0$ and $\delta k \ll k_0$.

In Fig. 1 we show the delity decay in the expected FGR regime $\delta K \ll K_0$. In spite of the fact that the classical motion is chaotic, it is clearly seen that the behavior does not obey the FGR which, according to [4, 12], should be $\sim \exp(-\gamma t)$ with $\gamma = 2\sigma^2$. The same conclusion can be drawn from Fig. 2 where we plot the decay rate of the delity as a function of σ . Indeed at $K_0 = 0.4$ the decay rate versus σ appears quite different from the quadratic one [16].

Deviations are present even at $K_0 = 1$ and only at $K_0 = 2$ one has good FGR decay. Moreover, above the FGR regime, where one expects Lyapunov decay, there are strong oscillations above and below the decay rate (for $K_0 = 1$, and 2). Only at larger σ values, one enters the Lyapunov regime.

In order to explain the above numerical results, we start from the standard semiclassical approach [7, 14]. For simplicity, we consider a finite configuration space, with dimension d and volume $V = \int dr$. The momentum space is also finite, with a volume V_p . In the semiclassical approach, an initial state $\psi_0(r_0)$ is propagated by the semiclassical Van Vleck-Gutzwiller propagator, $\psi(r;t) = \int dr_0 K_{sc}(r;r_0;t) \psi_0(r_0)$, where $K_{sc}(r;r_0;t) =$

$\int_{s_0}^s K_s(r;r_0;t)$, with

$$K_s(r;r_0;t) = \frac{C_s^{1/2}}{(2\pi i\hbar)^{d/2}} \exp\left\{ \frac{i}{\hbar} S_s(r;r_0;t) - \frac{i}{2} \mu_s \right\}; \quad (4)$$

The labels s in Eq. (4) (more exactly $s(r;r_0;t)$), indicate classical trajectories starting at r_0 and ending at r in a time t ; $S_s(r;r_0;t)$ is the time integral of the Lagrangian along the trajectory s , $S_s(r;r_0;t) = \int_0^t dt L$, $C_s = \sqrt{|\det(\partial^2 S_s / \partial r_0 i \partial r_j)|}$, and μ_s is the Maslov index counting the conjugate points.

In Ref. [14], it is shown that the semiclassical approximation to $m(t)$ for initial Gaussian wavepackets has a simple and convenient expression in the initial momentum space. Following similar arguments for initial point sources, $\psi_0(r) = \delta(r-r_0)$, (the theory can be extended to general initial states), one can write $m(t)$ as

$$m(r_0;t) \sim \int_{V_p} \frac{1}{V_p} dp_0 \exp\left\{ \frac{i}{\hbar} S(p_0;r_0;t) \right\}; \quad (5)$$

where $S(p_0;r_0;t)$ is the action difference along the trajectory starting at $(r_0;p_0)$ for the two systems H_0 and H . In the first order classical perturbation theory, $S(p_0;r_0;t) = \int_0^t dt V[r(t^0)]$, with V evaluated along the trajectory.

The averaged (over r_0) delity can be separated into a mean-value part and a fluctuating part [1], denoted by $\overline{M}_a(t)$ and $\overline{M}_f(t)$ respectively, $\overline{M}(t) = \overline{M}_a(t) + \overline{M}_f(t)$, where

$$\overline{M}_a(t) = \overline{|\overline{m}(t)|^2}; \quad \text{with } \overline{m}(t) = \frac{1}{V} \int dr_0 m(r_0;t); \quad (6)$$

From Eqs. (5) and (6), it is seen that the mean-value part $\overline{M}_a(t)$ can be expressed in terms of the distribution $P(S)$ of the action difference S ,

$$\overline{M}_a(t) \sim \int dS e^{iS/\hbar} P(S); \quad \text{where} \quad (7)$$

$$P(S) = \frac{1}{V} \int dr_0 dp_0 \delta[S - S(p_0;r_0;t)]; \quad (8)$$

It is usually assumed that for chaotic systems $P(S)$ is close to a Gaussian with a variance $[2^2 K(E)t]$, where $K(E) = \int_0^1 dt |V[r(t)]| V[r(0)]$ is the classical action diffusion constant [3]. As a result, $\overline{M}_a(t) \sim e^{-t}$, where $t = 2^2 K(E)$. At small t , the fluctuation is small compared with the average value, because the phase on the right hand side of Eq. (5) is proportional to S ; then, $\overline{M}(t) \sim \overline{M}_a(t)$ has the FGR-decay.

Let us now consider a fixed \hbar , and divide the space of the initial momenta p_0 into connected, disjoint subspaces, denoted by A , where each A is the largest possible subspace such that the correspondence between p_0 and the final position r is one-to-one, i.e. different p_0




FIG. 3: Comparison between the exact $\overline{M}(t)$, its semiclassical mean-value part \overline{M}_a , calculated by using Eqs. (5) and (6), and the fluctuation part $\overline{M}_f = \overline{M}_{sc} \overline{M}_a$, where \overline{M}_{sc} is the semiclassical approximation to the density, computed by the expression (5). Here $N = 131072$, $K_0 = 2$, and $\sigma = 0.9$ (left panel), $\sigma = 3$ (right panel). The exact $\overline{M}(t)$ is in good agreement with its semiclassical approximation $\overline{M}_{sc}(t)$. The average is taken over 500 initial point sources.

inside each single component A_j gives different initial position r . It is always possible to make such a division. The number of subspaces A_j is denoted by N_j . Note also that the sizes of A_j decrease exponentially with increasing time t . When p_0 runs over a subspace A_j , r may run over part of the configuration space, denoted by V_j . Note that, with this division of the p_0 subspace, the trajectories starting at r_0 are divided into N_j groups and "near" trajectories typically belong to the same group.

The amplitude $m(r_0; t)$ in Eq. (5) can now be written as $m(r_0; t) = \overline{m}(r_0; t)$, where

$$\overline{m}(r_0; t) = \frac{1}{V_p} \int_V dr C_s \exp \left(\frac{i}{\hbar} S_s(r; r_0; t) \right) \quad (9)$$

with integration over the subspace V_j , in which the change of variable $p_0 \rightarrow r$ within the subspaces A_j has been done and $S_s(r; r_0; t)$ coincides with $S(p_0; r_0; t)$ for the same trajectories starting at $(r_0; p_0)$ with $p_0 \in A_j$. $\overline{M}_f(t)$ is written as

$$\overline{M}_f(t) = \sum_j \overline{m}_f^2 \quad \text{with } \overline{m}_f = \overline{m}(r_0; t) \quad \frac{\overline{m}(t)}{N} \quad (10)$$

When σ is large enough, above a critical border σ_c , $m(r_0; t)$ can be regarded as possessing random phase, and therefore \overline{M}_f can be approximated by its diagonal




FIG. 4: Similar to Fig. 3, for $K_0 = 1$ and $\sigma = 6$. At this large σ , $\overline{M}_a(t)$ is negligible compared with $\overline{M}_f(t)$.

part

$$\overline{M}_f(t) = \sum_j \overline{m}_f^2 = \sum_j \int_{V_j} dr C_s \exp \left(\frac{i}{\hbar} S_s(r; r_0; t) \right)^2 \quad (11)$$

where the second approximation is obtained by noticing that $\overline{m}(t) = N^{-1} \sum_j m_j$ at large σ .

When the phase space is homogeneous with constant local (maximum) Lyapunov exponent λ , as in the sawtooth map, the number $N_s(r_0; r)$ of trajectories connecting two points r_0 and r in the configuration space is approximately $N_s \sim N e^{-\lambda t}$ [17]. The summation over j in (11) gives a contribution approximately proportional to N_s . At t large enough, the main time dependence of $\int_V dr C_s e^{\frac{i}{\hbar} S_s} j$ is given by $C_s / e^{-\lambda t}$. Combining these results, it is seen that at $\sigma > \sigma_c$, $\overline{M}_f(t)$ has the Lyapunov decay, $\overline{M}_f(t) \sim e^{-\lambda t}$.

In order to have the Lyapunov decay for $\overline{M}(t)$, the term $\overline{M}_a(t)$ must be small. To this end one needs to further increase σ above a critical value σ_c , so that the variance of the phase of $m(r_0; t)$ with respect to r_0 will become so large that $\overline{M}_a(t)$ is negligible.

The right panel of Fig. 3 gives an example of $\overline{M}_a(t)$ and $\overline{M}_f(t)$. This explains the fluctuation versus σ shown in Fig. 2 at $K_0 = 2$ and $\sigma < 3$. Fig. 4 instead gives an example with σ large enough ($\sigma > \sigma_c$), so that $\overline{M}_a(t)$ is negligible and $\overline{M}(t) \sim \overline{M}_f(t)$.

The deviation from FGR decay observed in Figs. 1 and 2 is due to the deviation of $P(S)$ from the Gaussian behavior. Indeed, when chaos in the underlying classical dynamics is strong enough ($K_0 > 1$), correlations be-




FIG. 5: Distribution $P[(S - \langle S \rangle)/\epsilon]$ of the classical action difference S , at $t = 10$, calculated by taking randomly 10^7 initial points in the phase space, where $h S_i = \text{thV}(i) = 2t = 6$, with an average over the phase space.

tween non-overlapping parts of a trajectory decay very rapidly and the distribution $P(S = \dots)$ reaches, in a relatively short time, the Gaussian distribution. This is the case of Fig. 5 for $K_0 = 2$, where $K(E) = 4 = 90 \cdot 1.08$ with $\epsilon = 2K(E)^2 \cdot 2 \cdot 16^2$, in agreement with the numerical results in Ref. [12] and in Fig. 2. However, when K_0 is not sufficiently large, e.g., $K_0 = 0.4$, a considerable deviation of $P(S = \dots)$ from the Gaussian distribution appears for times comparable to the delity decay times (Fig. 5). According to Eq. (7), this leads to deviations from the FGR-decay as observed in Fig. 1. We would like to draw the reader attention to the fact that, for $K_0 < 1$ the saw-tooth map, even though completely chaotic, it possess a structure of cantori which, in the quantum case, can act as perfect barriers to quantum motion thus leading to localization of wavefunctions.

Notice that the deviation of $P(S = \dots)$ from the Gaussian distribution depends on K_0 but not on ϵ . Therefore, by increasing ϵ , the effect of this deviation becomes more and more important, since the FGR exponential decay has a decay rate proportional to ϵ^2 while the deviation from the Gaussian remains unchanged. Therefore, for a given system, there is a critical value ϵ_d , below which the FGR decay is obeyed with good accuracy and above which FGR breaks down. This case is illustrated in Fig. 2 for the case $K_0 = 1$, which coincides with the well-known Arnold cat map, the paradigmatic model of chaos. Here the distribution $P(S = \dots)$ (at $t = 10$) is slightly different from the Gaussian distribution and the decay rate of delity deviates from the FGR decay for $\epsilon > 0.3$. In cases of weak classical chaos, the value of ϵ_d can be so small that FGR is never observed (e.g. the case with $K_0 = 0.4$). The left panel in Fig. 3 shows instead a case at $K_0 = 2$ and $\epsilon = 0.9$, in which $\overline{M}_a(t)$ obeys the FGR

decay and $\overline{M}_f(t)$ is negligible.

To summarize: above the perturbative border, the delity has a FGR decay for $\epsilon < \epsilon_d$, while for $\epsilon > \epsilon_d$, it has the Lyapunov decay. In the intermediate region, for $\epsilon_d < \epsilon < \epsilon_f$, the delity deviates from FGR and can decay even faster than Lyapunov. For $\epsilon_f < \epsilon < \epsilon_r$, $\overline{M}_a(t) \sim \overline{M}_f(t)$ and the decay rate of $\overline{M}(t)$ fluctuates around the Lyapunov exponent. It may be useful to recall here the physical meaning of different borders. Above ϵ_d , the distribution $P(S)$ deviates from the Gaussian and this induces deviations from the expected FGR decay. Below ϵ_r , $\overline{M}_a(t)$ is non negligible as compared to $\overline{M}_f(t)$ and this induces deviations from the expected Lyapunov decay.

It may be interesting to remark that the relation between the decomposition in two parts of $\overline{M}(t)$ here and that in Ref. [7] is the following. At small enough $\overline{M}(t) \sim \overline{M}_a(t) \sim \overline{M}^{nd}(t)$, with $\overline{M}_f(t)$ and $\overline{M}^d(t)$ negligible; while at large enough $\overline{M}(t) \sim \overline{M}_f(t) \sim \overline{M}^d(t)$, with $\overline{M}_a(t)$ and $\overline{M}^{nd}(t)$ negligible. In the intermediate regime of ϵ , in particular, in the crossover from the FGR decay to the Lyapunov decay, there may be considerable difference between the two divisions.

In this paper, by using the sawtooth map, we have demonstrated that the delity decay in a generic chaotic system can have a very complex behavior. In particular, deviations from the Fermi golden rule (for weak chaos) and Lyapunov decay have been discussed as well as the existence of perturbation borders separating different regimes. It is our opinion that delity is an important quantity which characterizes the stability of classical and quantum systems. It therefore deserves deeper analytical and numerical studies in order to fully understand its behavior in different dynamical regimes.

The authors are grateful to V. Sokolov for valuable discussions. This work was supported in part by the Academic Research Fund of the National University of Singapore. Support was also given by the EC RTN contract HPRN-CT-2000-0156, the NSA and ARDA under ARO contracts No. DAAD 19-02-1-0086, the project ED IQ IP of the IST-FET programme of the EC, the PRIN-2002 "Fault tolerance, control and stability in quantum information processing", and the Natural Science Foundation of China No.10275011.

-
- [1] A. Peres, Phys. Rev. A 30, 1610 (1984).
 - [2] T. Prosen, Phys. Rev. E 65, 036208 (2002); T. Prosen and M. Žnidarič, J. Phys. A 35, 1455 (2002).
 - [3] N. R. Cernuti and S. Tomsović, Phys. Rev. Lett. 88, 054103 (2002); J. Phys. A 36, 3451 (2003).
 - [4] Ph. Jacquod, P.G. Silvestrov, and C.W.J. Beenakker, Phys. Rev. E 64, 055203 (2001).
 - [5] Ph. Jacquod, I. Adagideli, and C.W.J. Beenakker, Phys. Rev. Lett. 89, 154103 (2002).

- [6] J. Emerson et al, Phys. Rev. Lett. 89, 284102 (2002).
- [7] R. A. Jalabert and H. M. Pastawski, Phys. Rev. Lett. 86, 2490 (2001).
- [8] F. M. Cucchietti, H. M. Pastawski, and D. A. Wisniacki, Phys. Rev. E 65, 045206 (2002);
- [9] D. A. Wisniacki and D. Cohen, Phys. Rev. E 66, 046209 (2002).
- [10] D. A. Wisniacki et al, Phys. Rev. E 65, 055206 (2002);
- [11] F. M. Cucchietti et al, Phys. Rev. E 65, 046209 (2002).
- [12] G. Benenti and G. Casati, Phys. Rev. E 65, 066205 (2002); F. Borgonovi, G. Casati, and B. Li, Phys. Rev. Lett. 77, 4744 (1996).
- [13] W. G. Wang and B. Li, Phys. Rev. E 66, 056208 (2002).
- [14] J. Van oek and E. Heller, Phys. Rev. E 68, 056208 (2003).
- [15] F. M. Cucchietti et al, quant-ph/0307752.
- [16] Deviations from Fermi golden rule have also been observed in the Bunimovich stadium billiard [10]. The delity in that model was found to decay as $\exp(-t/2)$, with being the halfwidth of the LDOS. In our case the decay rate of delity is close to . However, for $K_0 = 0.4$ considerable deviations in tail region of LDOS from the Lorentzian shape are present.
- [17] F. Haake, Quantum Signatures of Chaos, 2nd ed. (Springer-Verlag, Berlin, 2001).