M in in um decoherence cat-like states in Gaussian noisy channels

A. Sera nil, S. De Sienal, F. Illum inatil, and M. G. A. Paris²

Dipartimento di Fisica \E.R.Caianiello", Universita di Salemo, INFM UdR Salemo, INFN Sez. Napoli, G.C.Salemo, Via S.Allende, 84081 Baronissi (SA), Italy
²I.S.I.S.\A.Sorbelli", via G.M. atteotti 2, I-41026 Pavullo nel Frignano (MO), Italy

September 29, 2003

A bstract

We address the evolution of cat-like states in general Gaussian noisy channels, by considering superpositions of coherent and squeezed-coherent states coupled to an arbitrarily squeezed bath. The phase space dynamics is solved and decoherence is studied keeping track of the purity of the evolving state. The in uence of the choice of the state and channel parameters on purity is discussed and optimal working regimes that minimize the decoherence rate are determined. In particular, we show that squeezing the bath to protect a non squeezed cat state against decoherence is equivalent to orthogonally squeezing the initial cat state while letting the bath be phase insensitive.

1 Introduction

In Schrodinger original paper [1], a bipartite entangled state of the form

where A i and B i denote the \alive" and \dead" states of a cat and j i two orthogonal states of a microscopic system, was suggested to illustrate the counterintuitive consequences of quantum mechanics in a macroscopic setting. More generally, in the literature, any single-system coherent superposition j i / j i+ j + i of two pure quantum states j i that are mesoscopically distinguishable, is often referred to as a Schrodinger cat or a cat-like state. The interest in the study of such superpositions, possibly involving states of a microscopic system as well, stems from both theoretical and experimental considerations. A ctually, a cat state is one of the simplest and most fundamental con gurations allowing to probe the archetypal aspects of quantum theory: the superposition principle and quantum entanglement.

As far as quantum optical systems are concerned, the possibility of realizing superposition states of the radiation eld, rst envisioned by Yurke and Stoler [2], has been extensively investigated in later years [3, 4]. However, pure state superpositions are in general corrupted by the interaction with the environment. Therefore, cat-like states that are available for experiments are usually mixed states that su ered a partial decoherence, and it is crucial to know whether and to which extent superpositions can survive the environmental noise. The them e of decoherence of cat{like states spurred relevant theoretical works [5, 6, 7, 8], especially aimed to select schemes of quantum control and feedback stabilizing coherent superpositions

against decoherence [7]. Recent promising experimental results and perspectives continue to keep a widespread interest in this subject [9, 10].

In the present paper, we study the non-unitary evolution of coherent and squeezed-coherent single-m ode Schrodinger cat-like states in generic Gaussian noisy channels, namely, either therm all or squeezed-therm all baths of harm onic oscillators [11, 12]. In particular, we will focus our attention on the evolution of the purity (or linear entropy) of the states, showing how the quantum superposition is corrupted by the interaction with a noisy environment and how to optim ize the state and channel parameters to minimize the decoherence rate.

The paper is structured as follows. In Section 2 we introduce notations and evaluate the W igner function of generalized cat-like states. In Section 3 the time evolution in G aussian noisy channels is studied, whereas in Section 4 the dependence of decoherence and purity on the parameters of the initial state and of the channel is discussed in detail. In Section 5 optm ized regimes to minimize decoherence are discussed and, nally, Section 6 provides some concluding remarks.

2 Cat-like states

The simplest example of a Schrodinger cat state of a single-mode radiation eld is the following normalized superposition of coherent states

$$j_0; i = \frac{j_0 i + e^i j_0 i}{2 + 2 \cos(j_0 e^{2j_0 f})};$$
 (1)

We denote by $%_0$; the corresponding density matrix, whose symmetrically ordered characteristic function is given by

where D () = \exp (a^{\prime} a) denotes the displacement operator. The corresponding W igner function is defined as

$$W_{0}$$
; () $\frac{1}{2}$ e $_{0}$; () d^{2} : (3)

From now on, let us move to quadrature variables x and p, de ned through $= (x + ip) = \frac{p}{2}$. By de ning

e
$$\begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$$
; X $\begin{pmatrix} x \\ p \end{pmatrix}$; (4)

one can write the W igner function W: (x;p) as follows

The rst two G aussian terms are related to the projectors join ojand join oj; they are the W igner functions of the two coherent states joi and joi. The remaining two terms correspond to non diagonal operators and are responsible for the intereference elects which characterize a coherent superposition.

We now move to the study of a \squeezed cat", de ned as the superposition of two squeezed coherent states. Let us introduce the operator b by means of a Bogoliubov transform ation

b
$$a + 4;$$
 with j^{2} j^{2} $j = 1:$ (6)

and the states j i as its eigenvectors: bj i = j i. Such states are known in the literature as 'two{photon coherent states' and are indeed squeezed coherent states, according to the following well-known relation [13]

$$j i = D ()S (r_0; r_0; r_0) \mathcal{D}i;$$
 (7)

with the requirem ents

= ;
$$\cosh r_0 =$$
; $e^{i2'_0} \sinh r_0 =$; (8)

and the squeezing operator de ned as S (r;') = $\exp(\frac{1}{2}re^{i2'}a^2 - \frac{1}{2}re^{i2'}a^{y2})$.

W e consider the following superposition

$$j_0; i = \frac{j_0i + e^i j_0i}{2 + 2\cos()e^{2j_0}\hat{j}};$$
 (9)

where the states j 0 i are eigenstates of b: such a state will be referred to as to a \squeezed cat" state.

The W igner representation of the state j $_0$ i can be easily found by recalling that a two{ photon coherent state j $_0$ i m ay as well be written as

$$j_0 i = S(r_0; r_0)D(r_0) \forall i;$$

with the squeezing param eters r_0 and r_0 determined by Eqs. (8). This means that one can promptly derive the W igner function W r_0 ; of a squeezed cat state by simply replacing r_0 with r_0 in Eq. (3) and then applying a squeezing transform ation. In the following we will set $r_0 = 0$, without loss of generality, as a reference choice for phase space rotation.

The squeezing transform ation im plemented by S $(r_0;0)$ corresponds, in terms of the phase{ space variables $X = {x \atop p}$ to the map

$$X ! R ^{1}X ; w ith R = diag (e^{r_0}; e^{r_0}) :$$
 (10)

Applying such a transform ation to the coherent cat W igner function eventually yields

with

$$_{0}$$
 ReR: (12)

Of course, for $r_0 = 0$ and 0 = 0, one recovers Eq. (3) for a coherent cat.

3 Time evolution in noisy channels

We now consider the evolution in time of an initial squeezed cat state put in a noisy channel, in presence of damping and/or pumping toward an asymptotic squeezed thermal state. The system is governed, in the interaction picture, by the following master equation [13]

$$\frac{8}{2} = -N L [a^{y}] + - (N + 1) L [a] - M D [a] + M D [a^{y}] ;$$
 (13)

where the dot stands for tim e{derivative, the Lindblad superoperators are de ned by L [0]% $20\%0^{\circ}$ 0° 0% 0° 0% 0

As is well known, Eq. (13) can be transformed into a linear Fokker{P lanck equation for the W igner function of the system [13]. M oreover, the G aussian solutions of such equation have been thoroughly analysed in previous works [6, 14]. The initial condition we consider here is described by the W igner function W $_{0}$; of Eq. (12), which is just a linear combination of G aussian terms. Therefore, its evolution in the noisy channel can be followed straightforwardly by exploiting the general results derived in Ref. [14]: each G aussian terms evolves independently and it su cess to follow the time dependence of its second statistical moments.

Let $_{ij}$ $\frac{1}{2}h\hat{x}_i\hat{x}_j+\hat{x}_j\hat{x}_{i}i$ $h\hat{x}_ih\hat{x}_ji$ and X_{0i} $h\hat{x}_ii$ be, respectively, the covariance m atrix and the vector of the rst m oments of a Gaussian state $(\hat{x}_1;\hat{x}_2=\hat{x}_i;\hat{p})$ being the quadrature phase operators). Then, the time equadrature of (t) and X_0 (t) in the squeezed therm alchannel is described by the following Eqs. [14]

$$X_0(t) = e^{\frac{1}{2}t}X_0(0);$$
 (14)

$$(t) = {}_{1} 1 e^{t} + (0) e^{t};$$
 (15)

The time-dependent solution for the W igner function W $_{\scriptscriptstyle 0}$; (t) of an initial squeezed cat is thus readily found and reads

with (t) given by Eqs. (15, 16). The rst moments of each Gaussian term are exponentially damped in the channel. Any initial cat state is attracted toward an asymptotic centered

squeezed therm alstate with Wigner function

$$W_{1}(x;p) = \frac{e^{\frac{1}{2}X^{T} \cdot 1^{1}X}}{P_{Det_{1}}} :$$
 (18)

This state, like all asymptotic quantities, is a property of the channel and does not depend on the initial state.

4 Decoherence of an initial cat state

In order to quantify decoherence of the state caused by environm ental noise, we consider the loss of purity. The degree of purity of a continuous variable quantum state % can be e ectively characterized either by its Von Neum ann entropy S_V Tr(% ln %) or by its linear entropy S_1

$$S_1 \quad 1 \quad Tr(^2) \quad 1 = 1 - \frac{2}{2 \cdot R^2} W^2 dx dp$$
: (19)

In the following we will adopt linear entropy, which can be conveniently evaluated. The quantity = Tr^2 , conjugate to S_1 , will be referred to as the 'purity' from now on.

For an initial pure cat state (=1) at time t=0, the asymptotic value $_1$ can be obtained from Eq. (18) by straightforward integration [14]

$$_{1} = \frac{1}{2^{p} \frac{1}{D \text{ et }_{1}}} = \frac{1}{p \frac{1}{(2N + 1)^{2} + 4 \frac{M}{2}}}$$
: (20)

At nite times the purity of a decohering cat can be determined integrating the function W $_{0}$; (x;p) given by Eq. (17), according to Eq. (19). The integration can be promptly performed with the help of the following thum b rule

Exploiting the above rule one eventually has

with

A (t)
$$\frac{S_{xx}(t) - S_{pp}(t) - 2iS_{xp}(t)}{4}$$
 (23)

and

$$S(t) R (t)^{1}R ; T(t) (Det)^{1}S(t)^{1}$$
 (24)

Eq. (22) shows that $_0$; is a decreasing function of the initial parameters x_0 and p_0 . This should be expected: the bigger' the cat is, the faster it decoheres. For $_0$ = 0 the superposition




Figure 1: Evolution of the purity for di erent initial cat states and channels. In all cases the asymptotic purity of the bath is $_1=0.5$. The dotted line shows the behaviour of an initial cat with $\mathbf{x}_0=\mathbf{p}_0=1$ and $\mathbf{r}_0=0$ in a non squeezed channel. The dashed and the continuous lines refer to an initial cat with $\mathbf{x}_0=\mathbf{p}_0=100$ and $\mathbf{r}_0=0$ evolving, respectively, in a non squeezed channel and in a squeezed channel with $\mathbf{M}=2+2i$. The dot{dashed line refers to an initial cat state with $\mathbf{x}_0=\mathbf{p}_0=10$ and $\mathbf{r}_0=2$, evolving in a non squeezed channel.

disappears and the cat state reduces to a centered squeezed state; the latter, whose evolution in noisy channels has been studied in Ref. [14], decoheres more slow by than the corresponding cat state. The numerical analysis shows that the phase has little election the behaviour of purity at large times: in fact, as it is evident from Eq. (22), all the terms involving are suppressed by Gaussian terms of the form Exp (x_0^2 y_0^2). In Fig. 1 the behaviour of purity over the full temporal range up to the asymptotic regime is showed for various choices of the parameters of the channel and of the initial state.

One feature which is most evident in all instances is the fast initial fall of the purity. Although the minimum value of the purity attained in such a steep descent can vary, the temporal scale in which the minimum is reached is always of the order of $^{-1}$, which is indeed the only time characterizing the losses in the channel. Besides, it can be seen that the general behaviour of purity in squeezed baths is quite the same as in non squeezed ones. One can see as well that the value of the rst minimum of the purity depends drastically on the squeezing parameter r_0 , decreasing with greater r_0 , while, for a given squeezing parameter, an increase in the parameters x_0 and p_0 delays the reaching of the asymptotic purity (see Fig. 1).

5 Optim alregim es

Optim al regimes with minimized decoherence can be determined by maximizing the purity at any given time for xed values of the parameters of the channel and of the initial state. Notice that, as we have shown in the previous section, a cat like state decoheres on a time scale of the order of the photon lifetime ¹, regardless of the choice of the parameters of both the Gaussian reservoir and the initial pure cat state. This fact is a manifestation of a fundamental feature of quantum mechanics: once a single photon is added or lost, all the information contained in a coherent superposition leaks out to the environment and is therefore lost as well, together with the possibility of detecting such a coherent behaviour by means of interferometry [15, 16]. A simple, meaningful example in this respect is just a coherent even cat j₀;0i subjected to

damping. Under the loss of a photon, such a state jumps into aj $_0$; 0i / j $_0$; i, which is an odd cat and has got opposite interference terms. Therefore, as soon as the probability of losing a photon reaches 0.5, the original superposition turns into an incoherent m ixture of an even and an odd cat, whose interference terms cancel out each other exactly [16].

Actually, a more detailed analysis would reveal that decoherence times are even shorter than 1 : for a coherent cat j ${}_0$; i evolving in a thermal environment, coherence is lost at $t_{\rm dec} = {}^1 = 2 \, j_0 \, j^2$ [5]. In view of these considerations, we are interested in maximizing the purity in the time region $t \cdot 1 = 2 \, j_0 \, j^2$.

A relevant question in such a context is: given an initial bath and an initial squeezing of the cat{like state, which is the optimal phase space direction of $_0$ j_0je^i at xed j_0j^2 N ote that the last condition can be seen as a constraint on the energy of the cat{like state. O by iously, for a non squeezed cat in a thermal channel the symmetry of the problem forbids the existence of a privileged direction.

Interesting issues come instead from the consideration of a squeezed cat in a thermal channel and a non squeezed cat in a squeezed bath. For the moment being, let us consider the instance of an 'even' cat, i.e. of a cat with coherent phase = 0. The dependence of the purity on 0 is essentially contained in exponentials of quadratic form s, see Eq. (22). The algebraic analysis of such terms in the case of a squeezed cat in thermal baths and of a coherent cat in squeezed baths is quite easy (see App. A). If the squeezing is performed on the inital cat state, the coherence is better preserved if $_0$ is chosen in the same phase space direction of the variance which is suppressed by squeezing. With our notation, this = $'_0$ + =2 (where $'_0$ is the squeezing angle). On the contrary, if the squeezing is performed on the bath in the direction $'_1$, the choice $= '_1$ turns out to be the best one to slow down the rate of decoherence. This som ehow counterintuitive situation is due to the existence of complex fringe patterns of a cat state in phase space, especially in the presence of squeezing. Actually, preserving quantum coherence is crucially related to the persistance of the interference fringes: reducing quantum uctuations in a phase space direction protects the fringes from degradation and the cat state from decoherence. With the above expedient choices of the phase , squeezing does actually in proves the coherence of the superposition at small times with respect to dissipation in a phase insensitive setting [6].

Now, let us suppose that for an even cat state the phase — is optimally chosen and let us consider a channel with asymptotic purity $_1$. It can be easily shown (cfr. App. A) that the purity of an initially squeezed cat (with squeezing r) in such a thermal channel equals, at any given time, the one of a non squeezed cat evolving in a squeezed channel with the same squeezing r. The same protection against decoherence provided by the squeezing of the bath can be obtained, in a thermal phase insensitive channel, by squeezing the initial even cat state of the same amount in an orthogonal direction.

We nally remark that, with an optimal phase setting, an optimal nite value of the squeezing parameter r does indeed exist (see Fig. 2). This fact can be best appreciated from Eq. (22): even if the exponential terms increase with increasing r, the factor D et $^{1-2}$ decreases with r. The value of r allowing the maximum slowing down of decoherence increases with increasing j₀j.

We now brie y consider the instance = =2, which is the one considered originally by Yurke and Stoler as being produced by a unitary evolution in Kerrmedia. In such a case, the last term in Eq. (22) vanishes, so that the optimal choice for can be easily found for any choice of r_0 , r_1 and r_1 . To this end, it is su cient to determ in the eigenvector corresponding to the smallest eigenvalue of the matrix S (t) r_1 , which coincides with the analogous eigenvector of R r_1 and does not depend on time. However, we once again stress that, as soon




Figure 2: Com parison between the evolution of purity of an initial non squeezed cat in a non squeezed bath (continuous line) and of initial cats in squeezed con gurations with optimal phase choices. The dashed line refers to an initial squeezed cat with $r_0=1$, whereas the dotted line refers to an initial squeezed cat with $r_0=1.5$. In all instances $r_0=0.5$, $r_0=0.5$, $r_0=0.5$, $r_0=0.5$, $r_0=0.5$, in good agreement with the initial decrease of purity. The choice $r_0=0.5$ 1 appears to be optimal for such a value of $r_0=0.5$, $r_0=0.5$.

as one deals with m esoscopic cat states (so that Exp ($j_0 \hat{j}$) 1), the dependence on the coherent phase is severely suppressed and all the considerations made for even cat states still hold, regardless of the choice of . Results for the evolution of purity at small times are sum marized in Fig. 2.

6 Conclusions

The study of the decoherence of initial coherent and squeezed coherent cat states in arbitrary G aussian reservoirs has been carefully carried out by determining the exact time evolution of the purity of the state. Optimal settings that minimize the rate of decoherence in relevant congurations have been determined.

In particular, we have shown that the same protection against decoherence granted by a squeezed bath can be achieved by squeezing the initial cat{like state. In view of the well known diculties involved in the experimental realization of squeezed baths, even as elective descriptions of feedback schemes [7], this equivalence could provide a relevant alternative option for experimental purposes [4].

A cknow ledgem ents

We thank David Vitali for useful discussions. AS, FI and SDS thank INFM, INFN, and MIUR under national project PRIN-COFIN 2002 for nancial support. MGAP is a research fellow at Collegio Volta, and he thanks R.F.Antoni for continuing inspiration.

A Appendix

In this appendix we analytically single out the optim alphase space orientations of the cat state for the con gurations discussed in Section 5. The orientation of the cat state is determined by the angle = $A \operatorname{rg}(0)$. In the following we will set = 0 and de ne, for ease of notation, u $(1 \operatorname{Exp}(t))=2_1$ and $\operatorname{Exp}(t)=2_1$. The squeezing parameter $\operatorname{rg}(t)=2_1$ of the bath is determined by $\cosh(2r_1)=\frac{1}{1+4}$ in $\operatorname{rg}(t)=2_1$.

We rst consider a squeezed cat in a thermal channel, with $r_0 \notin 0 = r_1$. In this instance, one has

$$S(t) = diagf e^{2r_0} (u + e^{2r_0}v)^{-1}; e^{-2r_0} (u + e^{-2r_0}v)^{-1}q;$$

and

A (t) =
$$(2D \text{ et } (t))^{-1} u \sinh (2r_0)$$
:

Substituting these expressions in Eq. (22), it is easy to see that, for $x = 2j_0^2 = x_0^2 + p_0^2$, all the exponential terms are maximized by the choice $x_0 = 0$, $p_0 = 2j_0^2$. This corresponds to $x_0 = 2 = 2j_0^2 + 2j_0^2$.

A nalogously, for a non squeezed cat in a squeezed channel (with r_0 = 0 f r_1), one gets 1

$$S(t) = diagfe^{2r_1} (u + e^{2r_1} v)^{-1}; e^{2r_1} (u + e^{2r_1} v)^{-1}q;$$

and

A (t) =
$$(2D \text{ et } (t))^{1} u \sinh (2r_{1})$$
:

Eq. (22) shows that the choice $x_0 = \begin{pmatrix} p \\ 2j_0 j_1 p_0 = 0 \end{pmatrix}$ (corresponding to $p_0 = p_0 = p_0 = p_0$) is optimal in this case.

Finally, it is easy to verify that, adopting such optim alchoices and putting $r_0=r_1$, all the exponential term s entering in Eq. (22) take the same values for a squeezed cat in a therm alchannel and a non squeezed cat in a squeezed channel. Since Det depends only on the dierence jr_0 r_1 j [14], this implies that the time evolutions of purity are identical in these two instances.

R eferences

R eferences

- [1] E. Schrodinger, Naturwissenschaften 23, 812 (1935).
- [2] B. Yurke and D. Stoler, Phys. Rev. Lett. 57, 13 (1986).
- [3] A. Mecozzi and P. Tombesi, Phys. Rev. Lett. 58, 1055 (1987); B. Yurke, W. Schleich, and D. F. Walls, Phys. Rev. A 42, 1703 (1990); S. Song, C. M. Caves, and B. Yurke, Phys. Rev. A 41, 5261 (1990); T. Ogawa, M. Ueda, N. Imoto, Phys. Rev. A 43, 6458 (1991); M. Dakna, T. Anhut, T. Opatmy, L. Knoll, and D. G. Welsch, Phys. Rev. A 55, 3184 (1997); E. Solano, R. L. de Matos Filho, and N. Zagury, Phys. Rev. Lett. 87, 060402 (2001); M. Patemostro, M. S. Kim, and B. S. Ham, Phys. Rev. A 67, 023811 (2003).
- [4] M.G.A.Paris, J.Opt.B:Quantum Semiclass.Opt.1, 662 (1999).

 $^{^{1}}$ W e can set the squeezing angle of the bath $'_{1}$ = 0 because the reference choice is still free, being the initial cat non squeezed. This corresponds to choosing the squeezing M real.

- [5] D.F.Walls, G.J.Milbum, Phys. Rev. A 31, 2403 (1985).
- [6] T.A.B.Kennedy and D.F.Walls, Phys. Rev. A 37, 152 (1988).
- [7] P. Goetsch, P. Tombesi, and D. Vitali, Phys. Rev. A 54, 4519 (1996); S. Zippilli, D. Vitali, P. Tombesi, and J.M. Raimond, ibid. 67, 052101 (2003).
- [8] F.A.A.ElOrany, Phys. Rev. A 65, 043814 (2002).
- [9] M. Brune, E. Hagley, J. Dreyer, X. Ma^tre, A. Maali, C. Wunderlich, JM. Raim ond, and S. Haroche, Phys. Rev. Lett. 77, 4887 (1996); A. Au eves, P. Maioli, T. Meunier, S. Gleyzes, G. Nogues, M. Brune, JM. Raim ond, and S. Haroche, quant-ph/0307185 (2003).
- [10] E.Solano, G.S.Agarwal, and H.Walther, Phys.Rev.Lett.90,027903 (2003); P.Lougovski, E.Solano, and H.Walther, quant-ph/0308059 (2003); P.Lougovski, C. Casagrande, A.Lulli, B.G. Englert, E.Solano, and H.Walther, quant-ph/0309040 (2003).
- [11] M.-A. Dupertuis and S. Stenholm, J. Opt. Soc. Am. B 4, 1094 (1987); M.-A. Dupertuis, S. M. Barnett, and S. Stenholm, ibid. 4, 1102 (1987); Z. Ficek and P. D. Drummond, Phys. Rev. A 43, 6247 (1991); C. W. Gardiner and P. Zoller, Quantum Noise, (Springer Verlag, Berlin, 1999).
- [12] P. Tom besi and D. Vitali, Phys. Rev. A 50, 4253 (1994); N. Lutkenhaus, J. I. Cirac, and P. Zoller, ibid. 58, 548 (1998).
- [13] D.F.W alls and G.J.M ilbum, Quantum Optics, Springer Verlag, Berlin (1994).
- [14] M.G.A. Paris, F. Illum inati, A. Sera ni, and S. De Siena, Phys. Rev. A 68, 012314 (2003).
- [15] C.T.Lee, Phys. Rev. A 44, R2275 (1991).
- [16] S.H aroche, Introduction to quantum optics and decoherence, lectures held at Les Houches Sum mer School Quantum entanglement and information processing (2003).