

# Experimental demonstration of an efficient quantum phase-covariant cloning and its possible applications to simulating eavesdropping in quantum cryptography

Jiangfeng Du,<sup>1,2,3,\*</sup> Thomas Durt,<sup>4</sup> Ping Zou,<sup>1</sup> L.C. Kwek,<sup>5</sup> C.H. Lai,<sup>2</sup> C.H. Oh,<sup>2</sup> and Artur Ekert<sup>3,2</sup>

<sup>1</sup>Structural Research Laboratory and Department of Modern Physics,  
University of Science and Technology of China, Hefei, 230027, P.R. China

<sup>2</sup>Department of Physics, National University of Singapore, 2 Science Drive 3, Singapore 117542

<sup>3</sup>Centre for Quantum Computation, DAMTP, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA U.K.

<sup>4</sup>TONA-TENA Free University of Brussels, Pleinlaan 2, B-1050 Brussels, Belgium.

<sup>5</sup>Department of Natural Sciences, National Institute of Education,  
Nanyang Technological University, 1 Nanyang Walk, Singapore 637616

We describe a nuclear magnetic resonance (NMR) experiment which implements an efficient one-to-two qubit phase-covariant cloning machine (QPCCM). In the experiment we have achieved remarkably high fidelities of cloning, 0.848 and 0.844 respectively for the original and the blank qubit. This experimental value is close to the optimal theoretical value of 0.854. We have also demonstrated how to use our phase-covariant cloning machine for quantum simulations of bit by bit eavesdropping in the four-state quantum key distribution protocol.

PACS numbers:

The “no-cloning” result [1, 2] asserts that due to the linearity of quantum mechanics unknown quantum states cannot be copied perfectly. This notwithstanding one can design approximate quantum cloning machines and address their optimality. The most notable example is the universal quantum cloner (UQC) proposed by Bužek and Hillery [3]. It has been studied in great details [4] and a number of experimental implementations of a  $1 \rightarrow 2$  qubit UQC have been proposed [5, 6, 7, 8]. Another important example is the optimal quantum phase-covariant cloning machine (QPCCM) [9, 10, 11]. Unlike the UQC, it clones only subsets of states for which we have some *a priori* information. In the special case of the QPCCM operating on qubits it has been shown that a class of states  $|\psi\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{i\varphi}|1\rangle)$ , called equatorial states, can be cloned up to the fidelity 0.854. As expected this value is slightly higher than the optimal fidelity of the UQC (0.833). This is because even partial information about the original state allows to optimize the cloning process and to obtain higher fidelities of the clones. The phase-covariant cloners, which are the subject of this paper, are of significant importance in quantum cryptography as they provide the optimal eavesdropping technique for a large class of attacks on the four state protocol (BB84) [16]. The properties of the QPCCM have been extensively studied from the theoretical perspective [12, 13], however, on the experimental side, apart from an interesting recent optical proposal by Fiurasek [14], no actual realization of the QPCCM has been reported.

Here we describe the first experimental implementation of the QPCCM. We use the NMR technology to implement a modified two qubit network originally designed by Niu and Griffiths [10] (see Fig. 1). The simplicity of the network allows to reduce the effects of decoherence and to obtain remarkably high fidelities of the clones.




FIG. 1: Quantum network of the efficient phase-covariant cloning. It consists of two controlled-NOT gates together with one controlled-rotation gate, where  $R(\theta) = e^{-i\theta\sigma_y/2}$  is a rotation by an angle  $\theta$  about the  $y$  axis in Bloch-sphere. The upper and the lower horizontal lines correspond to the original and the blank qubits respectively.

This is in contrast with earlier approaches which were based on more complicated three qubit networks [9]. If the complexity of related three qubits experiments is of any guidance here, e.g. the NMR implementation of the UQC [5], then substantial losses due to inhomogeneities of the magnetic field and decoherence cannot be avoided with the current state of the art technology. This, together with the stringent precision requirements, lowers the fidelity (to about inconclusive 58% in [5]). A three qubit network for the  $1 \rightarrow 2$  QPCCM, for example the one proposed by Fuchs et al. [9], would face similar problems.

Our version of the network is shown in Fig. 1. The net unitary operator has the form

$$U(\theta) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

When  $\theta = \frac{\pi}{4}$ , this unitary transform defined as  $U^{opt}$  corresponds to an efficient optimal QPCCM. In fact,  $U^{opt}$  is just a 2-qubit square root of SWAP gate. Consider now an equatorial state of  $a$  qubit, i.e., a state with a definite spin in the direction  $n = (\cos \varphi, \sin \varphi, 0)$ . This

state has the form  $|n\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{i\varphi}|1\rangle)$ , whereas the  $b$  qubit is in the state  $|0\rangle$ .  $U^{opt}$  transforms the input state to  $\rho_{ab}^{out} = U^{opt}|n\rangle\langle 0| \langle n| U^{opt\dagger}$ . The reduced density matrix of two copies can be calculated (by tracing out another qubit) as

$$\rho_a^{out} = \rho_b^{out} = \begin{pmatrix} 3/4 & \sqrt{2}e^{-i\varphi}/4 \\ \sqrt{2}e^{i\varphi}/4 & 1/4 \end{pmatrix}.$$

Note that the two copies are in fact symmetric. We use the fidelity  $F = \langle n|\rho^{out}|n\rangle$  to define the quality of the copies. As expected, the optimal fidelity for  $1 \rightarrow 2$  QPCCM is  $F_{QPCCM}^{opt} = \frac{1}{2} + \frac{\sqrt{2}}{8} = 0.854$ , which is higher than the optimal value  $F_{QUCM}^{opt} = 0.833$ .

This quantum circuit of QPCCM is realized by using a two-qubit NMR quantum computer, based on  $^{13}\text{C}$  and the  $^1\text{H}$  nuclei in Carbon-13 labelled chloroform (Cambridge Isotopes) dissolved in  $\text{d}_6$  acetone. The  $^{13}\text{C}$  nucleus was used as qubit  $a$ , and  $^1\text{H}$  as qubit  $b$ . The reduced Hamiltonian of the 2-spin ensemble is given by  $H = \omega_a I_z^a + \omega_b I_z^b + 2\pi J I_z^a I_z^b$ , where the first two terms describe the free precession of spin  $a$  ( $^{13}\text{C}$ ) and  $b$  ( $^1\text{H}$ ) around the static magnetic field with frequencies 100MHz and 400 MHz.  $I_z^a$  ( $I_z^b$ ) is the angular momentum operator of  $a$  ( $b$ ) in direction  $\hat{z}$ , and the third term is the  $J$  coupling of the two spins with  $J = 214.5\text{Hz}$ .  $^{13}\text{C}$  nucleus's  $T_1$  relaxation time is 17.2s and it's  $T_2$  relaxation time is 0.35s.  $^1\text{H}$  nucleus's  $T_1$  relaxation time is 4.8s and it's  $T_2$  relaxation time is 3.3s. In the following, we describe how we experimentally realize the optimal  $1 \rightarrow 2$  QPCCM shown in Fig. 1.

(E1) Prepare the initial state: Initially the two qubits are in thermal equilibrium with the environment and their state is described by the density operator  $\rho_{th} \propto \sigma_z^a + 4\sigma_z^b$ . We use the spatial averaging technique [19] to create the effective pure state  $|\uparrow\rangle_a \otimes |\uparrow\rangle_b$ , or, in the density operator form,  $\frac{1}{2}(1 + \sigma_z^a) \otimes \frac{1}{2}(1 + \sigma_z^b)$ . The sequence of operations leading to this state is shown in Fig. 2(a). We then perform a single hard  $\frac{\pi}{2}$  radio frequency ( $rf$ ) pulse on  $a$  qubit to generate one of the desired equatorial state  $|n(\varphi)\rangle_a^{in} = (\cos\varphi, \sin\varphi, 0)$  with  $\varphi = \cos(n\pi/12)$ ,  $n = \{0, 1, \dots, 23\}$ .

(E2) Clone the input equatorial state: The quantum circuit of optimal  $1 \rightarrow 2$  QPCCM is described in Fig. 1 by fixing  $\theta = \frac{\pi}{4}$ . This corresponds to a 2-qubit square root of SWAP gate. NMR pulse sequences are developed by replacing this operation with an idealized sequence of NMR pulses and delays. The resulting sequences are then simplified by combining  $rf$  pulses appropriately. Figure 2(b) shows the final pulse sequence to demonstrate the optimal  $1 \rightarrow 2$  QPCCM. All the  $rf$  pulses are hard pulses which hardly affect the state of  $b$  qubit due to the heteronuclear sample we used.

(E3) Measure and analyze: In principle, the quality of the copies, defined as fidelity, can be calculated by  $F = \langle n|\rho_{a(b)}^{out}|n\rangle$ , where  $\rho_{a(b)}^{out}$  is the reduced density ma-




FIG. 2: NMR pulse sequences. The white and black boxes are  $90^\circ$  and  $180^\circ$  pulses, while the grey boxes are pulses with other flip angles shown above each box; pulse phases and gradient directions are shown below each pulse. All  $rf$  pulses are hard pulses with rectangular shape and  $5\mu\text{s}$  pulse width. Delay times are  $\tau_1 = 1/(4J)$  and  $\tau_2 = \theta \cdot \tau_1/\pi$ . In the QPCCM experiment we take  $\theta = \pi/4$  and change the value of  $\theta \in [0, 2\pi]$ .

trix of a single qubit and can be obtained from the density matrix  $\rho_{ab}^{out}$ . In NMR, one can use state tomography technique to get  $\rho_{ab}^{out}$  by applying a set of readout pulses, but this has the disadvantage of requiring separate experiments. In our experiment, we use a simpler method described in Ref.[5]: we measure the two spectra of two output qubits individually, here the receiver phase are set with the same phase as that of the input qubit measurement. Therefore, the tracing out process can be implemented by integrating the entire multiplet in each spectrum, comparing to the integration of the input state spectrum, we can obtain the relative length of the output state vector  $r'_{a(b)}$  in the same orientation as its input state vector, so the fidelity between the input and output state can be calculated as  $F_{a(b)} = \frac{1}{2}(1 + r'_{a(b)})$ . Figure 3 shows the experimental results from cloning the input equatorial state  $|n(0)\rangle_a^{in} = (1, 0, 0)$ . There are three spectra, corresponding to the observable NMR signals of one input state and its two copies, that are measured by setting the same receiver phase experimentally. The spectra do have similar expected form (in-phase absorption signals at the outmost positions of each multiplet).

An important feature of QPCCM is that all equatorial states are cloned equally well and so it is necessary to study the behavior of pulse sequence when applied to a wide range of states on equator. We have prepared a total of 24 input equatorial states  $|n(\varphi)\rangle_a^{in} = (\cos\varphi, \sin\varphi, 0)$  by changing the value  $\varphi$  with a spacing of  $15^\circ$  as shown in (E1). For each input state, we measure its spectrum and denote it as a reference to calculate the quality of the two copies after the cloning transformation described in (E2). Finally we measure each copy and calculate the fidelity. Experimentally, we get the mean fidelity of this phase-covariant cloning are  $F_a = 0.848 \pm 0.015$  for  $a$  qubit




FIG. 3: Experimental spectra of cloning the input state  $|n(0)\rangle_a^{in} = (1, 0, 0)$ . The left and middle spectra are the carbon spectra corresponding to the input and output state of  $a$  qubit, where the vertical scales are in the same arbitrary units. The right spectrum is for the hydrogen nucleus representing the output of  $b$  qubit, where the vertical scale does not share the same arbitrary units with those of the carbon spectra. From the integration of the each multiplet, we obtain the fidelities of the two copies,  $F_a = 0.842$  and  $F_b = 0.839$ .

and  $F_b = 0.844 \pm 0.015$  for  $b$  qubit, which are both close to the optimal theoretical value 0.854.

Compared to the low fidelity of the NMR experiment for UQCM [5], our near-optimal fidelity arises from the following reasons: (1) Less decoherence effect – the time used for cloning in our experiment is about  $5.3ms$ , which is well within the decoherence time (about  $350ms$  for  $^{13}C$  nucleus and  $3.3s$  for  $^1H$  nucleus); while the time used for UQCM in Ref. [5] was estimated about  $400ms$ , which is close to the decoherence time with the value  $720ms$  for two  $^1H$  nuclei; (2) Simplicity – it is simpler to realize our economic 2-qubit QPCCM than to realize the 3-qubit UQCM in Ref. [5]; (3) Pulses – in our experiment all the  $rf$  pulses are hard pulses, which are more perfect than selective pulses, this is simply achieved by using heteronuclear sample. In our experiments, small errors arise as a result of the inhomogeneity of the static with  $rf$  magnetic fields as well as the variability of the measurement.

One important application of our efficient QPCCM is the quantum simulator of a bit by eavesdropping on the four-state protocol of quantum cryptography [10, 17]. (The UQCM plays the same role for the six-state protocols [18]). The four-state protocol (also known as BB84) uses four quantum states, say  $|\pm\rangle_x = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$  and  $|\pm\rangle_y = \frac{1}{\sqrt{2}}(|0\rangle \pm i|1\rangle)$ , that constitute two maximally conjugate bases. Alice chooses one of these four states in her qubit (denoted by  $a$ ) at random, then sends it to Bob. Whatever the state is, Eve approximately clone two copies of this input state by inserting a  $1 \rightarrow 2$  QPCCM in the quantum channel. She then sends one copy to Bob and stores another copy in her qubit  $b$ . Thereafter, Bob measures qubit  $a$  in one of the two bases chosen at random. Finally Alice announces publicly the basis she used for transmission of the signal, and in those cases in which Bob measures in the same bases (these cases are useful, the others are discarded by Alice and Bob). Eve, who now knows the bases Alice employed, measures  $b$  qubit in order to estimate which signal Alice sent.

Experimentally, we realize the above process of eaves-




FIG. 4: The normalized observable NMR signals  $\langle\sigma\rangle$  versus the rotation angle  $\theta$  of the phase-covariant cloning machine, for two bases, X-base(at Left) and Y-base(at right). The lines correspond to theoretical calculation. The filled and empty boxes (circles) correspond to the experimental measurement from  $a$  qubit ( $b$  qubit), while the filled (empty) circles and boxes correspond to the input state is  $|+\rangle$  ( $|-\rangle$ ).

dropping attack as following: (1) Prepare one of four BB84 states by using the same method that we described in (E1); (2) Perform the quantum  $1 \rightarrow 2$  QPCCM described in Fig. 1. The pulse sequence to realize the network is shown in Fig. 2(b). Note that  $\theta$  is not fixed to  $\pi/4$  as in the previous optimal  $1 \rightarrow 2$  QPCCM realization, we set various rotation angles  $\theta \in [0, 2\pi]$  of the pulse [marked as grey box in Fig.2(b)] with a space of  $\pi/12$ ; (3) For each experiment, we perform two measurements of the NMR observable signals  $\langle\sigma_i\rangle$  for Bob and Eve individually, here  $i \in \{x, y\}$  is the same base as that of the input state. Recall another measurement for the input state which is used as reference spectra, there are 3 measurements. Totally, we perform  $4 \times 24 \times 3 = 288$  experiments distinguished by 4 BB84 states and 24 rotation angles  $\theta \in [0, 2\pi]$  and 3 measurements. Theoretically, if Alice sends the state  $|\pm\rangle_x$ , Bob measures  $\langle\sigma_x^{Bob}\rangle = \pm \frac{\cos\theta}{2}$  and Eve measures  $\langle\sigma_x^{Eve}\rangle = \pm \frac{\sin\theta}{2}$  in X-base; if Alice send state  $|\pm\rangle_y$ , Eve measures  $\langle\sigma_y^{Eve}\rangle = \pm \frac{\sin\theta}{2}$  while Bob measures  $\langle\sigma_y^{Bob}\rangle = \pm \frac{\cos\theta}{2}$  in Y-base. Both the theoretical and experimental results are plotted in Fig. 3, where the symmetry between  $|+\rangle$  and  $|-\rangle$  in each base is clearly seen.

A main concern of the eavesdropping is to determine how much information an eavesdropper can obtain from a given level of noise. For the above optimal eavesdropping attack, regardless of the input BB84 state, Bob guesses correctly the state sent by Alice with probability  $F_{Bob} = \frac{1}{2} + \langle\sigma_i^{Bob}\rangle$  and makes an error  $D_{Bob} = 1 - F_{Bob} = \frac{1}{2} - \langle\sigma_i^{Bob}\rangle$ , where  $i \in \{x, y\}$  is one of the maximally conjugate bases; while Eve guesses correctly the state sent by Alice with probability  $F_{Eve} = \frac{1}{2} + \langle\sigma_i^{Eve}\rangle$  and makes an error  $D_{Eve} = \frac{1}{2} - \langle\sigma_i^{Eve}\rangle$ . As we know, the mutual information is defined as  $I = \frac{1}{2} + D \log_2 D + (1 - D) \log_2 (1 - D)$ . From our experimental dates shown in Fig. 4, we extract the Alice-Bob and Alice-Eve mutual information as a function of the




FIG. 5: Mutual information versus the quantum bit error rate. The lines correspond to theoretical calculation. The boxes correspond to the experimental obtained mutual information of Alice and Bob ( $a$  qubit), while the circles correspond to the experimental mutual information of Alice and Eve ( $b$  qubit).

value of noise (QBER) defined as  $QBER = \frac{1-\cos\theta}{2}$ . Here  $\theta \in [0, \pi/2]$  characterize the strength of Eve's attack. The experimental results are shown in Fig. 5. We show the relation between the mutual information and QBER, in agreement with the theoretical results.

In summary, we provide the first experimental demonstration of an efficient and nearly optimal  $1 \rightarrow 2$  QPCCM by using a 2-qubit NMR quantum computer. Our approach cannot be extended to the UQC as it is known that a 3-qubit  $1 \rightarrow 2$  UQC cannot be reduced to an efficient 2-qubit network [13]. However, our efficient QPCCM has potential applications as a simulator of eavesdropping techniques in quantum key distributions.

This project was supported by the National Nature Science Foundation of China (Grants. No. 10075041 and No. 10075044) and Funded by the National Fundamental Research Program (2001CB309300). We also thank supports from the ASTAR Grant No. 012-104-0040 and Temasek Project in Quantum Information Technology (Grant No. R-144-000-071-305). T.D. thanks supports from the Flemish Fund for Scientific Research, the Inter-University Attraction Pole Program of the Belgian government under grant V-18, the Concerted Research Ac-

tion Photonics in Computing, and the research council (OZR) of the VUB.

\* Electronic address: djf@ustc.edu.cn

- [1] W.K. Wootters and W.H. Zurek, Nature (London) **299**, 802.
- [2] G.C. Ghirardi and T. Weber, Nuovo Cimento Soc. Ital. Fis. B **78**, 9 (1983). The impossibility of Exact cloning was first mentioned in a report by G.C. Ghirardi (1981) in response to a paper submitted to Found. Phys.
- [3] V. Bužek and M. Hillery, Phys. Rev. A **54**, 1844 (1996).
- [4] N. Gisin and S. Massar, Phys. Rev. Lett. **79**, 2153 (1997); D. Bruss et al., Phys. Rev. Lett. **81**, 2598 (1998); L.M. Duan and G.C. Guo, Phys. Rev. Lett. **80**, 4999 (1998); C. Simon et al., Phys. Rev. Lett. **84**, 2993 (2000); V. Bužek and M. Hillery, Phys. Rev. Lett. **81**, 5003 (1998); S.L. Braunstein, N.J. Cerf et al., Phys. Rev. Lett. **86**, 4938 (2001); G.M. D'Ariano et al., Phys. Rev. Lett. **86**, 914 (2001).
- [5] H.K. Cummins et al., Phys. Rev. Lett. **88**, 187901 (2002).
- [6] A. Lamas-Linares et al., Science **296**, 712 (2002).
- [7] S. Fasel et al., Phys. Rev. Lett. **89**, 107901 (2002).
- [8] Y. F. Huang, Phys. Rev. A **64**, 012315 (2001).
- [9] C. A. Fuchs et al., Phys. Rev. A **56**, 1163 (1997); R. B. Griffiths and C.-S. Niu, Phys. Rev. A **56**, 1173 (1997).
- [10] C.-S. Niu and R.B. Griffiths, Phys. Rev. A **60**, 2764 (1999).
- [11] D. Bruss et al., Phys. Rev. A **62** 012302 (2000).
- [12] G. M. D'Ariano and C. Macchiavello, Phys. Rev. A **64** 042308 (2001); H. Fan et al., Phys. Rev. A **65** 012304 (2002).
- [13] T. Durt and J. Du, e-print quant-ph/0309072.
- [14] J. Fiurasek, Phys. Rev. A **67**, 052314 (2003).
- [15] N. Gisin et al., Rev. Mod. Phys. **74**, 145 (2002).
- [16] C.H. Bennett and G. Brassard, in Proceedings of IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India (IEEE, New York, 1984) p. 175.
- [17] V. Scarani and N. Gisin, Phys. Rev. Lett. **87**, 117901 (2001)
- [18] D. Bruss, Phys. Rev. Lett. **81**, 3018 (1998); H. Bechmann-Pasquinucci and N. Gisin Phys. Rev. A **59**, 4238 (1999).
- [19] D.G. Cory et al., Physica D **120**, 82 (1998); J. Du et al., Phys. Rev. Lett. **91**, 100403 (2003).