

Decoherence from Internal Dynamics in Macroscopic Matter-Wave Interferometry

Sang Wook Kim¹ and Hwa-Kyun Park²¹Department of Physics Education and Research Center for Dielectric and Advanced Matter Physics
Pusan National University, Busan 609-735, Korea²Korea Institute for Advanced Study, 207-43 Cheongryangri-dong, Dongdaem-un-gu, Seoul 130-722, Korea
(Dated: March 29, 2024)

Usually, decoherence is generated from the coupling with an outer environment. However, a macroscopic object generically possesses its own environment in itself, namely the complicated dynamics of internal degrees of freedom. We address a question: when and how the internal dynamics decohere interference of the center of mass motion of a macroscopic object. We will show that weak localization of a macroscopic object in disordered potentials can be destroyed by such decoherence.

PACS numbers: 03.75.+b, 03.65.Yz, 42.25.Dd

I. INTRODUCTION

Superposition of states lies at the heart of quantum mechanics and gives rise to many of its paradoxes. Not only can a particle go through two paths simultaneously, but the wavefunction of a pair of particles flying apart from each other is also entangled into a non-separable superposition of states. However, such strange phenomena have never been observed in our macroscopic world. It has been an important question why and how quantum weirdness disappears in large systems [1].

Environment, usually described by a huge number of variables, can destroy coherence among the states of a quantum system. This is decoherence. The environment is watching the path followed by the system, and thus suppressing interference effects and quantum weirdness. In macroscopic systems, such process is so efficient that we see only its final result: the classical world around us [2, 3]. For truly macroscopic superpositions, decoherence occurs on a very short time-scale that it is almost impossible to observe quantum coherences. However, mesoscopic systems present the possibility of investigating the process of decoherence and the transition from quantum to classical behavior [4]. So far many experiments have been realized to generate mesoscopic superpositions [5, 6] and to decohere them in a controlled way [7]. Recently considerably large molecules have been used to investigate the decoherence, the transition from quantum to classical. For example, the researchers in Wien have observed interference of de Broglie waves of fullerenes (C_{60} or C_{70} molecules) and even bigger ones [8, 9, 10, 11, 12, 13]. In this experiment the fullerenes are quite hot as well as big, which means they contain complicated dynamics of their internal degrees of freedom. The internal thermal energy is almost one order of magnitude larger than the kinetic energy of its center of mass (CM) motion. A question naturally arises: is the complex dynamics of the internal degrees of freedom harmful for the interference of the CM motion?

Usually, decoherence has been generated from the coupling with an outer environment such as other particles or fluctuating electromagnetic fields. However, a macroscopic object generically possesses its own environment

in itself, namely the internal dynamics (ID) [14] when only small part of the total system, e.g. its CM, is under consideration. In this paper, we would like to address a question: when and how the ID decoheres interference of a macroscopic object. We also show nontrivial expectation that the weak localization of large molecules in a disordered potential can be destroyed by the decoherence generated from the ID without any external perturbation breaking the time reversal symmetry.

Let us consider a macroscopic object consisting of N particles exposed to the external potential V_{ex} . The total Hamiltonian can be written as

$$\begin{aligned} H &= \sum_i^N \frac{p_i^2}{2m_i} + V_{\text{in}}(\mathbf{f}; \mathbf{g}) + \sum_i^N V_{\text{ex}}(\mathbf{x}_i) \\ &= \frac{P^2}{2M} + V_{\text{ex}}(\mathbf{X}) + \sum_i^N \frac{p_i^2}{2m_i} + V_{\text{in}}(\mathbf{f}; \mathbf{g}) + \sum_i^N V_{\text{ex}}(\mathbf{x}_i) \\ &= H_{\text{CM}} + H_{\text{in}} + \dots; \end{aligned} \quad (1)$$

where V_{in} is an internal (or confinement) potential. P and X are a momentum and a coordinate of the CM with a mass $M (= \sum_i m_i)$ respectively, while p_i and x_i are the same quantities of the internal degrees of freedom with the reduced mass m_i . V_{in} represents the coupling between the CM and the ID. Since one finds V_{in} is determined only from the relative coordinates \mathbf{f}, \mathbf{g} , i.e. $V_{\text{in}}(\mathbf{f}; \mathbf{g}) = V_{\text{in}}(\mathbf{f} - \mathbf{g})$, the coupling term depends only on the external potential V_{ex} . It is easy to show that if V_{ex} does not correspond to a simple form such as constant, linear, and harmonic, the non-zero coupling always exists [15]. The CM motion can be entangled with its ID when the anharmonic external potential is applied. We call such a non-trivial external potential "nonlinear" since the corresponding force is nonlinear. It is noted that existence of the external potential has nothing to do with generation of the decoherence.

In a usual two slit experiment, a macroscopic object freely flies to a screen. Therefore, no entanglement between CM and ID arises, neither does the decoherence from ID. However, one can ask what happens if the repulsive potential produced by the slits is considered. For example, the van der Waals interaction between the C_{70}




FIG. 1: Various setups of two-slit interferometry (a) with the path 2 containing a usual external bath denoted by the grey circular region, (b) with no bath but delay, (c) with two paths containing the same nonlinear regions denoted by the grey boxes. (d) The same as (c) with delay in one of the paths after (A) and before (B) the interaction within the nonlinear region.

and the grating was assumed to correctly explain experimental results [10]. Since C_{70} molecule is too small to see the anharmonic shape of the external potential the coupling between the CM and the ID is hardly expected. One can still insist, however, that not only if the slit wall contains more rapidly varying repulsive potential but also if the size of the object becomes larger, for example by using an insulin, then the coupling term might manifest itself. In this paper we consider such situation that the coupling between CM and ID is non-negligible. It must be mentioned that we ignore all the other sources of decoherences except those originating from the ID for our discussion, but will briefly comment it. We also note that neither we derive any new formula nor estimate any values of this effect. We would like to show basic physical mechanism and possibility of the phenomena.

II. REVIEW OF DECOHERENCE

First, we briefly review the decoherence from an external environment. Figure 1(a) shows usual two path interferometry, where one of the paths interacts with a "bath" of particles, i.e. an environment. One can write an initial total wavefunction in the following way

$$|0\rangle = \frac{1}{\sqrt{2}} [|1(0;X)\rangle + |2(0;X)\rangle] \quad |0; \rangle; \quad (2)$$

where $|1(2)\rangle$ and $|0; \rangle$ denote a wavefunction of a system moving along the path 1(2) and that of a bath described

by variables \mathbf{r} , respectively. Under the dynamics, which includes interaction with the bath only along the path 2, the wavefunction becomes

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}} [|1(t;X)\rangle |1(t; \mathbf{r})\rangle + |2(t;X)\rangle |2(t; \mathbf{r})\rangle]; \quad (3)$$

where $|1(2)\rangle$ is a state of the bath with the system going around the path 1(2). Here, we assume the coupling with the bath is small enough to have no influence on the system, but changes only a state of the bath [17, 18].

From the reduced density matrix the interference term is given as

$$\text{Re} \int d\mathbf{r} |1(t;X)\rangle \langle 2(t;X)| \rho_{12}(\mathbf{r}) : \quad (4)$$

The physical meaning of Eq. (4) is obvious. The first term contains usual information of the interference pattern of the system going through the two paths. The second term corresponds to the visibility which measures the decoherence. Eq. (4) allows one to interpret the reduction of the interference in terms of a reduction in the overlap of the bath states for the two paths. Stern, Aharony and Imry [17] argue that one can make the identification

$$\rho_{12}(\mathbf{r}) = \int d\mathbf{r}' |1(t; \mathbf{r}')\rangle \langle 2(t; \mathbf{r}')|; \quad (5)$$

where $\rho_{12}(\mathbf{r}) = \int d\mathbf{r}' |1(t; \mathbf{r}')\rangle \langle 2(t; \mathbf{r}')|$, and $\rho_{12}(\mathbf{r}) = \int d\mathbf{r}' |1(t; \mathbf{r}')\rangle \langle 2(t; \mathbf{r}')|$. Here V_I denotes the interaction potential between the system and the environment in the interaction picture, i.e. $V_I = \exp(iH_{env}t)V(X; \mathbf{r})\exp(-iH_{env}t)$ in which H_{env} and V represent the Hamiltonian of the environment and its interaction with the system, respectively. Eq. (5) implies that the reduction of the interference can also be ascribed to the accumulating phase uncertainty of the system on the interacting path being subject to an uncertain potential. In this sense decoherence is often referred to as "dephasing".

III. DECOHERENCE FROM INTERNAL DYNAMICS

A. Two slit interferometry

Now let us consider two path interferometry of a macroscopic object with its complicated ID. The CM and the ID now play roles of a system and an environment, respectively. We take into account the case that the object is moving freely as shown in Fig. 1(b), so that there is no entanglement between the CM and the ID. The motion of the CM is easily described by a plane wave, $|0(t;X)\rangle$. The CM of the object then shows perfect coherence since the CM dynamics is completely isolated. When one increases the path length of one of the two paths by an amount of l , clear interference pattern is expected from the term

$\psi_1(t; X) \psi_2(t; X) = \psi_0(t + t; X) \psi_0(t; X)$ between two CM states, where $t = L/v$ (v is the velocity of the CM). The final wavefunction is given as

$$\psi(t) = \frac{1}{\sqrt{2}} [\psi_0(t + t; X) + \psi_0(t; X)] \quad (t;) : \quad (6)$$

The time delay t is only applicable to the CM since the ID independently evolves in time. The overlap of the ID always yields that of the same states, i.e. complete coherence. Without entanglement with the ID the decoherence of the CM cannot be generated.

Let us consider now the case an object goes through nonlinear external potentials along the paths as shown in Fig. 1(c). We assume these nonlinear potentials are equivalent for the two paths. In a two slit interference, in general, the two slits are made to have the same geometry. Even though the CM motion is entangled with the ID during the passage through such nonlinear region, this does not generate any decoherence. The reason is that the evolution of the ID is always equal for the two paths, so that $\psi_1(t;) = \psi_2(t;)$, i.e. $\langle \psi_1(t;) | \psi_2(t;) \rangle = 1$. The entanglement with internal environment makes the phases of each of the partial waves of the CM uncertain in viewpoint of Eq. (5), but does not alter the relative phase. In Fig. 1(d), we introduce additional delay for one of the paths. When the delay is given after the passage through the nonlinear region, the situation exactly corresponds to the case described in Fig. 1(b). The only difference is that one starts not with an initial ID state $\psi_1(0;) = \psi_2(0;)$ but with $\psi_1(t_0;) = \psi_2(t_0;)$, where t_0 is the time when the object departs from the nonlinear region. When the delay is given before, one can see that the situation is also the same as the case shown in Fig. 1(b) by considering the argument related to Eq. (6).

It should be noted that in usual decoherence generated from an outer environment it is not easy to find the case that the two paths have the same environment. If the decoherence occurs mainly by interacting with other particles, the system going through two paths accumulates different random phases from scattering with different particles of different states. The system going through the two paths thus see different environments. This is the reason why the decoherence from an outer environment has been dealt with the setup shown in Fig. 1(a). One example that the same environment is applied to the two interfering waves is the interaction of an interfering electron with zero-point (or vacuum) fluctuation, where the electron does not decohere [19, 20, 21].

B. Quantification of decoherence from internal dynamics

In the above discussion it has been shown that it is not easy to see the decoherence generating from the ID with usual simple geometry of interference. The only way to observe the decoherence from the coupling with ID is that the ID's should see different nonlinear interactions

for each path. Without loss of generality this situation can be represented as the case that only one path contains the nonlinear region. The overlap of the ID can then be written as

$$\langle \psi(0;) | \psi(t;) \rangle = \int \psi_0^*(t;) \psi(t;) dt = \int \psi_0^*(t;) \exp(i \int_0^t (H_{in} + V) dt) \psi_0(t;) dt \quad (7)$$

where t and \hat{T} denote the interaction time within the nonlinear potential and the time ordering operator, respectively. A symmetric geometry of interference is sometimes useful, for example, measurement of phase of the transmission coefficient through a quantum dot [22], where a quantum dot is plugged into one of the arms of an Aharonov-Bohm ring. From the Aharonov-Bohm oscillation one can determine the phase shift of electron passing through the quantum dot.

The quantity given in Eq. (7) is known as so called fidelity [23]. The decay of such a quantity determines the decoherence rate. One important remark is that even for the ID with a few degrees of freedom the fidelity decays exponentially if its dynamics is chaotic [24, 25]. It opens possibility that the decoherence can occur in molecules consisting of even several atoms from entanglement with its ID [15] (See also [26, 27, 28, 29, 30, 31] for decoherence generated from chaos) in a certain special condition. When a single coherent state, a minimum wavepacket, is chosen as an initial state of ID, one can expect golden rule or Lyapunov decay depending on the strength of the coupling [32], in which the fidelity decay does not much depend on the initial condition for a given energy as far as the completely chaotic dynamics is concerned for the ID. This situation corresponds to the ID governed by rather small number of degrees of freedom at low temperature. As we mentioned in the beginning, however, for small molecules it is hard to expect the coupling between the CM and the ID. To see this effect in the system with small degrees of freedom we need something different from usual molecules.

For rather bigger systems, which we are interested in, it is not easy to directly calculate how much the interaction between the CM and the ID make an influence on the state of the ID because the internal degrees of freedom consist of many particles. First let us consider the zero temperature case. If the interaction is strong enough to generate any kind of elementary excitations such as phonons, charge density waves, magnons, ferromagnetic systems, and so on, then the CM will lose his coherence completely. Nothing happens for the system going through the path 1 in Fig. 1, while the state of the ID through the path 2 in Fig. 1 is changed from the ground state to the excited state of the elementary excitation. By checking the state of the ID one can see which path the system go through. It is nothing but a complete decoherence. In this case it is crucial to know whether the excitation is gapless or not. At finite temperature the situation is much more complicated. In the beginning we assumed that there is no other sources of decoherence




FIG. 2: (a) Two time reversal paths, χ_1 and χ_2 , multiply reflecting from random scatterers in a disordered system. (b) Simple example with three scatterers labeled by 0, 1, and 2. U_n denotes an unitary operator freely propagating between two scatterers.

except those originating from the ID. Unfortunately this is no longer true since an object with finite temperature is always coupled with outer environment by emitting black body radiation or possibly other radiation from thermal vibration. It is another issue how much this effect degrades the coherence of the motion of the CM. Now one needs to compare the decoherence from the coupling with the outer environment and that with the ID. If the radiation is not so harmful for the coherence, e.g. the case that the wavelength of the radiation is much larger than the difference of two paths, it is meaningful to calculate Eq. (7). This calculation can be done by using the well theoretical technique once the time dependent interaction is known. Surely it is still hard task. We do not want to calculate it in this paper, but only point out the possible existence of the decoherence generating from the coupling with the ID.

Another remark on Eq. (7) is its physical interpretation in terms of quantum measurement. Here, the total system including CM and ID is isolated from the external world. The decoherence is generated from entanglement only within a macroscopic object itself. In this sense no external observer exists. No information is transferred from the object to outside. Since a system (the CM) does not know the difference between an internal and an outer environment, the ID plays a role of an observer. Information thus flows from the CM to the ID. Following Zurek's argument [3], one can say the ID is watching the CM.

Finally, let us consider one interesting example, weak localization in disordered systems [33]. Two time reversal paths multiply reflecting from the random scatterers leads to the localization of a wavefunction due to their constructive interference as shown in Fig. 2 (a). Such localization is fragile for both the decoherence and the perturbation breaking time reversal symmetry. A random potential generates complicated and chaotic dynamics, which can give rise to the entanglement between CM and ID. At first glance the decoherence from the ID is hardly expected to arise since the total system has time reversal symmetry. One might think that the state of the ID of the CM rotating clockwise must be the same as that of counter-clockwise. It can be shown, however, that the coupling to the ID can destroy the weak localization by generating the decoherence from the ID.

To prove the appearance of the decoherence let us consider the overlap between two ID, namely χ_1 and χ_2 , of the time reversal paths of CM after a round trip along a closed loop. Since we are interested only in the ID, the influence from its coupling to the CM can be regarded as a time dependent external perturbation, i.e. $(X_{1(2)}(t); \chi) = \chi_{1(2)}(t; \chi)$. Note $\chi_{1(2)}(t + \tau; \chi) = \chi_{1(2)}(t; \chi)$, where τ denotes the duration time taken for a round trip around the closed loop. The final state $\chi_{1(2)}(\tau)$ is then given as

$$\chi_{1(2)}(\tau; \chi) = \hat{T} \exp \left[-i \int_0^\tau dt H_{in}(\chi) + X_{1(2)}(t; \chi) \right] \chi_0 : \quad (8)$$

Even though $X_1(t) = X_2(\tau - t)$ holds, i.e. $\chi_1(t; \chi) = \chi_2(\tau - t; \chi)$, one finds $\chi_1(\tau) \notin \chi_2(\tau)$ due to existence of the time ordering operator \hat{T} . To make it more clear, let us consider a simple example: three scatterers well localized in space as shown in Fig. 2 (b). During free propagation between two scatterers the CM is decoupled from the ID. We assume the process of collision with the scatterers is short enough to be described by delta-function in time. The interaction term V_1 for the clockwise propagation can then be given as

$$V_1(t; \chi) = f_1(\chi)(t - t_1) + f_2(\chi)(t - t_2); \quad (9)$$

where t_1 and t_2 are the collision times upon the first and the second scatterer, respectively, and $0 < t_1 < t_2 < \tau$. After one round trip, the states of the ID for the clockwise and the counter-clockwise χ_1 and χ_2 are respectively given as

$$\begin{aligned} \chi_1(\tau) &= U_3 K_2 U_2 K_1 U_1 \chi_0 \\ \chi_2(\tau) &= U_1 K_1 U_2 K_2 U_3 \chi_0; \end{aligned} \quad (10)$$

where by using the eigenstates of H_0 ($H_0 |j\rangle = E_j |j\rangle$) one obtains $U_{n,ij} = \exp[iE_i t_n] \delta_{ij}$, and $K_{n,ij} = \delta_{ij} \exp[i\phi_n(\chi)] \delta_{ij}$. Here, $t_1 = t_1$, $t_2 = t_2 - t_1$, and $t_3 = \tau - t_2$. It is obvious that in general $U_3 K_2 U_2 K_1 U_1 \chi_0 \notin U_1 K_1 U_2 K_2 U_3 \chi_0$ since K_1 and K_2 are not

diagonal. Consequently one can find that in general $\langle |j_1\rangle\langle j_2| \rangle_i < 1$. The weak localization of the CM of a macroscopic object in disordered potentials can be destroyed due to the coupling to the ID.

IV. SUMMARY

In summary, we have investigated the decoherence generated from the internal dynamics of a macroscopic object. In a usual setup of two slit interferometry, it is hard to expect the appearance of such decoherence. Only asymmetric geometry of the interfering paths containing anharmonic external potential allows one to observe the decoherence from the internal dynamics. Such decoherence can then be measured by the fidelity given in

Eq. (7). In this case, the internal degrees of freedom of a macroscopic object are watching its center of mass motion. The weak localization of the center of mass motion of a macroscopic object in disordered potentials can also be destroyed by such decoherence without any external perturbation breaking time reversal symmetry.

Acknowledgments

SWK would like to thank Fritz Haake and Klaus Homberger for helpful discussions. The part of this work has been done when SWK has participated in the focus program of Asia Pacific Center for Theoretical Physics (APCTP) in Pohang, Korea.

-
- [1] W. H. Zurek, *Phys. Today* 44 (10), 36 (1991).
 - [2] E. Joos, in *Decoherence: Theoretical, Experimental and Conceptual Problems* edited by P. Blanchard, D. Giulini, E. Joos, and I.-O. Stamatescu, *Lecture Notes in Physics* Vol. 538 (Springer-Verlag, Heidelberg, 2000) p.1.
 - [3] W. H. Zurek, *Rev. Mod. Phys.* 75, 715 (2003).
 - [4] S. Haroche, *Phys. Today* 51 (7), (1998).
 - [5] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* 77, 4887 (1996).
 - [6] C. Monroe, D. M. Meekhof, B. E. King, and D. J. Wineland, *Science* 272, 1131 (1996).
 - [7] C. J. Myatt, B. E. King, Q. A. Turchette, C. A. Sackett, D. Kielpinski, W. M. Itano, and D. J. Wineland, *Nature* 403, 269 (2000).
 - [8] M. Amdt, O. Nariz, J. Vos-Andrae, C. Keller, G. van der Zouw, and A. Zellinger, *Nature* 401, 680 (1999).
 - [9] O. Nariz, B. Brezger, M. Amdt, and A. Zellinger, *Phys. Rev. Lett.* 87, 160401 (2001).
 - [10] B. Brezger, L. Hackemüller, S. Uttenthaler, J. Petschinka, M. Amdt, and A. Zellinger, *Phys. Rev. Lett.* 88, 100404 (2002).
 - [11] K. Homberger, S. Uttenthaler, B. Brezger, L. Hackemüller, M. Amdt, and A. Zellinger, *Phys. Rev. Lett.* 90, 160401 (2003).
 - [12] L. Hackemüller, S. Uttenthaler, K. Homberger, E. Reiger, B. Brezger, A. Zellinger, and M. Amdt, *Phys. Rev. Lett.* 91, 090408 (2003).
 - [13] L. Hackemüller, K. Homberger, B. Brezger, A. Zellinger, and M. Amdt, *Nature* 427, 711 (2004).
 - [14] A. R. K. Klovsky, *Europhys. Lett.* 27, 79 (1994).
 - [15] H.-K. Park and S. W. Kim, *Phys. Rev. A* 67, 060102(R) (2003).
 - [16] R. E. Grisenti, W. Schollkopf, J. P. Toennies, G. C. Hegerfeldt, and T. Kohler, *Phys. Rev. Lett.* 83, 1755 (1999).
 - [17] A. Stern, Y. Aharony, and Y. Imry, *Phys. Rev. A* 41, 3436 (1990).
 - [18] G. A. Fiete and E. J. Heller, *Phys. Rev. A* 68, 022112 (2003).
 - [19] N. Kumar, D. V. Baxter, R. Richter, and J. O. Stom Olsen, *Phys. Rev. Lett.* 59, 1853 (1987).
 - [20] J. Rammer, A. L. Shelankov, and A. Schmid, *Phys. Rev. Lett.* 60, 1985 (1988); G. Bergman, *Phys. Rev. Lett.* 60, 1986 (1988).
 - [21] Y. Imry, cond-mat/0202044 (unpublished).
 - [22] R. Schuster, E. Buks, M. Heiblum, D. Mahalu, V. Umansky, and H. Shtrikman, *Nature* 385, 417 (1997).
 - [23] H. M. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
 - [24] A. Peres, *Phys. Rev. A* 30, 1610 (1984).
 - [25] R. A. Jalabert and H. M. Pastawski, *Phys. Rev. Lett.* 86, 2490 (2001).
 - [26] S. Adachi, M. Toda, and K. Ikeda, *Phys. Rev. Lett.* 61, 659 (1988).
 - [27] A. R. Klovsky, *Phys. Rev. E* 50, 3569 (1994).
 - [28] H. Kubotani, T. Okamura, M. Sakagami, *Physica A* 214, 560 (1995).
 - [29] H. Nakazato, M. Namiki, S. Pascazio, and Y. Yamataka, *Phys. Lett. A* 222, 130 (1996).
 - [30] A. Tanaka, H. Fujisaki, and T. Miyadera, *Phys. Rev. E* 66, 045201(R) (2002).
 - [31] D. Cohen and T. Kottos, *Phys. Rev. E* 69, 055201(R) (2004).
 - [32] Ph. Jacquod, P. G. Silvestrov, and C. W. J. Beenakker, *Phys. Rev. E* 64, 055203(R) (2001).
 - [33] G. Bergman, *Phys. Rep.* 107, 1 (1984).