Comment on "A mathematical theorem as a basis for the second law: Thomson's

formulation applied to equilibrium"

by

Dan Solomon

Rauland-Borg Corporation 3450 W. Oakton Skokie, IL USA

Email: dan.solomon@rauland.com March 3, 2004

Abstract

A. E. Allahverdyan and Th . M. Nieuwenhuizen [1] in their paper "A mathematical theorem as a basis for the second law: Thomson's formulation applied to equilibrium" present a proof of the second law of thermodynamics based on quantum mechanics. In this comment on their paper I offer a counterexample to their proof.

In their paper [1] A. E. Allahverdyan and Th . M. Nieuwenhuizen present a proof of the second law of thermodynamics as a rigorous theorem of quantum mechanics. Allahverdyan and Nieuwenhuizen consider a closed statistical system . At t=0 the initial Hamiltonian is \hat{H}_0 and the initial equilibrium state is a Gibbs distribution so that,

$$\hat{\rho}(0) = \frac{e^{-\beta \hat{H}_0}}{Z}; \quad Z = \text{tr}e^{-\beta \hat{H}_0}$$
 (1)

where $\beta = 1/T$ and T is the temperature. At time t = 0 an external interaction is turned on and then turned off at time t. W is the work put into the system during this interaction and is given by [1],

$$W = tr[\hat{H}_0 \hat{V} \hat{\rho}(0) \hat{V}^{\dagger}] - tr[\hat{H}_0 \hat{\rho}(0)]$$
 (2)

where \hat{V} is a unitary operator that describes the interaction. Using the mathematical identity tr[AB] = tr[BA] this can be rewritten as,

$$W = tr[\hat{V}^{\dagger} \hat{H}_0 \hat{V} \hat{\rho}(0)] - tr[\hat{H}_0 \hat{\rho}(0)]$$
(3)

Allahverdyan and Nieuwenhuizen claim that W must be non-negative for all possible interactions. Since the interaction is described by \hat{V} this means that W must be non-negative for all possible unitary operators \hat{V} . I will give an example where this is not the case.

I will examine this problem for a quantized fermion field interacting with a quantized electromagnetic field in the temporal gauge. In the temporal gauge the gauge condition is given by the relationship $A_0 = 0$ [2,3,4,5] where A_0 is the scalar component of the electric potential. We will work in the Schrödinger picture and use natural units so

that $\hbar=c=1$. In the Schrödinger picture the field operators are time independent and the time dependence of the quantum system is reflected in the state vectors $|\Omega(t)\rangle$. The Hamiltonian is given by,

$$\hat{\mathbf{H}}_{0} = \hat{\mathbf{H}}_{0,D} + \hat{\mathbf{H}}_{0,M} - \int \hat{\mathbf{J}}(\mathbf{x}) \cdot \hat{\mathbf{A}}(\mathbf{x}) d\mathbf{x}$$
(4)

where,

$$\hat{\mathbf{H}}_{0,D} = \frac{1}{2} \int \left[\hat{\mathbf{\psi}}^{\dagger} (\mathbf{x}), \mathbf{H}_{0,D} \hat{\mathbf{\psi}} (\mathbf{x}) \right] d\mathbf{x}; \quad \mathbf{H}_{0,D} = -i\boldsymbol{\alpha} \cdot \nabla + \beta \mathbf{m}$$
 (5)

$$\hat{\mathbf{H}}_{0,\mathbf{M}} = \frac{1}{2} \int (\hat{\mathbf{E}}^2 + \hat{\mathbf{B}}^2) d\mathbf{x}; \qquad \hat{\mathbf{B}}(\mathbf{x}) = \nabla \times \hat{\mathbf{A}}(\mathbf{x})$$
 (6)

$$\hat{\mathbf{J}}(\mathbf{x}) = \frac{\mathbf{q}}{2} \left[\hat{\mathbf{v}}^{\dagger}(\mathbf{x}), \alpha \hat{\mathbf{v}}(\mathbf{x}) \right] \tag{7}$$

In the above expressions m is the fermion mass, α and β are the usual 4x4 matrices, q is the electric charge, $\hat{H}_{0,D}$ is the Dirac Hamiltonian, $\hat{H}_{0,M}$ is the Hamiltonian for the electromagnetic field, and $\hat{J}(x)$ is the current operator. The Schrödinger picture time independent fermion field operators are $\hat{\psi}(x)$ and $\hat{\psi}^{\dagger}(x)$. The Schrödinger picture time independent field operators for the electromagnetic field are $\hat{A}(x)$ and $\hat{E}(x)$. The electromagnetic field operators are real so that $\hat{A}^{\dagger}(x) = \hat{A}(x)$ and $\hat{E}^{\dagger}(x) = \hat{E}(x)$.

The field operators obey the following relationships [3,4],

$$\left[\hat{A}^{i}(\mathbf{x}), \hat{E}^{j}(\mathbf{y})\right] = -i\delta_{ij}\delta^{3}(\mathbf{x} - \mathbf{y}); \left[\hat{A}^{i}(\mathbf{x}), \hat{A}^{j}(\mathbf{y})\right] = \left[\hat{E}^{i}(\mathbf{x}), \hat{E}^{j}(\mathbf{y})\right] = 0 \quad (8)$$

and

$$\left\{\hat{\psi}_{a}^{\dagger}(\mathbf{x}), \hat{\psi}_{b}(\mathbf{y})\right\} = \delta_{ab}\delta(\mathbf{x} - \mathbf{y}); \left\{\hat{\psi}_{a}^{\dagger}(\mathbf{x}), \hat{\psi}_{b}^{\dagger}(\mathbf{y})\right\} = \left\{\hat{\psi}_{a}(\mathbf{x}), \hat{\psi}_{b}(\mathbf{y})\right\} = 0$$
 (9)

where "a" and "b" are spinor indices. In addition, all commutators between the electromagnetic field operators and fermion field operators are zero, i.e.,

$$\left[\hat{\mathbf{A}}(\mathbf{x}), \hat{\mathbf{\psi}}(\mathbf{y})\right] = \left[\hat{\mathbf{E}}(\mathbf{x}), \hat{\mathbf{\psi}}(\mathbf{y})\right] = \left[\hat{\mathbf{A}}(\mathbf{x}), \hat{\mathbf{\psi}}^{\dagger}(\mathbf{y})\right] = \left[\hat{\mathbf{E}}(\mathbf{x}), \hat{\mathbf{\psi}}^{\dagger}(\mathbf{y})\right] = 0$$
(10)

Define the quantity,

$$\hat{G}(\mathbf{x}) = \nabla \cdot \hat{\mathbf{E}}(\mathbf{x}) - \hat{\rho}(\mathbf{x}) \tag{11}$$

where the current operator $\hat{\rho}(\mathbf{x})$ is defined by,

$$\hat{\rho}(\mathbf{x}) = \frac{\mathbf{q}}{2} \left[\hat{\psi}^{\dagger}(\mathbf{x}), \hat{\psi}(\mathbf{x}) \right]$$
 (12)

All valid state vectors $|\Omega\rangle$ must satisfy the Gauss constraint [2],

$$\hat{\mathbf{G}}(\mathbf{x})|\Omega\rangle = 0 \tag{13}$$

The unitary operator \hat{V} converts the state vector $|\Omega\rangle$ into the state vector $\hat{V}|\Omega\rangle$. The Gauss constraint must be still be satisfied so that in the temporal gauge \hat{V} is restricted to those operators which satisfy,

$$\hat{G}(\mathbf{x})\hat{V}|\Omega\rangle = 0$$
 for all $|\Omega\rangle$ for which $\hat{G}(\mathbf{x})|\Omega\rangle = 0$ (14)

I will show that for the Hamiltonian operator defined in (4) it is possible to find a unitary operator \hat{V} which satisfies the Gauss constraint for which W of equation (3) is negative. This is contrary to the assertion of reference [1].

Now let,

$$\hat{\mathbf{V}} = \hat{\mathbf{U}}\hat{\mathbf{R}} \tag{15}$$

where \hat{U} and \hat{R} are unitary operators. Let \hat{U} satisfy the Gauss constraint (14) and define \hat{R} by,

$$\hat{R} = e^{i\hat{U}^{\dagger}\hat{C}\hat{U}} \tag{16}$$

where

$$\hat{\mathbf{C}} = \int \hat{\mathbf{E}} \cdot \nabla \chi d\mathbf{x} \tag{17}$$

where $\chi(\vec{x})$ is an arbitrary real valued function. Note that $\hat{C} = \hat{C}^{\dagger}$ so that $\hat{R}^{\dagger} = e^{-i\hat{U}^{\dagger}\hat{C}\hat{U}}$ therefore $\hat{R}^{\dagger}\hat{R} = 1$ so that \hat{R} is unitary.

Using the commutator relations we obtain,

$$\left[\hat{\mathbf{H}}_{0,D},\hat{\mathbf{C}}\right] = \left[\hat{\mathbf{J}}(\mathbf{x}),\hat{\mathbf{C}}\right] = \left[\rho(\mathbf{x}),\hat{\mathbf{C}}\right] = 0 \tag{18}$$

and,

$$\left[\hat{\mathbf{A}}(\mathbf{x}),\hat{\mathbf{C}}\right] = -i\nabla\chi(\mathbf{x}) \tag{19}$$

Use this result to obtain,

$$\left[\hat{\mathbf{B}}(\mathbf{x}), \hat{\mathbf{C}}\right] = \nabla \times \left[\hat{\mathbf{A}}(\mathbf{x}), \hat{\mathbf{C}}\right] = -i\nabla \times \nabla \chi(\mathbf{x}) = 0$$
(20)

Therefore,

$$\left[\hat{H}_{0,M},\hat{C}\right] = 0 \tag{21}$$

From the above we obtain,

$$\left[\hat{G}(\mathbf{x}),\hat{C}\right] = 0 \tag{22}$$

Next we want to determine if $\hat{V} = \hat{U}\hat{R}$ satisfies the Gauss constraint. To do this we use the obvious relationship,

$$\left[\hat{\mathbf{U}}^{\dagger}\hat{\mathbf{O}}_{1}\hat{\mathbf{U}},\hat{\mathbf{U}}^{\dagger}\hat{\mathbf{O}}_{2}\hat{\mathbf{U}}\right] = \hat{\mathbf{U}}^{\dagger}\left[\hat{\mathbf{O}}_{1},\hat{\mathbf{O}}_{2}\right]\hat{\mathbf{U}} \tag{23}$$

where \hat{O}_1 and \hat{O}_2 are operators. Use this in (22) to obtain,

$$\left[\hat{\mathbf{U}}^{\dagger}\hat{\mathbf{G}}(\mathbf{x})\hat{\mathbf{U}},\hat{\mathbf{U}}^{\dagger}\hat{\mathbf{C}}\hat{\mathbf{U}}\right] = \hat{\mathbf{U}}^{\dagger}\left[\hat{\mathbf{G}}(\mathbf{x}),\hat{\mathbf{C}}\right]\hat{\mathbf{U}} = 0 \tag{24}$$

Therefore $\hat{V}=\hat{U}\hat{R}$ satisfied Gauss constraint since \hat{U} has been assumed to satisfy this constraint and $\hat{U}^{\dagger}\hat{C}\hat{U}$, and thereby \hat{R} , commutes with $\hat{U}^{\dagger}\hat{G}(x)\hat{U}$.

Next we want to evaluate $\,\hat{V}^{\dagger}\hat{H}_{0}\hat{V}\,.\,$ Use (4) to obtain,

$$\hat{\mathbf{V}}^{\dagger} \hat{\mathbf{H}}_{0} \hat{\mathbf{V}} = \hat{\mathbf{V}}^{\dagger} \left(\hat{\mathbf{H}}_{0,D} + \hat{\mathbf{H}}_{0,M} - \int \hat{\mathbf{J}} \left(\mathbf{x} \right) \cdot \hat{\mathbf{A}} \left(\mathbf{x} \right) d\mathbf{x} \right) \hat{\mathbf{V}}$$
 (25)

We will require the Baker-Campell-Hausdorff relationships [6] which states that,

$$e^{+\hat{O}_1}\hat{O}_2e^{-\hat{O}_1} = \hat{O}_2 + \left[\hat{O}_1, \hat{O}_2\right] + \frac{1}{2}\left[\hat{O}_1, \left[\hat{O}_1, \hat{O}_2\right]\right] + \dots$$
 (26)

Use this along with (18), (21), and (24) to obtain,

$$\hat{\mathbf{V}}^{\dagger} \left(\hat{\mathbf{H}}_{0,D} + \hat{\mathbf{H}}_{0,M} \right) \hat{\mathbf{V}} = \hat{\mathbf{U}}^{\dagger} \left(\hat{\mathbf{H}}_{0,D} + \hat{\mathbf{H}}_{0,M} \right) \hat{\mathbf{U}}$$
 (27)

Next we obtain,

$$\hat{\mathbf{V}}^{\dagger} \left(\hat{\mathbf{J}} (\mathbf{x}) \cdot \hat{\mathbf{A}} (\mathbf{x}) \right) \hat{\mathbf{V}} = \left(\hat{\mathbf{V}}^{\dagger} \hat{\mathbf{J}} (\mathbf{x}) \hat{\mathbf{V}} \right) \cdot \left(\hat{\mathbf{V}}^{\dagger} \hat{\mathbf{A}} (\mathbf{x}) \hat{\mathbf{V}} \right)$$
(28)

Use (10), (19), (24), and (26) to yield,

$$\left(\hat{\mathbf{V}}^{\dagger}\hat{\mathbf{J}}(\mathbf{x})\hat{\mathbf{V}}\right) = \left(\hat{\mathbf{U}}^{\dagger}\hat{\mathbf{J}}(\mathbf{x})\hat{\mathbf{U}}\right) \tag{29}$$

and,

$$(\hat{\mathbf{V}}^{\dagger} \hat{\mathbf{A}}(\mathbf{x}) \hat{\mathbf{V}}) = \hat{\mathbf{U}}^{\dagger} (\hat{\mathbf{A}}(\mathbf{x}) - \nabla \chi(\mathbf{x})) \hat{\mathbf{U}}$$
(30)

Use these results in (28) to obtain

$$\hat{\mathbf{V}}^{\dagger} (\hat{\mathbf{J}}(\mathbf{x}) \cdot \hat{\mathbf{A}}(\mathbf{x})) \hat{\mathbf{V}} = (\hat{\mathbf{U}}^{\dagger} \hat{\mathbf{J}}(\mathbf{x}) \hat{\mathbf{U}}) \cdot \hat{\mathbf{U}}^{\dagger} (\hat{\mathbf{A}}(\mathbf{x}) - \nabla \chi(\mathbf{x})) \hat{\mathbf{U}}$$

$$= \hat{\mathbf{U}}^{\dagger} (\hat{\mathbf{J}}(\mathbf{x}) \cdot (\hat{\mathbf{A}}(\mathbf{x}) - \nabla \chi(\mathbf{x}))) \hat{\mathbf{U}}$$
(31)

Use this and (27) in (25) to obtain,

$$V^{\dagger} \hat{H}_{0} V = \hat{U}^{\dagger} \left(\hat{H}_{0} - \int \hat{\mathbf{J}}(\mathbf{x}) \cdot \nabla \chi(\mathbf{x}) d\mathbf{x} \right) \hat{U}$$
(32)

Use this in (3) to yield,

$$W = W_0 - \int tr \left[\hat{U}^{\dagger} \hat{\mathbf{J}}(\mathbf{x}) \hat{U} \hat{\rho}(0) \right] \cdot \nabla \chi(\mathbf{x}) d\mathbf{x}$$
(33)

where,

$$W_0 = \operatorname{tr}[\hat{U}^{\dagger} \hat{H}_0 \hat{U} \hat{\rho}(0)] - \operatorname{tr}[\hat{H}_0 \hat{\rho}(0)]$$
(34)

Integrate the second term in (33) with respect to parts and assume reasonable boundary conditions to obtain,

$$W = W_0 + \int \chi(\mathbf{x}) \nabla \cdot \hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x}) d\mathbf{x}$$
(35)

where $\hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x})$ is defined by,

$$\hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x}) = \operatorname{tr}[\hat{\mathbf{U}}^{\dagger}\hat{\mathbf{J}}(\mathbf{x})\hat{\mathbf{U}}\hat{\boldsymbol{\rho}}(0)] \tag{36}$$

Note that W_0 and the quantity $\hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x})$ are independent of $\chi(\mathbf{x})$. Therefore $\chi(\mathbf{x})$ can be varied in an arbitrary manner without affecting W_0 or $\hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x})$. Assume that $\hat{\mathbf{U}}$ is chosen so that $\nabla \cdot \hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x})$ is non-zero. In this case we can always find a $\chi(\mathbf{x})$ which makes W a negative number. For example let $\chi(\mathbf{x}) = -\lambda \nabla \cdot \hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x})$ where λ is a constant. In this case (35) becomes,

$$W = W_0 - \lambda \int \left(\nabla \cdot \hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x}) \right)^2 d\mathbf{x}$$
(37)

Therefore, since $\nabla \cdot \hat{\mathbf{J}}_{\hat{\mathbf{U}}}(\mathbf{x})$ is non-zero the quantity under the integral must be positive. Therefore for sufficiently large λ the work W will be negative.

This result depends on the assumption that we can find a unitary transformation \hat{U} such that $\nabla \cdot \hat{\mathbf{J}}_{\hat{U}}(\mathbf{x})$ is non-zero. How do we know that this can be done? $\hat{\mathbf{J}}_{\hat{U}}(\mathbf{x})$ is the current averaged over the distribution after an interaction associated with the unitary operator \hat{U} . Therefore we must have an interaction the takes the average current from an initial equilibrium state, where the divergence of the average current is zero, to some non-equilibrium state where $\nabla \cdot \hat{\mathbf{J}}_{\hat{U}}(\mathbf{x})$ is non-zero. Since there are many interaction that can take a system into a non-equilibrium state it is reasonable to assume that a \hat{U} exists.

In conclusion a counterexample has been found to the proof presented in [1] that the quantity W in equation (2) must be non-negative for any unitary operator \hat{V} . The counterexample is for a quantized fermion field coupled to a quantized electromagnetic field in the temporal gauge. It is shown that in this case a unitary operator \hat{V} exists for which W is negative.

References

- 1. A.E. Allahverdyan and Th. M. Nieuwenhuizen, Physica A, **305**, 542 (2002). See also arXiv:cond-mat/0110422.
- 2. G. Leibbrandt, Rev of Modern Phys, 59, 1067 (1987).
- 3. M. Creutz, Ann. Phys., **117**, 471 (1979)
- 4. B. Hatfield. "Quantum Field Theory of Point Particles and Strings", Addison-Wesley, Reading, Massachusetts (1992).
- 5. C. Kiefer and A. Wiepf, Annals Phys., 236, 241 (1994).
- 6. W. Greiner and J. Reinhardt. "Field Quantization", Springer-Verlag, Berlin (1996).