

Global entanglement and quantum criticality in spin chains

Tzu-Chieh Wei, Dyutiman Das, Swagatam Mukhopadhyay, Smitha Vishveshwara, and Paul M. Goldbart

Department of Physics, University of Illinois at Urbana-Champaign,
1110 West Green Street, Urbana, Illinois 61801-3080, U.S.A.

(Dated: May 26, 2004)

Entanglement in quantum XY spin chains of arbitrary length is investigated via a recently-developed *global* measure suitable for generic quantum many-body systems. The entanglement surface is determined over the phase diagram, and found to exhibit structure richer than expected. Near the critical line, the entanglement is peaked (albeit analytically), consistent with the notion that entanglement—the non-factorization of wave functions—reflects quantum correlations. Singularity does, however, accompany the critical line, as revealed by the divergence of the field-derivative of the entanglement along the line. The form of this singularity is dictated by the universality class controlling the quantum phase transition.

PACS numbers: 03.67.Mn, 03.65.Ud, 73.43.Nq, 05.70.Jk

Introduction—Quantum entanglement, a term coined by Schrödinger, has been recognized over the past decade as the central actor in many quantum information processing tasks, such as teleportation [1], dense coding [2], secure key distribution [3], and perhaps quantum computation [4]. More recently, entanglement has emerged as an actor on the nearby stage of quantum many-body physics, especially for systems that exhibit quantum phase transitions [5, 6, 7, 8], where it can play the role of a diagnostic of quantum correlations. Quantum phase transitions [9] are transitions between qualitatively distinct phases of quantum many-body systems, driven by quantum fluctuations. In view of the connection between entanglement and quantum correlations, one anticipates that entanglement will furnish a dramatic signature of the quantum critical point. From the viewpoint of quantum information, the more entangled a state, the more resources it is likely to possess. It is thus desirable to study and quantify the degree of entanglement near quantum phase transitions. By employing entanglement to diagnose many-body quantum states one may obtain fresh insight into the quantum many-body problem.

To date, progress in quantifying entanglement has taken place primarily in the domain of bipartite systems [10]. Much of the previous work on entanglement in quantum phase transitions has been based on bipartite measures, i.e., focus has been on entanglement either between pairs of parties [5, 6] or between a part and the remainder of a system [7]. For multipartite systems, however, the complete characterization of entanglement requires the consideration of multipartite entanglement, for which a consensus measure has not yet emerged.

Singular and scaling behavior of entanglement near quantum critical points was discovered in important work by Osterloh and co-workers [6], who invoked Wootters' *bipartite* concurrence [11] as a measure of entanglement. In the present letter, we apply a recently-developed *global* measure that provides a *holistic*, rather than bipartite, characterization of the entanglement of quantum many-body systems. Our focus is on one-dimensional spin systems, specifically ones that are exactly solvable and

exhibit quantum criticality. For these systems we are able to determine the entanglement analytically, and to observe that it varies in a singular manner near the quantum critical line. This supports the view that entanglement—the non-factorization of wave functions—reflects quantum correlations. Moreover, the boundaries between different phases can be detected by the entanglement.

Global measure of entanglement—To introduce a measure for characterizing the *global* entanglement, consider a general, n -partite, normalized pure state: $|\Psi\rangle = \sum_{p_1 \dots p_n} \Psi_{p_1 p_2 \dots p_n} |e_{p_1}^{(1)} e_{p_2}^{(2)} \dots e_{p_n}^{(n)}\rangle$. If the parties are all spin-1/2 then each can be taken to have the basis $\{|\uparrow\rangle, |\downarrow\rangle\}$. Our scheme for analyzing the entanglement involves considering how well an entangled state can be approximated by some unentangled (normalized) state (e.g. the state in which every spin points in a definite direction): $|\Phi\rangle \equiv \otimes_{i=1}^n |\phi^{(i)}\rangle$. The proximity of $|\Psi\rangle$ to $|\Phi\rangle$ is captured by their overlap; the entanglement of $|\Psi\rangle$ is revealed by the maximal overlap [12]

$$\Lambda_{\max}(\Psi) \equiv \max_{\Phi} |\langle \Phi | \Psi \rangle|; \quad (1)$$

the larger Λ_{\max} is, the less entangled is $|\Psi\rangle$. (Note that for a product state, Λ_{\max} is unity.) If the entangled state consists of two separate entangled pairs of subsystems, Λ_{\max} is the product of the maximal overlaps of the two. Hence, it makes sense to quantify the entanglement of $|\Psi\rangle$ via the following *extensive* quantity [13]

$$E_{\log_2}(\Psi) \equiv -\log_2 \Lambda_{\max}^2(\Psi), \quad (2)$$

This normalizes to unity the entanglement of EPR-Bell and N -party GHZ states, as well as gives zero for unentangled states. Finite- N entanglement is interesting in the context of quantum information processing. To characterize the properties of the quantum critical point we use the thermodynamic quantity \mathcal{E} defined by

$$\mathcal{E} \equiv \lim_{N \rightarrow \infty} \mathcal{E}_N, \quad \mathcal{E}_N \equiv N^{-1} E_{\log_2}(\Psi), \quad (3)$$

where \mathcal{E}_N is the *entanglement density*, i.e., the entanglement per particle.

Quantum XY spin chains and entanglement—We consider the family of models governed by the Hamiltonian

$$\mathcal{H}_{XY} = - \sum_{j=1}^N \left(\frac{1+r}{2} \sigma_j^x \sigma_{j+1}^x + \frac{1-r}{2} \sigma_j^y \sigma_{j+1}^y + h \sigma_j^z \right), \quad (4)$$

where r measures the anisotropy between x and y couplings, h is the transverse external field, lying along the z -direction, and we impose periodic boundary conditions. At $r = 0$ we have the isotropic XY limit (also known as the XX model) and at $r = 1$, the Ising limit. All anisotropic XY models ($0 < r \leq 1$) belong to the same universality class, i.e., the Ising class, whereas the isotropic XX model belongs to a different universality class. XY models exhibit three phases (see Fig. 1): oscillatory, ferromagnetic and paramagnetic. In contrast to the paramagnetic phase, the first two are ordered phases, with the oscillatory phase being associated with a characteristic wavevector, reflecting the modulation of the spin correlation functions (see, e.g., Ref. [14]). We shall see that the global entanglement detects the boundaries between these phases.

As is well known [9, 14, 15], the energy eigenproblem for the XY spin chain can be solved by a Jordan-Wigner transformation, via which the spins are recast

as fermions, followed by a Bogoliubov transformation, which diagonalizes the quadratic Hamiltonian. Having found the eigenstates, Λ_{\max} of Eq. (1) and hence the entanglement can be found. To do this, we parametrize the separable states via

$$|\Phi\rangle \equiv \bigotimes_{i=1}^N [\sin(\xi_i/2)|\uparrow\rangle_i + e^{i\phi_i} \cos(\xi_i/2)|\downarrow\rangle_i], \quad (5)$$

where $|\uparrow/\downarrow\rangle$ denote spin states parallel/antiparallel to the z -axis. Instead of maximizing the overlap with respect to the $2N$ real parameters $\{\xi_i, \phi_i\}$, for the lowest two states it is adequate to appeal to the translational symmetry and reality of the wave functions. Thus taking $\xi_i = \xi$ and $\phi_i = 0$ we make the Ansatz: $|\Phi(\xi)\rangle \equiv e^{-i\frac{\xi}{2}\sum_{j=1}^N \sigma_j^y} |\uparrow\uparrow \dots \uparrow\rangle$ for searching the maximal the overlap $\Lambda_{\max}(\Psi)$ [16]. This form shows that this separable state can be constructed as a global rotation of the ground state at $h = \infty$, viz., the separable state $|\uparrow\uparrow \dots \uparrow\rangle$. The energy eigenstates are readily expressed in terms of the Jordan-Wigner fermion operators, and so too are the Ansatz states $|\Phi(\xi)\rangle$. By working in this fermion basis we are able to evaluate the overlaps between the two lowest states and the Ansatz states. With $|\Psi_0\rangle$ ($|\Psi_1\rangle$) denoting the lowest state in the even (odd) fermion-number sector, we arrive at the overlaps

$$\langle \Psi_a(r, h) | \Phi(\xi) \rangle = f_N^{(a)}(\xi) \prod_{m=a}^{m < \frac{N-1}{2}} \left[\cos \theta_m^{(a)}(r, h) \cos^2(\xi/2) + \sin \theta_m^{(a)}(r, h) \sin^2(\xi/2) \cot(k_{m,N}^{(a)}/2) \right], \quad (6)$$

$$k_{m,N}^{(a)} \equiv \frac{2\pi}{N} \left(m + \frac{a}{2} \right),$$

$$\tan 2\theta_m^{(a)}(r, h) \equiv r \sin k_{m,N}^{(a)} / (h - \cos k_{m,N}^{(a)});$$

$$f_N^{(0)}(\xi) \equiv 1, \quad f_N^{(1)}(\xi) \equiv \sqrt{N} \sin(\xi/2) \cos(\xi/2), \quad (N \text{ even});$$

$$f_N^{(0)}(\xi) \equiv \cos(\xi/2), \quad f_N^{(1)}(\xi) \equiv \sqrt{N} \sin(\xi/2), \quad (N \text{ odd});$$

where $a = 0, 1$ and $m \in [0, N-1]$ is an integer. The above results are exact for arbitrary N , obtained with periodic boundary conditions on spins rather than the so-called c -cyclic approximation [15]. Given these overlaps, we can readily obtain the entanglement of the ground state, the first excited state, and any linear superposition, $\cos \alpha |\Psi_0\rangle + \sin \alpha |\Psi_1\rangle$ of the two lowest states, for arbitrary (r, h) and N , by maximizing the magnitude of the overlap with respect to the single real parameter ξ .

The formulas that we have just established contain all the results that we explore in the present letter. By analyzing the structure of Eq. (6), we find that the global entanglement does provide information on the phase structure and critical properties of the quantum spin chains. Two of our key results, as captured in Figs. 1 and 2, are

that: (i) although the entanglement itself is, generically, not maximized at the quantum critical line in the (r, h) plane, *the field-derivative of the entanglement diverges as the critical line $h = 1$ is approached*; and (ii) the entanglement *vanishes* at the disorder line $r^2 + h^2 = 1$, which separates the oscillatory and ferromagnetic phases.

Entanglement and quantum criticality—From Eq. (6) it follows that the thermodynamic limit of the entanglement density is given by

$$\mathcal{E}(r, h) = -\frac{2}{\ln 2} \max_{\xi} \int_0^{\frac{1}{2}} d\mu \ln [\cos \theta(\mu, r, h) \cos^2(\xi/2) + \sin \theta(\mu, r, h) \sin^2(\xi/2) \cot \pi\mu], \quad (8)$$

where $\tan 2\theta(\mu, r, h) \equiv r \sin 2\pi\mu / (h - \cos 2\pi\mu)$.

Figure 2 shows the thermodynamic limit of the entanglement density $\mathcal{E}(r, h)$ and its h -derivative in the ground state, as a function of h for three values of r , i.e., three slices through the surface shown in Fig. 1. As the $r = 1$ slice shows, in the Ising limit the entanglement density is small for both small and large h . It increases with h from zero, monotonically, albeit very slowly for small h , then




FIG. 1: (Color online) Entanglement density (upper) and phase diagram (lower) vs. (r, h) for the XY model with $N = 10^4$ spins, which is essentially in the thermodynamic limit. There are three phases: **O**: ordered oscillatory, for $r^2 + h^2 < 1$ and $r \neq 0$; **F**: ordered ferromagnetic, between $r^2 + h^2 > 1$ and $h < 1$; **P**: paramagnetic, for $h > 1$. As is apparent, there is a sharp rise in the entanglement across the line $h = 1$, which signifies a quantum phase transition. The arc $h^2 + r^2 = 1$, along which the entanglement density is zero (see also Fig. 2), separates phases **O** and **F**. Along $r = 0$ lies the XX model, which belongs to a different universality class from the anisotropic XY model.

swiftly rising to a maximum at $h \approx 1.13$ before decreasing monotonically upon further increase of h , asymptotically to zero. The entanglement maximum *does not* occur at the quantum critical point. However, the derivative of the entanglement with respect to h *does* diverge at the critical point $h = 1$, as shown in the inset. The slice at $r = 1/2$ (shifted, for clarity, half a unit to the right) shows qualitatively similar behavior, except that it is finite (although small) at $h = 0$, and starts out by decreasing to a shallow minimum of zero at $h = \sqrt{1 - r^2}$. By contrast, the slice at $r = 0$ (XX) starts out at $h = 0$ at a maximum value of $1 - 2\gamma_C/(\pi \ln 2) \approx 0.159$. (where γ_C is the *Catalan* constant), the globally maximal value of the entanglement over the entire (r, h) plane. For larger h it falls monotonically until it vanishes at $h = 1$, remaining zero for larger h .

We find that along the line $r^2 + h^2 = 1$ the entan-




FIG. 2: (Color online) Entanglement density and its h -derivative (inset) for the ground state of three systems at $N = \infty$. Solid blue line: Ising ($r = 1$) limit; dashed black line: anisotropic ($r = 1/2$) XY model; dash-dotted red line: ($r = 0$) XX model. For the sake of clarity, the XY-case curves are shifted to the right by 0.5, indicated by the green arrow. For the $r = 1/2$, at $h^2 + r^2 = 1$ the entanglement density vanishes, which is a general property for the anisotropic XY model. Note that whilst the the entanglement itself has a non-singular maximum at $h \approx 1.1$ (Ising), $h \approx 1.04$ (XY $r = 1/2$), $h = 0$ (XX), respectively, it has a singularity at the quantum critical point at $h = 1$, as revealed by the divergence of its derivative.

glement density vanishes in the thermodynamic limit. In fact, this line exactly corresponds to the boundary separating the oscillatory and ferromagnetic phases; the boundary can be characterized by a set of ground states with total entanglement of order unity, and thus of zero entanglement density. The entanglement density is also able to track the phase boundary ($h = 1$) between the ordered and disordered phases. Associated with the quantum fluctuations accompanying the transition, the entanglement density shows a drastic variation across the boundary and the field-derivative diverges all along $h = 1$. The two boundaries separating the three phases coalesce at $(r, h) = (0, 1)$, i.e., the XX critical point. Figures 1 and 2 reveal all these features.

The singular behavior of the entanglement density (8) can be analyzed in the vicinity of the quantum critical line, and we find the asymptotic behavior (for $r \neq 0$)

$$\frac{\partial \mathcal{E}}{\partial h} \approx -\frac{1}{2\pi r \ln 2} \ln |h - 1|, \text{ for } |h - 1| \ll 1. \quad (9)$$

From the arbitrary- N results (6) we analyze the approach to the thermodynamic limit, in order to further understand connections with quantum criticality. We focus on the exponent ν , which governs the divergence at criticality of the correlation length: $L_c \sim |h - 1|^{-\nu}$. To do this, we compare the divergence of the slope $\partial \mathcal{E}_N / \partial h$ (i) near $h = 1$ (at $N = \infty$), given above, and (ii) for large N at the value of h for which the slope is maximal (viz. $h_{\max, N}$), i.e., $\partial \mathcal{E}_N / \partial h|_{h_{\max, N}} \approx 0.230r^{-1} \ln N + \text{const.}$, obtained by analyzing Eq. (6) for various values of r . Then, noting that $(2\pi \ln 2)^{-1} \approx 0.2296$ and that the logarithmic

scaling hypothesis [17] identifies ν with the ratio of the amplitudes of these divergences, $0.2296/0.230 \approx 1$, we recover the known result that $\nu = 1$.

Compared with $r \neq 0$ case, the nature of the divergence of $\partial\mathcal{E}/\partial h$ at $r = 0$ belongs to a different universality class:

$$\frac{\partial}{\partial h}\mathcal{E}(0, h) \approx -\frac{\log_2(\pi/2)}{\sqrt{2}\pi} \frac{1}{\sqrt{1-h}}, \quad (h \rightarrow 1^-). \quad (10)$$

From this divergence, the scaling hypothesis, and the assumption that the entanglement density is intensive, we can infer the known result that the critical exponent $\nu = 1/2$ for the XX model. In keeping with the critical features of the XY-model phase diagram, for any small but nonzero value of the anisotropy, the critical divergence of the entanglement derivative is governed by Ising-type behavior. It is only at the $r = 0$ point that the critical behavior of the entanglement is governed by the XX universality class. For small r , XX behavior ultimately crosses over to Ising behavior.

As is to be expected, at finite N the two lowest states $|\Psi_0\rangle$ and $|\Psi_1\rangle$ featuring in Eq. (6) do not spontaneously break the Z_2 symmetry. However, in the thermodynamic limit they are degenerate for $h \leq 1$, and linear combinations are also ground states. The question then arises as to whether linear combinations that explicitly break Z_2 symmetry, i.e., the physically relevant states with finite spontaneous magnetization, show the same entanglement properties. In fact, we see from Eq. (6) that, in the thermodynamic limit, overlaps for both $|\Psi_0\rangle$ and $|\Psi_1\rangle$ are identical, up to the prefactors $f_N^{(0)}$ and $f_N^{(1)}$. These prefactors do not contribute to the entanglement density, and the entanglement density is therefore the same for both $|\Psi_0\rangle$ and $|\Psi_1\rangle$. It follows that, in the thermodynamic limit, the results for the entanglement density are insensitive to the replacement of a symmetric ground state by a broken-symmetry one.

Concluding remarks—In summary, we have quantified the global entanglement of the quantum XY spin chain. This model exhibits a rich phase structure, the qualitative features of which are reflected by this entanglement measure. Perhaps the most interesting aspect is the divergence of the field-derivative of the entanglement as the critical line ($h = 1$) is crossed. Furthermore, in the thermodynamic limit, the entanglement density vanishes on the disorder line ($r^2 + h^2 = 1$). The structure of the entanglement surface, as a function of the parameters of the model (the magnetic field h and the coupling anisotropy r), is surprisingly rich.

We close by pointing towards a deeper connection between the global measure of entanglement and the correlations among quantum fluctuations. The maximal overlap (1) can be decomposed in terms of correlation functions:

$$\Lambda_{\max}^2 = \frac{1}{2^N} + \max_{|\vec{r}|=1} \frac{N}{2^N} \left\{ \langle \vec{r} \cdot \vec{\sigma}_1 \rangle + \sum_{j=2}^N \langle \vec{r} \cdot \vec{\sigma}_1 \otimes \vec{r} \cdot \vec{\sigma}_j \rangle + \dots \right\},$$

where translational invariance is assumed and the Cartesian coordinates of \vec{r} can be taken to be $(\sin \xi, 0, \cos \xi)$. The two-point correlations appearing in the decomposition are related to a bipartite measure of entanglement, namely, the concurrence, which shows similar singular behavior [6]. It would be interesting to establish the connection between the global entanglement and correlations more precisely, e.g., by identifying which correlators are responsible for the singular behavior in the entanglement and how they relate to the better known critical properties.

Acknowledgments—We thank A. Abanov, J.-S. Caux, D. Chang, E. Kim, P. Kwiat, G. Ortiz, M. Randeria, M. Stone, N. Trivedi, W. Zhang, and especially E. Fradkin and A.A. Ludwig for valuable discussions. This work was supported by NSF EIA01-21568 and DOE DEFG02-96ER45434.

-
- [1] C. H. Bennett et al., Phys. Rev. Lett. **70**, 1895 (1993).
 - [2] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. **69**, 2881 (1992).
 - [3] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
 - [4] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge Univ. Press, 2000).
 - [5] T. J. Osborne and M. A. Nielsen, Phys. Rev. A **66**, 032110 (2002).
 - [6] A. Osterloh et al., Nature **416**, 608 (2002).
 - [7] G. Vidal et al., Phys. Rev. Lett. **90**, 227902 (2003).
 - [8] H. Barnum et al., Phys. Rev. Lett. **92**, 107902 (2004).
 - [9] S. Sachdev, *Quantum phase transitions* (Cambridge University Press, 1999).
 - [10] For a review, see M. Horodecki, Quantum Inf. Comput. **1**, 3 (2001), and references therein.
 - [11] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
 - [12] T.-C. Wei and P. M. Goldbart, Phys. Rev. A **68**, 042307 (2003).
 - [13] E_{\log_2} provides a lower bound to another multipartite measure: the relative entropy of entanglement, proposed by V. Vedral et al., Phys. Rev. Lett. **78**, 2275 (1997). For further discussion, see T.-C. Wei et al., quant-ph/0405002.
 - [14] M. Henkel, *Conformal Invariance and Critical Phenomena* (Springer, 1999).
 - [15] E. Lieb, T. Schultz, and D. Mattis, Ann. Phys. **60**, 407 (1961).
 - [16] We have confirmed this Ansatz numerically for small numbers of spins.
 - [17] M. N. Barber, in *Phase Transitions and Critical Phenomena*, Vol. 8, p. 161, Eq. (3.20), C. Domb and J. L. Lebowitz (eds.) (Academic, London, 1983).