

Entanglement of electron spins in superconductors

Sangchul Oh* and Jaewan Kim†

School of Computational Sciences, Korea Institute for Advanced Study, Seoul 130-012, Korea

(Dated: November 5, 2021)

We investigate entanglement of two electron spins forming Cooper pairs in an s -wave superconductor. The two-electron space-spin density matrix is obtained from the BCS ground state using a two-particle Green's function. It is demonstrated that a two spin state is not given by a spin singlet state but by a Werner state. It is found that the entanglement length, within which two spins are entangled, is not the order of the coherence length but the order of the Fermi wave length.

PACS numbers: 03.67.Mn, 03.67.-a, 03.65.Ud, 74.50.+r

Entanglement, referring to the nonlocal quantum correlation between subsystems, is one of the key resources in quantum teleportation, quantum communication, and quantum computation [1]. A study of entanglement in many-body systems is of importance for not only its application to quantum information processing but also giving us new insights on its relevant physics. For example, entanglement in many spin systems has been investigated in connection with quantum phase transition [2]. For a non-interacting electron gas, the entanglement length within which two electron spins are entangled is the order of the Fermi wave length $\lambda_F = 2\pi/k_F$ [3, 4].

A solid state entangler, analogous to the parametric down conversion producing a pair of entangled photons in quantum optics, is a device to generate a pair of entangled electrons in controlled way. It is of interest for the realization of scalable solid-state quantum computers. There have been various proposals to create entangled pairs in solid state systems; spin-entanglement via a quantum dot [5, 6] or via a magnetic impurity [7], generation of entangled electron spins by extracting a Cooper pair out of a superconductor [8, 9, 10, 11, 12, 13, 14], entangled electron-hole pairs in a degenerate electron gas [15, 16], two particle orbital entanglement [17, 18], and entangled spins in electron gases due to exchange interaction [3, 4, 19]. Usually entangling process takes two steps, generation of an entangled electron pair via some kind of interaction and separation of them from each other.

A Cooper pair of a BCS superconductor is composed of two electrons with opposite momenta and a spin singlet state, $\mathbf{k} \uparrow$ and $-\mathbf{k} \downarrow$ [20]. Since the size of a Cooper pair is the order of the coherence length ξ ($\sim 10^{-4}$ cm), entanglement of electron spins may survive within that scale. This means the entanglement length may be about the coherence length. If one can extract a Cooper pair and separate two electrons from each other, the superconductor may be a good natural resource of entangled spin states. In Refs. [8, 9, 10, 11, 12, 13, 14] it is implicitly assumed that the distance between two tunnel

junctions attached to the superconductor should be less than the coherence length ξ . Also it is unclear whether a spin state of a tunneled electron pair is a spin singlet state (one of four Bell states) or a mixed state.

In this Letter we address this problem and find the two-spin state of the BCS ground state is not given not by a Bell state but by a Werner state. We investigate entanglement of the two-spin state as a function of the relative distance between two electrons. Surprisingly, we find the entanglement length is not the order of the coherence length ξ but the order of the Fermi wave length λ_F .

Let us start with the pairing Hamiltonian of the BCS theory [20, 21]

$$H = \sum_{\mathbf{k}s} \epsilon_{\mathbf{k}} c_{\mathbf{k}s}^\dagger c_{\mathbf{k}s} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow}, \quad (1)$$

where $c_{\mathbf{k}s}^\dagger$ is a creation operator for electrons of wave vector \mathbf{k} and z -component of spin s . The normalized BCS ground state of Eq. (1) is given by

$$|\psi_0\rangle = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger) |0\rangle, \quad (2)$$

where coefficients $u_{\mathbf{k}}$ and $v_{\mathbf{k}}$ are written by

$$u_{\mathbf{k}}^2 = \frac{1}{2} \left(1 + \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} \right), \quad (3a)$$

$$v_{\mathbf{k}}^2 = \frac{1}{2} \left(1 - \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} \right). \quad (3b)$$

Here $E_{\mathbf{k}} = \sqrt{(\epsilon_{\mathbf{k}} - \mu)^2 + \Delta_{\mathbf{k}}^2}$ is the excitation energy of a quasi-particle of wave vector \mathbf{k} and $\Delta_{\mathbf{k}}$ the superconducting gap.

In order to investigate the entanglement of two-electron spins forming Cooper pairs, we introduce the two-electron space-spin density matrix in the second quantization

$$\rho^{(2)}(x_1, x_2; x'_1, x'_2) = \frac{1}{2} \langle \hat{\psi}^\dagger(x'_2) \hat{\psi}^\dagger(x'_1) \hat{\psi}(x_1) \hat{\psi}(x_2) \rangle, \quad (4)$$

where $\langle \cdots \rangle = \langle \psi_0 | \cdots | \psi_0 \rangle$ at zero temperature and $x = (\mathbf{r}, s)$. The two-particle Green's function is defined by

$$G(1, 2; 1', 2') = -\langle T[\hat{\psi}_H(1) \hat{\psi}_H(2) \hat{\psi}_H^\dagger(2') \hat{\psi}_H^\dagger(1')] \rangle, \quad (5)$$

*Electronic address: scoh@kias.re.kr

†Electronic address: jaewan@kias.re.kr

where 1 refers to $x_1 t_1$, $\hat{\psi}_H^\dagger$ is the creation operator for electrons in the Heisenberg representation, and T the time-ordering operator. The relation between the two-electron space-spin density matrix and the two-particle Green's function reads

$$\rho^{(2)}(x_1, x_2; x'_1, x'_2) = -\frac{1}{2} G(x_1 t_1, x_2 t_2; x'_1 t_1^+, x'_2 t_2^+), \quad (6)$$

where t^+ denotes time infinitesimally later than t .

The two-particle Green's function for the superconducting state can be factored into single-particle Green's functions [22, 23]

$$G(1, 2; 1', 2') = G(1, 1')G(2, 2') - G(1, 2')G(2, 1') - F(1, 2)F^\dagger(1', 2'). \quad (7)$$

The single-particle Green's function is given by

$$G(1, 1') \equiv -i\langle T[\psi_H(x_1 t_1)\psi_H^\dagger(x'_1 t'_1)] \rangle \quad (8a)$$

$$= \delta_{s_1 s'_1} G(\mathbf{r}_1 t_1, \mathbf{r}'_1 t'_1), \quad (8b)$$

where its spin dependence for a non-magnetic system becomes a unit matrix $\delta_{s_1 s'_1}$. The anomalous Green's function is written by

$$F^\dagger(1, 2) \equiv -i\langle T[\psi_H^\dagger(x_1 t_1)\psi_H(x_2 t_2)] \rangle \quad (9a)$$

$$= I_{s_1 s_2} F^\dagger(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2), \quad (9b)$$

where its spin dependence is given by an antisymmetric matrix

$$I_{ss'} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = i\sigma_y. \quad (9c)$$

Also notice that $F(1, 2) = I_{s_1 s_2} F(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2)$.

Since the system is translational-invariant and the Hamiltonian of the system is time-independent, the Green's functions G and F depend only on the relative coordinates $\mathbf{r}_1 - \mathbf{r}'_1$ and the time difference $t_1 - t'_1$. In the limit that t'_1 goes to t_1 , the spatial part of G becomes

$$iG(\mathbf{r}) = \langle \psi_0 | \hat{\psi}_\sigma^\dagger(\mathbf{r}_1) \hat{\psi}_\sigma(\mathbf{r}'_1) | \psi_0 \rangle \quad (10a)$$

$$= \frac{1}{V} \sum_{\mathbf{k}} v_{\mathbf{k}}^2 e^{i\mathbf{k} \cdot \mathbf{r}}, \quad (10b)$$

where

$$\hat{\psi}_s^\dagger(\mathbf{r}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} c_{\mathbf{k}s}^\dagger e^{i\mathbf{k} \cdot \mathbf{r}}.$$

is the field operator in Schrödinger representation. In the continuum limit one obtains

$$iG(r) = \frac{1}{2\pi^2 r} \frac{m}{\hbar^2} \int_0^\infty v_k^2 \sin(kr) k dk. \quad (11)$$

Due to the similarity between $v_{\mathbf{k}}^2$ of the BCS ground state and the Fermi function of an ideal electron gas at the critical temperature T_c , as shown in Fig. 1, Eq. (11) has

a similar form of Eq. (17) in Ref. [4]. The electron density $n \equiv N/V$ can be calculated by $iG(0) = n/2$. Similarly, the spatial part of F^\dagger is given by

$$iF^\dagger(\mathbf{r}_1 - \mathbf{r}_2) = \langle \psi_0 | \hat{\psi}_\uparrow^\dagger(\mathbf{r}_1) \hat{\psi}_\downarrow^\dagger(\mathbf{r}_2) | \psi_0 \rangle \quad (12a)$$

$$= \frac{1}{V} \sum_{\mathbf{k}} v_{\mathbf{k}} u_{\mathbf{k}} e^{i\mathbf{k} \cdot (\mathbf{r}_1 - \mathbf{r}_2)}. \quad (12b)$$

Note that $F^\dagger(\mathbf{r}_1 - \mathbf{r}_2) = [F(\mathbf{r}_1 - \mathbf{r}_2)]^*$. In the continuum limit one has

$$iF(r) = \frac{1}{\pi^2 r} \int_{\mathcal{R}} \frac{\sin(kr)k}{\sqrt{(\xi/\Delta)^2 + 1}} dk \quad (13)$$

where $\xi_{\mathbf{k}} \equiv \epsilon_{\mathbf{k}} - \mu$, $\Delta_{\mathbf{k}} = \Delta \theta(\hbar\omega_D - |\xi_{\mathbf{k}}|)$ with a step function $\theta(x)$, and the integration over k should be done on the range such that $-\hbar\omega_D \leq \xi_{\mathbf{k}} \leq \hbar\omega_D$ [20, 21, 22]. Here ω_D is the Debye frequency. Thus Eq. (13) becomes approximately

$$iF(r) \approx N(0) \Delta \frac{\sin(k_F r)}{k_F r} K_0\left(\frac{r}{\pi \xi_0}\right) \quad (14)$$

where $N(0)$ is the density of states for one spin projection at the Fermi surface, given by [22]

$$N(0) = \frac{1}{2\pi^2} \left[k^2 \frac{dk}{d\epsilon_k} \right]_{\epsilon_k = \epsilon_F} = \frac{mk_F}{2\pi^2 \hbar^2}, \quad (15)$$

and $K_0(y)$ is a Bessel function of order 0

$$K_0(y) = \int_0^\infty dt \frac{\cos(yt)}{\sqrt{1+t^2}} \approx \sqrt{\frac{\pi}{2y}} e^{-y}. \quad (16)$$

From Eqs. (7), (8) and (9), we have the two-electron space-spin density matrix

$$\begin{aligned} \rho_{s_1, s_2; s'_1, s'_2}^{(2)}(\mathbf{r}_1, \mathbf{r}_2; \mathbf{r}'_1, \mathbf{r}'_2) \\ = -\frac{1}{2} \left[\delta_{s_1 s'_1} \delta_{s_2 s'_2} G(\mathbf{r}_1 - \mathbf{r}'_1) G(\mathbf{r}_2 - \mathbf{r}'_2) \right. \\ \left. - \delta_{s_1 s'_2} \delta_{s_2 s'_1} G(\mathbf{r}_1 - \mathbf{r}'_2) G(\mathbf{r}_2 - \mathbf{r}'_1) \right. \\ \left. - I_{s_1 s_2} I_{s'_1 s'_2} F(\mathbf{r}_1 - \mathbf{r}_2) F^*(\mathbf{r}'_1 - \mathbf{r}'_2) \right]. \end{aligned} \quad (17)$$

Eq. (17) has the same form of the two-electron space-spin density matrix for a non-interacting electron gas, Eq. (8) in Ref. [4], except the last anomalous term. In the limit that $|\mathbf{r}_i - \mathbf{r}'_i| \rightarrow \infty$ for $i = 1, 2$, one has

$$\begin{aligned} \rho_{s_1, s_2; s'_1, s'_2}^{(2)}(\mathbf{r}_1, \mathbf{r}_2; \mathbf{r}'_1, \mathbf{r}'_2) \\ \rightarrow \frac{1}{2} I_{s_1 s_2} I_{s'_1 s'_2} F(\mathbf{r}_1 - \mathbf{r}_2) F^*(\mathbf{r}'_1 - \mathbf{r}'_2) \end{aligned} \quad (18)$$

which shows the off-diagonal long range order of a superconductor [24].

Let us consider the case $\mathbf{r}_1 = \mathbf{r}'_1$, $\mathbf{r}_2 = \mathbf{r}'_2$, which is equivalent to take only the diagonal elements of the space density matrix. For a solid state entangler that produce entangled spins out of a superconductor, two leads are




FIG. 1: (color online) $v_{\mathbf{k}}^2$ (solid line) and $u_{\mathbf{k}}v_{\mathbf{k}}$ (dotted line) as a function of $\epsilon_{\mathbf{k}}/\epsilon_F$. In plotting Δ/ϵ_F is taken to be 1/1000.

attached to two tunneling points \mathbf{r}_1 and \mathbf{r}_2 on the superconductor. This implies one electron is located at \mathbf{r}_1 and the other at \mathbf{r}_2 . The two spin state depending on the relative distance $\mathbf{r} \equiv \mathbf{r}_1 - \mathbf{r}_2$ is given by

$$\rho_{s_1, s_2; s'_1, s'_2}^{(2)}(r) = -\frac{1}{2} \left[\delta_{s_1, s'_1} \delta_{s_2, s'_2} G^2(0) - \delta_{s_1, s'_2} \delta_{s_2, s'_1} G^2(r) - I_{s_1, s_2} I_{s'_1, s'_2} |F(r)|^2 \right]. \quad (19)$$

Let us define functions $g(r) \equiv G(r)/G(0) = 2iG(r)/n$ and $f(r) \equiv F(r)/G(0)$. Eq. (19) becomes

$$\rho_{s_1, s_2; s'_1, s'_2}^{(2)}(r) = \frac{n^2 \mathcal{N}}{8} \rho_{12}, \quad (20)$$

where the two-spin state ρ_{12} of the BCS ground state with normalization factor $\mathcal{N} \equiv 4 - 2g^2 + 2f^2$ is given by

$$\rho_{12} = \frac{1}{\mathcal{N}} \begin{bmatrix} 1 - g^2 & 0 & 0 & 0 \\ 0 & 1 + f^2 & -g^2 - f^2 & 0 \\ 0 & -g^2 - f^2 & 1 + f^2 & 0 \\ 0 & 0 & 0 & 1 - g^2 \end{bmatrix}. \quad (21)$$

We find that the two-spin state ρ_{12} is not a spin singlet state but a Werner state characterized by a single parameter p ($0 \leq p \leq 1$)

$$\rho_{12} = (1 - p) \frac{\mathbf{I}}{4} + p |\Psi^{(-)}\rangle \langle \Psi^{(-)}|, \quad (22)$$

where \mathbf{I} is a 4×4 unit matrix and $|\Psi^{(-)}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$. The parameter p is given by a function of f and g

$$p = \frac{f^2 + g^2}{2 + f^2 - g^2}. \quad (23)$$

For a non-interacting electron gas, the two spin-state is also given by a Werner state and can be obtained from Eq. (21) by putting $f = 0$ [4].

The properties of a Werner state are well known [4, 25]. According to the Peres-Horodecki separability criterion [26, 27], a Werner state is entangled for $p > 1/3$.




FIG. 2: (color online) \tilde{F} (solid line), g (dotted line), C (thick solid line) as a function of $k_F r$. The plotting parameters are given in the text.

Thus ρ_{12} is entangled for $f^2 + 2g^2 > 1$. If $g^2 = 0$, ρ_{12} is entangled for $f^2 \geq 1$. As shown later, f^2 is always much less than 1. The concurrence for ρ_{12} , one of entanglement measures, is calculated as [4, 29]

$$C = \max \left\{ 0, \frac{3p - 1}{2} \right\}. \quad (24)$$

Fig. 2 depicts the concurrence as a function of $k_F r$. This implies that the two spin state ρ_{12} of the BCS ground state is not entangled if the relative distance r between two tunneling points on the superconductor is larger than the Fermi wave length λ_F and even though less than the coherence length ξ .

Let us calculate the magnitude of $f(r)$ which is rewritten as

$$f(r) = \frac{F(r)}{F(0)} \frac{F(0)}{G(0)}, \quad (25)$$

where $\tilde{F}(r) \equiv F(r)/F(0)$ is always less than 1 except at the origin as shown in Fig. 2. On the other hand we obtain

$$\frac{F(0)}{G(0)} \approx \frac{2}{n} N(0) \Delta \ln \frac{2\hbar\omega_D}{\Delta} \quad (26a)$$

$$= \frac{3}{2} \frac{\Delta}{\epsilon_F} \ln \frac{2\hbar\omega_D}{\Delta}. \quad (26b)$$

In the weak coupling limit [20, 21, 22], $\Delta \ll \hbar\omega_D \ll \epsilon_F$, the value of Eq. (26) is very small. For example, for the values $\Delta \approx 1$ meV, $\hbar\omega_D \approx 100$ meV, and $\epsilon_F \approx 1$ eV, we have $F(0)/G(0) \sim 10^{-2}$. This means f^2 is very small compared with g^2 . Thus we demonstrate that two spins are not entangled for $\lambda_F < r < \xi$.

Our results could be explained as follows. A superconductor has many Cooper pairs. Although the spin state of each Cooper pair is a Bell state, the spin correlation between two points \mathbf{r}_1 and \mathbf{r}_2 is due to many Cooper pairs not a single Cooper pair. Thus for solid state entangler which generate entangled spins from a superconductor,

one needs to extract a single Cooper pair or to confirm that two electrons are tunneled out from a single Cooper pair.

In conclusion, we obtained the two-electron reduced density matrix of the BCS ground state based on the Green's function method. We investigated entanglement of two electron-spins forming the Cooper pair in BCS superconductors. It has been found that the two-spin density matrix for a given relative distance between two electrons is given by a Werner state not by a Bell

state. Also the entanglement length is not the order of the coherence length ξ but the order of the Fermi wave length λ_F .

J.K. was supported by Korean Research Foundation Grant KRF-2002-070-C00029. S.O. was partially supported by R&D Program for Fusion Strategy of Advanced Technologies of Ministry of Science and Technology of Korean Government.

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2001)
 - [2] T. J. Osborne and M. A. Nielsen, Phys. Rev. A **66**, 032110 (2002); A. Osterloh, L. Amico, G. Falci, and R. Fazio, Nature (London) **416**, 608 (2002); G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, Phys. Rev. Lett. **90**, 227902 (2003); F. Verstraete, M. Popp, and J. I. Cirac, Phys. Rev. Lett. **92**, 027901 (2004).
 - [3] V. Vedral, e-print quant-ph/0302040.
 - [4] S. Oh and J. Kim, Phys. Rev. A **69**, 054305 (2004).
 - [5] D. S. Saraga and D. Loss, Phys. Rev. Lett. **90**, 166803 (2003).
 - [6] W. D. Oliver, F. Yamaguchi, and Y. Yamamoto, Phys. Rev. Lett. **88**, 037901 (2002).
 - [7] A. T. Costa, Jr. and S. Bose, Phys. Rev. Lett. **87**, 277901 (2001).
 - [8] P. Recher, E. V. Sukhorukov, and D. Loss, Phys. Rev. B **63**, 165314 (2001).
 - [9] P. Recher and D. Loss, Phys. Rev. B **65**, 165327 (2002).
 - [10] P. Recher and D. Loss, Phys. Rev. Lett. **91**, 267003 (2003).
 - [11] C. Bena, S. Vishveshwara, L. Balents, and M. P. A. Fisher, Phys. Rev. Lett. **89**, 037901 (2002).
 - [12] G. B. Lesovik, T. Martin, and G. Blatter, Eur. Phys. J. B **24**, 287 (2001).
 - [13] N. M. Chtchelkatchev, G. Blatter, G. B. Lesovik, and T. Martin, Phys. Rev. B **66**, 161320 (2002).
 - [14] V. Bouchiat, N. Chtchelkatchev, D. Feinberg, G. B. Lesovik, T. Martin, and J. Torrès, Nanotechnology **14**, 77 (2003).
 - [15] C. W. J. Beenakker, C. Emary, M. Kindermann, and J. L. van Velsen, Phys. Rev. Lett. **91**, 147901 (2003).
 - [16] C. W. J. Beenakker and M. Kindermann, Phys. Rev. Lett. **92**, 056801 (2004).
 - [17] P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, Phys. Rev. Lett. **91**, 157002 (2003).
 - [18] P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, Phys. Rev. Lett. **92**, 026805 (2004).
 - [19] A. V. Lebedev, G. B. Lesovik, and G. Blatter, cond-mat/0311423; A. V. Lebedev, G. Blatter, and G. B. Lesovik, cond-math/0311649.
 - [20] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **108**, 1175 (1957).
 - [21] J. R. Schrieffer, *Theory of Superconductivity* (Addison-Wesley, New York, 1988).
 - [22] A. L. Fetter and J. D. Waleka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York, 1971).
 - [23] A. A. Abrikosov, L. P. Gorkov, and I. E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics* (Dover Publications, New York, 1975).
 - [24] C. N. Yang, Rev. Mod. Phys. **34**, 694 (1962).
 - [25] R. F. Werner, Phys. Rev. A **40**, 4277 (1989).
 - [26] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
 - [27] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 1 (1996).
 - [28] R. Horodecki, P. Horodecki, and M. Horodecki, Phys. Lett. A **200**, 340 (1995).
 - [29] S. Hill and W. K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997); W. K. Wootters, *ibid.* **80**, 2245 (1998).