A hsan N azir, B rendon W . Lovett,<sup>y</sup> and G . A ndrew D . B riggs D epartm ent of M aterials, O xford U niversity, O xford O X 1 3PH, U nited K ingdom (D ated: Septem ber 17, 2021)

We show that two initially non-resonant quantum dots may be brought into resonance by the application of a single detuned laser. This allows for control of the inter-dot interactions and the generation of highly entangled excitonic states on the picosecond timescale. A long with arbitrary single qubit manipulations, this system would be su cient for the demonstration of a prototype excitonic quantum computer.

PACS num bers: 03.67 M n, 78.67 H c

Sem iconductor quantum dots (QD's) are offen described as 'arti cial atom s' due to the discrete energy level structure which results from their three-dimensional connement [1]. Consequently, many of the techniques of quantum optics are now being used in QD studies and have led to the observation of R abi oscillations [2], photon antibunching [3], and recently the optical Stark effect [4] in single QD's. Such experiments have stimulated a great deal of interest in possible applications of QD's in quantum information processing (Q IP) devices [5, 6, 7].

In this paper, we shall analyze the behaviour of two adjacent self-assem bled QD's addressed by an external classical laser eld, with the aim of controlling the electronic interactions between them . We shall demonstrate that it is possible to generate and maintain long-lived entangled excitonic states in such QD's through the inter-dot resonant (Forster) energy transfer [5, 8]. This is achieved with a single laser that dynam ically Stark shifts the exciton ground states in and out of resonance, e ectively switching the inter-dot interaction on and o .

Our model considers only the ground state (no exciton) and rst excited state (single exciton) in each dot, and these two states de ne our qubit as pi and jli respectively. Each QD is assumed to be within the strong-con nem ent regim e where their typical sizes are much smaller than the corresponding bulk exciton radius, which is determined by the electron-hole Coulomb interaction. As a result, the con nem ent energy due to QD size dom inates and m ixing of the single-particle electron and hole states due to their Coulomb interactions may be neglected [9]. Any associated energy shift can be absorbed into the exciton creation energy; this shift is in portant as it ensures that the resonance condition for single-particle tunneling is not the same as that for resonant exciton transfer. A dditionally, we consider only weak inter-dot interaction strengths ( 0:1 m eV) which would be expected for two dots with relatively large spacing ( 10 nm) [10]. Therefore, we may neglect inter-dot tunneling of electrons and holes.

The Ham iltonian for two coupled dots in the presence

of a single laser of frequency  $!_1 m$  ay be written in the computational basis f 00i; 01i; 10i; 10i; 11i as (h = 1):

where  $!_0$  is the ground state energy,  $!_{1(2)}$  the exciton creation energy for dot 1(2), and  $!_T = !_0 + !_1 + !_2$ . The coupling term s  $V_F$  and  $V_{XX}$  are the Forster (transition dipole-dipole) and biexciton [6, 7] (static dipole-dipole) interaction strengths respectively.

We have assumed that each dot may couple to the laser with a di erent strength, governed by the respective Rabifrequency  $_1$  or  $_2$ , with  $_i$ (t) 2d E (r;t), for i = 1;2. Here,  $d_i$  is the inter-band ground state transition dipole moment for dot i, and E (r;t) is the laser amplitude at time t and position r. Natural size and composition uctuations in self-assembled dot samples (for example in InG aAs QD's [11]) lead to a large range of possible transition dipole moments for each dot. The size of the ground-state dipole mism atch (and related exciton energy di erence) between two dots is an important factor in determining our ability to control their interactions.

We shall rst analyze the H am iltonian of Eq. 1 within the rotating wave approximation (RWA). This will allow us to derive approximate conditions governing the behaviour of the system, and elucidate the mechanism for controlling excitonic entanglement. Subsequently, we shall characterize the small corrections to these RWA solutions and perform a full numerical solution of H (t).

Transforming Eq. 1 into a frame rotating with the laser frequency  $!_1$  with respect to both qubits, we obtain (within the RWA):

where  $i = !_i !_i$  is the detuning of the laser from dot i, and an irrelevant constant has been subtracted from each term on the diagonal. In order to demonstrate control over the interaction  $V_F$  we would like to isolate the

E lectronic address: ahsan nazir@ m aterials.ox.ac.uk

<sup>&</sup>lt;sup>y</sup>E lectronic address: brendon.lovett@ m aterials.ox.ac.uk




FIG.1: Laser induced anticrossing in the fjD1; jD0ig subspace, for xed ratio  $_1=\ _2$  = 0:55. Eigenvalues are calculated from Eq.4 with  $_1$  = 292:59 meV,  $_2$  = 290:59 meV, and  $V_F$  = 0:1 meV.

behaviour of the fj01;j00ig subspace in which it acts. W e m ay proceed, utilizing degenerate perturbation theory, provided that the following conditions are satis ed:

$$j_1 _{2} \mathbf{j} \mathbf{j}_F \mathbf{j}_F \mathbf{j}_F \mathbf{j}_i = 2j \quad \min(\mathbf{j}_i \mathbf{j}_F \mathbf{j}_i + \mathbf{V}_{XX} \mathbf{j}): \quad (3)$$

In this case the three subspaces fjli,jl0ig, fjD0ig, and fjlig are decoupled and we can write an elective H am iltonian for the fjli; jl0ig subspace:

where = 1 = 1, =  $1 = (2 + V_{XX})$ , and  $\frac{0}{1} = 1 = 2$ .

As we may control both the detunings  $_{\rm i}$  of the laser from the QD's, and the Rabi frequencies  $_{\rm i}$ , it is possible to controllably modify the dynam ics within this subspace. As is shown in Fig.1, two regimes are of particular interest. When the laser is o, the difference in diagonal elements can be much larger than the elective interaction strength if

$$!_1 !_2 V_F$$
: (5)

In this case, the eigenstates of Eq. 4 approach the com – putational basis states 311 and 3101, shown away from the anticrossing in Fig. 1, that would be expected for non-interacting dots.

In contrast, under the condition:

$$_{1} _{2} = !_{1} !_{2} = '_{2} ''_{1} (+);$$
 (6)

the diagonal terms of Eq. 4 are equal and the two dots Stark shift into resonance under the action of the laser. The corresponding eigenstates lie at the anticrossing and




FIG.2: Populations of the four states j00i; j01i; j11i and j10i calculated from Eq.2 with input j01i, nm = hnm j jm i, V<sub>F</sub> = 0:1 meV and V<sub>XX</sub> = 0. Dashed line: state evolution when the laser is always on. The R abi frequencies and detunings of the laser are given by  $_2$  = 40:96 meV,  $_1$  = 0:55  $_2$ ,  $_2$ =2  $_1$  = 0:07, and  $_1$   $_2$  = 2 meV. Dotted line: state evolution when the same laser is on for a time of = (4V<sub>e</sub>) and then turned o. The small oscillations in population in the = (4V<sub>e</sub>) pulse case after the laser is switched o are due to some residual coupling between the dots which can be suppressed by increasing the energy selectivity  $_1$  2. Solid line: state evolution when the laser is always o. In this case the state is alm ost purely j01i throughout.

are maximally entangled due to the modied o -diagonal interaction

$$V_e V_F + {}^{0}_{1} {}^{0}_{2} () :$$
 (7)

They are given by j \_ i = 2 <sup>1=2</sup> (j)li + jl0i) and j i = 2 <sup>1=2</sup> (jl0i j)li). Hence, if the system is initialized in the state j)li, it will coherently evolve to ijl0i during the laser pulse, passing through the maxim ally entangled state 2 <sup>1=2</sup> (j)li + ijl0i). This happens with a coherent exciton transfer time oft = = (2V<sub>e</sub>).

Therefore, we may selectively couple the two initially non-resonant QD's (satisfying Eq. 5 before the laser is switched on) by the application of a single detuned laser satisfying Eq. 6 which non-adiabatically shifts the eigenstates to the anticrossing point in Fig. 1. As soon as we wish to decouple the dots again, we simply turn the laser o.

Thise ect is dem onstrated in Fig.2 where a num erical simulation of the evolution of an input state jD1i under the RW A Ham iltonian of Eq.2 is shown. W ithout laser coupling, the system remains in its initial state jD1i with a delity, F > 1 4  $(V_F = j_1 _2)^2 = 0.99$ . However, under the application of a laser satisfying Eq. 6, coherent oscillations are observed between the states 011 and 101 with an exciton transfer time of 10.9 ps, for the parameters chosen here. This transfer is very close to being 100% complete with little population leaking from the f011; 101 g subspace, justifying our perturbative treatment.

This behaviour could be observed in an experiment by applying a laser pulse for a series of di erent pulse lengths, and afterwards observing the emitted photons. As can be seen in Fig. 1, the two eigenstates when the laser is o (which are approximately 10i and 101i) are separated by some 2 meV. This is readily resolved in a modern spectrom eter { and so a measurem ent of the wavelength of the emitted photons can be use to determine which of the two dots each one came from . Plotting the number of each wavelength of detected photons as a function of the pulse length would allow a determ ination of the coherent transfer oscillations. Further, a  $t = (4V_e)$  pulse will create and maintain an (approxim ately m axim ally) entangled state from an initially separable one, and this is also shown in Fig 2. Even for the sm all coupling strength ( $V_F = 0:1 \text{ meV}$ ) considered here this operation is on the picosecond time scale. Therefore, we would expect such an entangled state to be long-lived in a pair of QD 's relative to the tim escale of its generation. Single and coupled dot exciton lifetim es have been m easured to be as long as nanoseconds at low tem peratures [12, 13, 14]. Additionally, pure phonon dephasing e ects are suppressed as the tem perature is low ered below 10 K [12, 15].

We therefore now account for the nite exciton lifetimes by including only spontaneous emission terms in the density matrix master equation [16], and neglecting pure dephasing processes:

$$= iH; ] + \frac{1}{2} X_{i} 2_{i} + \frac{1}{2} + \frac$$

Here, the i label the dipole allowed transitions in the coupled system,  $_{i}^{+}$  and  $_{i}$  are their raising and low ering operators, and the i are their transition rates. These term s can lead to a signi cant reduction of the degree of entanglem ent over time. We can see this by referring to Fig. 3, where we show the result of num erical calculations which use the full Ham iltonian (Eq.1) in the master equation (Eq.8). In Fig.3 (a) the populations of the four states are shown as a function of time for the input state j01i, subject to a square pulse of 5:45 ps duration (this satis est =  $= (4V_e)$  for our chosen parameters), and subject to signi cant decay. In Fig. 3 (b), we plot the entanglement of formation (EOF) [17] of the system as a function of time for the input state D1i, for a variety of decay rates. The EOF m easures the number of Bell states required to create the state of interest; for a maximally entangled state it is equal to unity while for a separable state it is zero. For a general two qubit state




FIG.3: (a) Num erical simulation of the four populations for an input state j01i and a =(4V<sub>e</sub>) pulse. The simulation uses Eq.8, which does not rely on the RWA and includes exciton decay, but not pure dephasing. The decay rates for dots 1 and 2 are given by  $_{1} = _{1}^{1} = (331 \text{ ps})^{1}$  and  $_{2} = _{2}^{1} = (100 \text{ ps})^{1}$  respectively. All other parameters are the same as for Fig. 2 except  $_{1} = _{2} = 2:18 \text{ meV}$  to account for the extra shifts, and  $!_{1} = 1500 \text{ meV}$ . (b) Entanglement of form ation of the input state j01i as a function of time for a =(4V<sub>e</sub>) laser pulse, and for a series of di erent decay rates. We keep a constant ratio of  $_{1} = _{2} = jl_{1}=d_{2} f = (_{1} = _{2})^{2}$ . The calculations are made by using Eqs. 8-10.

it is given by the equation

$$E_{\rm F}() = h \frac{1 + p 1}{2}$$
; (9)

where  $h(x) = x \log (x)$  (1 x)  $\log (1 x)$  is the Shannon entropy function. is the \tangle" or \concurrence" squared, which can be computed by using the equation:

$$= C^{2} = [m \operatorname{axf}_{1} _{2} _{3} _{4}; 0g]^{2}: (10)$$

 com putational basis j00i; j11i; j10i; j11i [18]. In Fig. 3 we see that at the end of the laser pulse the state has becom e alm ost maxim ally entangled and, in the absence of decay, stays so once the laser is switched o (the sm all deviation from unity is due to the residual e ect of the Forster coupling when the laser is o ). When decay is included the EOF decreases over time, but for typical experimental lifetimes of 1 ns [12, 13, 14], it retains a value which is higher than 0.94 for times up to 50 ps [19].

The num erical solution of Eq.8 required for Fig.3 has been computed without making use of the RWA. The behaviour of the system is exactly as expected from the RW A case, except for an extra small shift in the dot energies when the laser is on. This shift is the largest correction to the Stark shift in a perturbative expansion and arises from the counter-propagating terms in the state evolution which are discarded when the RWA is made [20]. The corrections to the RWA can all be derived by following the method of Shirley [21], who considers the case of a two-level system (which we label 0 and 1 and consider to have an energy separation of  $!_1$ ) coupled by an oscillating eld (of frequency  $!_1$ ). He shows that this kind of problem can be mapped on to a time independent one by using F loquet's theory to construct a Hamiltonian in an in nite dimensional Hilbert space given by

$$hnjH_F jmi = H^{nm} + n! \qquad nm: \qquad (11)$$

Here ; 2 f0;1g and the n;m represent Fourier com – ponents of the state evolution. H  $^n$  is the Fourier com – ponent with frequency n! of the oscillating H am iltonian (which is only non-zero for  $\frac{6}{5}$ ). Shirley showed that the time evolution operator of the two-level system can be written as

U 
$$(t;t_0) = hnjexp[iH_F(t_0)]j0ie^{in!t}$$
: (12)

He goes on to derive the B loch-Siegert shift, which is in portant for resonant interactions between the two-level system and the oscillating eld. However, we are interested in the o-resonant behaviour. In the absence of any interactions, our system will initially be in a state of the form a 100i + b 10i (in the notation j ni), and we can use the time evolution operator (Eq. 12) to determ ine the energy separation of the two (eigen) states, 101 and jlOi. We follow a similar procedure when the laser is switched on. In this case, if the laser is su ciently detuned from resonance (i.e. if =  $(!_1 !_1)$ =2,with the laser-system coupling), the jni are still approxim ate eigenstates and we can em ploy second order perturbation theory to obtain the energy shift in the separation of the two levels. W e obtain:

$$= \frac{2^{-\alpha}}{(+2!)} + \frac{2^{-\alpha}}{(+2!)}$$
(13)

where, as before,  $^{0} = = 2$ . The extra term represents a detuning from resonance by an amount  $2!_{1}+$ , com pared

to the term with detuning which is usually the only one kept. For the relatively large detunings considered in our two dot model, they become non-negligible contributions to the Stark shifts for each dot, i, with a magnitude of  $2 \frac{\alpha}{i} = (2!_1 + i) \cdot 0$  now we account for this extra shift, for example in this calculation by redening the parameters 1 and 2 to include it, the system behaves exactly as expected from our earlier analysis of Eq. 4.

We now assess the feasibility of our method by examining the state-of-the-art in real system s. In our previous work [8], we predicted that the Forster transfer energy can be as large as about 1 m eV in Q D's (corresponding to energy transfer times on the sub-picosecond timescale); we also showed how a static electric eld can be applied to the dots to suppress the interaction and so lengthen the transfer time as required. In addition to this, experimental work has suggested that Forster transfer times can approach picoseconds in structurally optimized pairs of C dSe Q D's [22], and that these transfer times can be on the subpicosecond timescale in photosynthetic biom olecular system s [23]. M olecular system s could therefore provide an alternative route to experimental realization of the elects we have predicted here.

QD exciton - laser interaction strengths (which correspond to the Rabi frequency when the laser is resonant with the exciton) of a few meV have been attained in an experiment which measured the optical Stark shift in A lA s/G aA s heterostructures [4]. Further, an experiment which directly observed Rabi oscillations in a In-G aA s/G aA s QD photodiode also measured a Rabi frequency of a similar magnitude [24]. However, these experiments were not designed to maxim ize the laser -exciton coupling strength, and so the value of 40 meV, which we used in our simulations, is not unrealistic.

The coherent exciton transfer process detailed in Fig.2 is equivalent to the realization of iSW AP logic operations between the pair of excitonic qubits [25]. A long with arbitrary single qubit rotations, this gate would be su cient for a demonstration of a prototype excitonic quantum computer. Single qubit operations in our two qubit system may be achieved by using their frequency addressability. A laser resonantly tuned to either of the two dots will induce a Rabioscillation in the dot to which it is tuned; in the language of Pauli spin matices this is a rotation around the X axis of the Bloch sphere, and represents one of the two rotations which are required for arbitrary single qubit operations. The other could be obtained in a number of ways. For example a slightly detuned laser will cause the qubit state to move around another trajectory on the Bloch sphere. A Itematively, a higher level Ji (of energy ! T relative to the ji) could be used. If this higher level has a di erent energy for each qubit, it could be resonantly excited from , say, the Di state in only one of the two qubits by applying an appropriately tuned laser pulse. Leaving the system to evolve then for a time before using another pulse to deexcite it causes the  $\eta$  to pick up a phase of  $!_{T}$  relative to the jli. This Z rotation is su cient to complete

the set required for universal quantum computing.

If the ratio of decoherence time to gate operation time in our system were large enough, full scale fault tolerant quantum computing (FTQC) [26] would be possible. It is well established that a ratio of around 1000 is good enough for FTQC, and recent estimates of this have been a low as 100 [27]. We have dem onstrated that an entangling gate can be performed in real systems in around 5 ps, which is around 200 times shorter than the longest m easurem ents of decoherence tim es [12]. Hence, our proposed scheme could be carried out with a precision which is already on the limit required for FTQC; increm ental im provem ents in either FTQC protocols or in experimental systems should enable the implementation of full quantum algorithms. The absolute speed at which gates can be carried out in our scheme also makes them ideal for smaller scale applications such as quantum repeaters [28], which require much less stringent gate delities than full scale FTQC.

To sum m arize, we have shown that two non-resonant QD's m ay be brought into resonance by the application of a single detuned laser which induces Stark shifts within

- [1] L. Jacak, P. Hawrylak, and A. Wojs, Quantum Dots (Springer-Verlag, Heidelberg, 1998).
- [2] T.H.Stievater, X.Li, D.G.Steel, D.Gammon, D.S. Katzer, D.Park, C.Pierm arocchi, and L.J.Sham, Phys. Rev.Lett. 87, 133603 (2001).
- [3] P. M ichler, A. Im am oglu, M. D. M ason, P. J. Carson, G.F. Strouse, and S.K. Buratto, Nature 406, 968 (2000).
- [4] T.Unok, K.Mueller, C.Lienau, T.Elsaesser, and A.D. Wieck, Phys. Rev. Lett. 92, 157401 (2004).
- [5] L.Quiroga and N.F.Johnson, Phys. Rev. Lett. 83, 2270 (1999).
- [6] E. Biolatti, I. D'Amico, P. Zanardi, and F. Rossi, Phys. Rev. B 65, 075306 (2002).
- [7] F. Troiani, U. Hohenester, and E. Molinari, Phys. Rev. B 62, 2263 (R) (2000).
- [8] B.W. Lovett, J.H. Reina, A. Nazir, and G.A.D. Briggs, Phys. Rev. B 68, 205319 (2003).
- [9] S. Schm itt-Rink, D. A. B. M iller, and D. S. Chem la, Phys. Rev. B 35, 8113 (1987).
- [10] A.Nazir, B.W. Lovett, S.D.Barrett, J.H.Reina, and G.A.D.Briggs, xxx lanlgov/quant-ph/0309099.
- [11] P.G.Eliseev, H.Li, A.Stintz, G.T.Lin, T.C.Newell, K.J.M alloy, and L.F.Lester, Appl. Phys.Lett. 77, 262 (2000).
- [L2] M. Bayer and A. Forchel, Phys. Rev. B 65, 41308 (R) (2002).
- [13] D. Birkedal, K. Leosson, and J. M. Hvam, Phys. Rev. Lett. 87, 227401 (2001).
- [14] P. Borri, W. Langbein, U. Woggon, M. Schwab,

each dot w ithout signi cant population excitation. This in turn allows for control over the inter-dot interactions, and hence the generation of highly entangled states on the picosecond timescale. The conditions of Eqs. 3, 5, and 6 set the upper limits on the energy selectivity and, neglecting incoherent processes, the delity possible with a particular dot sample. The Forster strength  $V_F$  sets the interaction timescale. In general, as the magnitude and di erence of the Rabi frequencies increases, and as  $V_F$  increases, so does the feasibility of the proposed idea. We believe that the means to demonstrate the e ects outlined above already exist. Moreover, these types of experimentsm ay prove invaluable in assessing the potential applicability of sem iconductor QD's for future QIP technologies.

AN and BW L are supported by EPSRC (BW L as part of the Foresight LINK Award \Nanoelectronics at the Quantum Edge," GR/R66029, and the QIPIRC, GR/S82176).We thank T.P.Spiller, S.D.Barrett, and W.J.Munro for useful and stimulating discussions, and R.G.Beausoleil for providing numerical simulation code.

M. Bayer, S. Fafard, Z. W asilewski, and P. Hawrylak, Phys. Rev. Lett. 91, 267401 (2003).

- [15] P.Borri, W.Langbein, S.Schneider, U.Woggon, R.L. Sellin, D.Ouyang, and D.Bimberg, Phys. Rev. Lett. 87, 157401 (2001).
- [16] G.S.Agarwal, Quantum Optics (Springer, Berlin, 1974).
- [17] C.H.Bennett, D.P.D iV incenzo, J.A.Smolin, and W.K. W ootters, Phys. Rev. A 54, 3824 (1996).
- [18] W .J.Munro, D.F.V. James, A.G. White, and P.G. Kwiat, Phys. Rev. A 64, 030302 (2001).
- [19] If j00i or j11i are used as the input states,  $E_F = 0$  throughout the subsequent evolution.
- [20] S.M. Barnett and P.M. Radmore, M ethods in Theoretical Quantum Optics (Clarendon Press, Oxford, 1997).
- [21] J.H. Shirley, Phys. Rev. 138, B 979 (1965).
- [22] S.A. Crooker, J.A. Hollingsworth, S. Tretiak, and V.I. K lim ov, Phys. Rev. Lett. 89, 186802 (2002).
- [23] J.L.Herek, W .W ohlleben, R.J.Cogdell, D.Zeidler, and M.Motzkus, Nature 417, 533 (2002).
- [24] A. Zrenner, E. Beham, S. Stu er, F. Findeis, M. Bickler, and G. Abstreiter, Nature 418, 612 (2002).
- [25] N. Schuch and J. Siewert, Phys. Rev. A 67, 032301 (2003).
- [26] A.M. Steane, Phys. Rev. A 68, 042322 (2003).
- [27] B.W. Reichardt, xxx.lanlgov/quant-ph/0406025.
- [28] W . Dur, H.-J. Briegel, J. I. Cirac, and P. Zoller, Phys. Rev. A 59, 169 (1999).