Noise cancellation effect in quantum systems

Paolo Solinas,[1] Nino Zanghì[1]
[1] *Istituto Nazionale di Fisica Nucleare (INFN) and Dipartimento di Fisica, Università di Genova, Via Dodecaneso 33, 16146 Genova, Italy*
(Dated: January 2, 2022)

We consider the time evolution of simple quantum systems under the influence of random fluctuations of the control parameters. We show that when the parameters fluctuate sufficiently fast, there is a cancellation effect of the noise. We propose that such an effect could be experimentally observed by performing a simple experiment with trapped ions. As a byproduct of our analysis, we provide an explanation of the robustness against random perturbations of adiabatic population transfer techniques in atom optics.

PACS numbers: 03.67.Lx

arXiv:quant-ph/0409020v1 3 Sep 2004

I. INTRODUCTION

In the last ten years the interest around the control and manipulation of quantum systems has grown very fast. The possibility to encode and process information has lead to innovative proposals. The major results have been achieved in Quantum Information Processing (QIP), including both theoretical and experimental ones. Quantum cryptography [1] and information transfer protocols [2] have enhanced our understanding of information processing and this in the near future will presumably lead to a significant technical advancement. Quantum Computation (QC) is still in a initial stage: even if the quantum computers seem to be able to solve quickly some problems which are intractable with classical computers [3], more quantum algorithms are required to extend its applicability.

Unfortunately, the quantum system are very delicate and they are subject to two different kinds of errors. On one hand, there is the loss of information due to decoherence the unavoidable interaction of quantum systems with their environments. This problem has been extensively studied over the past few years and proposals to overcome it have been put forward (and a few have been experimentally tested). These proposals include error avoiding [4], error correcting strategies [5] and decoupling techniques [6]. The other source of errors is the imprecise control of the parameters which perform the quantum operation (e.g. the laser or the magnetic field). How to handle such errors is an open problem, though some progress has been made in the framework of the so called geometrical quantum computation [7, 8, 9]. The goal of this paper is to approach the second problem in very simple and idealized situations.

A simple way to model the parameter noise is to consider a quantum system subject to a stochastic fluctuating field with zero mean. Such a model has been recently considered in order to study the effect of the noise on holonomic quantum gates: in Ref. [10] it has been shown that there is a cancellation effect for a fast fluctuating stochastic field and shown that such a cancellation is due to the geometrical dependence of the holonomic operator. Recently, the general validity of such cancellation effect has been clarified: in Ref. [11] is has be shown that for sufficient fast fluctuating stochastic field with zero mean the effects of the noise are wiped out.

In this paper we shall study simple quantum systems subjected to stochastic noise and discuss some applications. In Section II we consider random perturbations which are diagonal in the logical basis and propose an experiment to test the cancellation effect of the the noise—a simple modification of the experiment done by Kielpinski *et al.* [14]. Then we consider a more general noise and compute the *fidelity*. By elementary perturbation theory, we show that the effects of the noise are wiped out. In Section III we discuss how noise cancellation could be relevant for *adiabatic population transfer* experiments in atoms optics (in this case, the cancellation effect avoids the break of the adiabatic approximation, and allows for desired transformation, even in presence of fast fluctuating noise).

For sake of simplicity, all the simulations and analytical calculation are done with Gaussian noise distributions but presumably analogous results can be achieved with a generic stochastic noise with zero mean [11].

II. TWO-LEVEL SYSTEMS

Consider a two-level system evolving according to Hamiltonian

$$H(t) = H_0 + \delta H_I(t). \tag{1}$$

Suppose that $H_0 = B_z\sigma_z$ (with σ_z being a Pauli matrix) and that $\delta H_I(t)$ is of the form

$$\delta H_I(t) = \sum_{j=0}^{N} \delta A^j S_j(t)\,, \tag{2}$$

where δA^j are random 2×2 matrices and $S_j(t)$ are "box functions" with time step τ, i.e., they are functions equal to 1 in the time interval $(j\tau,(j+1)\tau)$ and zero otherwise. Let T be some "final" time at which we wish to consider the system, and let $\tau = T/N$; τ can be regarded the correlation time of the noise. Then the evolution operator

from time zero to time T, generated by $H(t)$ with $\delta H_I(t)$ given by (2), can be written as

$$U(T) = U_N U_{N-1} \cdots U_j \cdots U_2 U_1 \tag{3}$$

where

$$U_j = \exp\left[-i\tau\left(B_z\sigma_z + \delta A^j\right)\right] . \tag{4}$$

A. Diagonal Noise

First, we shall focus on the very simple case corresponding to the choice $\delta A^j = \delta B_z^j \sigma_z$, where the δB_z^j, $j = 1, \ldots, N$ are independent Gaussian distributed random variables with mean zero. In this case the noise is diagonal in the logical basis $\{|\uparrow\rangle, |\downarrow\rangle\}$ in which σ_z is diagonal. Then (3) is trivially computed (due to the commutativity of the δA^j),

$$U(T) = U^0(T)\delta U(T) \tag{5}$$

where $U^0(T) = \exp(-iTH_0)$ is the evolution generated by H_0, and

$$\delta U(T) = \begin{pmatrix} e^{-i\sum_j \frac{\delta B_z^j}{N} T} & 0 \\ 0 & e^{i\sum_j \frac{\delta B_z^j}{N} T} \end{pmatrix} \tag{6}$$

A standard performance estimator is the *fidelity* [15], which, in our case, is given by

$$\mathcal{F}_T = \sqrt{\left|\langle\psi^{(0)}(T)|\psi(T)\rangle\right|^2} \tag{7}$$

where $\psi^{(0)}(T)$ is the "ideal" final state evolved according to H_0 (that is, when the noise is turned off) and $\psi(T)$ is actual final state (that is, when the noise is turned on). The two final states are generated, of course, by the *same* (generic) initial state $|\psi(0)\rangle = \alpha|\uparrow\rangle + \beta|\downarrow\rangle$. From (5) and (6) one easily obtains

$$\mathcal{F}_T^2(\chi, \alpha, \beta) = |\alpha|^4 + |\beta|^4 + 2|\alpha|^2|\beta|^2 \cos\chi , \tag{8}$$

where

$$\chi = 2T\frac{1}{N}\sum_{j=0}^{N} \delta B_z^j . \tag{9}$$

Since the random variables δB_z^j in (9) are taken to be independent and identically distributed according to a Gaussian with zero mean and variance δB^2, the probability distribution of χ is

$$\mathcal{P}(\chi) = \frac{\sqrt{N}}{\sqrt{2\pi}2T\delta B} e^{-\frac{N\chi^2}{2(2T)^2\delta B^2}} , \tag{10}$$

By averaging (8) with respect to (10), we arrive at the mean square fidelity:

$$\begin{aligned} \mathcal{F}_T^2(\alpha, \beta) &= \int d\chi \mathcal{P}(\chi)\mathcal{F}_T^2(\chi, \alpha, \beta) = \\ &= |\alpha|^4 + |\beta|^4 + 2|\alpha|^2|\beta|^2 e^{-\frac{2T^2\delta B^2}{N}} \end{aligned} \tag{11}$$




FIG. 1: The *square fidelity* from noisy process described by (1). The solid line is the theoretical mean square fidelity, eq. (11), for $T = 100$ (in arbitrary units) and $\delta B = 0.1$.

As it can be easily seen, for $N \to \infty$ and $\tau \to 0$ (while keeping $N\tau = T$ of order 1), the mean square fidelity $\mathcal{F}_T^2(\alpha, \beta)$ approaches 1 (since $|\alpha|^2 + |\beta|^2 = 1$), independently of the initial state. This corresponds to a short correlation time of the noise, that is, to a fast random fluctuating field.

Now we'd like to provide some perspective on such limit behavior. Of course, the limit $\tau \to 0$ is only an idealization for τ small but finite. The above results are obtained for *constant* δB^2, but the Heisenberg uncertainty relation imposes strong constraint on the energy fluctuation happening in such a short time interval, $\delta B^2 \propto 1/\tau$. Note that for $\delta B^2 \propto N$ eq. (10) and (11) describe a system interacting with a white noise environment that is the standard way to model the decoherence effect. So, for very small τ the variance δB^2 should not be any more considered constant and our approximation of constant variance breaks down. However, rather than microscopic (environmental) noise, we are interested in modeling the macroscopic (parametric) noise due to imprecision in the control field, and for such our approximation should apply. In this case it is interesting to ask whether the condition of small τ could be physically relevant. In general, in a quantum evolution the final time T is fixed; the correlation time τ can be hardly controlled (i.e., stabilizing the control field) and we are not in the condition to have a cancellation effect. In Section II B we present an experimental proposal to test the presence of this effect; the experiment realizability lies in the control of the correlation time τ of the *simulated* noise. Moreover, there are situations in which this effect can be experimentally relevant: when we have a further degree of freedom and can change the evolution time T. In these cases, fixed τ with the above properties, we can prolong T in order to have $T \gg \tau$ (that is $N \gg 1$) and exploit the cancellation effect. We give an example of such situation in a adiabatic evolution in Section III.

We observe that another interesting feature of eq. (11) is that the cancellation of the noise does not depend on




FIG. 2: The decoherence effect during the evolution for two different value of $N = T/\tau$. Theoretical curves (dashed line) and those obtained by numerical simulations (solid line) are showed. For the theoretical curve (eq. 13) the decoherence time is $t_{deco} = \frac{N}{2T^2\delta B^2}$.

the strength δB: it is always possible to find a suitable N in order to obtain cancellation of the noise—invert equation (11) and express N in terms of δB for given $\mathcal{F}_T$ and evolution time T.

We recall that dependence of the fidelity on a specific choice of the initial state is usually eliminated by averaging $\mathcal{F}_T^2(\alpha,\beta)$ over all the possible initial states with respect to the uniform distribution on the unit sphere in the Hilbert space of the system [15] in our case the (projective) sphere $|\alpha|^2+|\beta|^2=1$. Performing this operation yields to

$$\mathcal{F}_T^2 \equiv < \mathcal{F}_T^2(\alpha,\beta) > = \frac{1}{3}(2 + e^{-\frac{2T^2\delta B^2}{N}}) \qquad (12)$$

The simple model we have been considering here is often used as a toy model for phenomenological decoherence [16, 17]. The relationship between noise cancellation and decoherence is easily seen by considering the time evolution at the discrete times $t = k\tau$, $k \leq N$. By proceeding as before, the average square fidelity at time $t = k\tau$ results

$$\mathcal{F}_t^2 = \frac{1}{3}(2 + e^{-\frac{2Tt\delta B^2}{N}}) \qquad (13)$$

with which it is quite natural to associate a 'decoherence time' $t_{deco} = \frac{N}{2T\delta B^2}$ (see Figure 2 [12, 13]).

B. Experimental proposal

In Ref. [14] Kielpinski *et al.* used the same idea to simulate the decoherence effect due to interaction of the quantum system with the environment degree of freedom. The authors used trapped ions and study the coherence of superposition of quantum state subject to simulated noise. The logical states were the hyperfine states of a trapped Beryllium ion $|F = 2, m_F = -2\rangle$ and $|F = 1, m_F = -1\rangle$ sublevels of the ground state $^2S_{1/2}$. The environment noise was simulated shining the ions with a off-resonant laser with random varying amplitude for the electromagnetic field E^j and random intensity (proportional to $(E^j)^2$). The laser electromagnetic field produce a AC Stark effect on the ions and let one state to acquire a random phase respect to the other. This effect is *quadratic* (*quadratic Stark effect*) in the electromagnetic field E^j. In fact, the two hyperfine states have the same angular momentum (they have both $L = 0$) and the difference is in the spin part of the wave function. The splitting of the energy level is *linear* (*linear Stark effect*) for state with different angular momentum since only these states have non-vanishing matrix element $\langle L = i|\vec{E}\cdot\vec{r}|L = k\rangle$ (with $i \neq k$). Then in the above example we have corrections to the energy level only quadratic and not linear in E^j. Because of the quadratic energy shift, we have a random phase difference proportional to $(E^j)^2$ for every τ interval. Even if this effect is sufficient to produce decoherence effect (as found by the authors), we cannot presumably see the cancellation effect discussed above in a transparent way since our new stochastic variable $\chi \propto \sum (E^j)^2$ has no zero mean.

A small modification of this experiment should allow us to see sharply this cancellation effect. To have an evolution described by Hamiltonian (1) it is sufficient to use states with different angular momentum in order to produce a *linear Stark effect*. At every time interval τ the perturbation of the laser produce a shift of the energy levels proportional to the random intensity of the laser E^j; this produces an evolution where the phase difference between the states is given by a known dynamical part plus a random phase $exp(2i\alpha E^j\tau)$ (where α is a proportional constant). In this case, the new stochastic variable $\chi \propto \sum_j E^j$ has still zero mean and we expect to obtain results shown in the previous section: fixed the evolution time T we should see an increase of $\mathcal{F}^2$ as the correlation time decreases (see Figure 1). Moreover, if the environment decoherence does not depend on the simulated noise, we should be able to see effect analogous to the ones in Figure 2. In particular, subtracted the effect of the environment, the decoherence time should increase as $t_{deco} \propto \frac{1}{\tau}$.

C. Off-Diagonal Noise

We now consider the case of off-diagonal noise, that is, the matrices δA^j in (2) are of the form $\delta A^j = \delta B_x^j\sigma_x + \delta B_y^j\sigma_y$, with δB_x^j, δB_y^j, $j = 1,\ldots,N$ independent Gaussian distributed random variables with mean zero, and σ_x, σ_y being the usual Pauli matrices. Then the one-step evolution operators U_j in (3) are

$$U_j = e^{-i\vec{B}^j\cdot\vec{\sigma}\tau} = \cos(B^j\tau) - i\hat{n}^j\cdot\vec{\sigma}\sin(B^j\tau)\,,$$

where B^j is the modulus of the vector $\vec{B}^j = (\delta B_x^j, \delta B_y^j, B_z)$ and $\hat{n}^j$ is the associated unit vector. By




FIG. 3: Cancellation effect for system subject to off-diagonal noise discussed in section II C. For the stochastic process $\delta B = 0.01$. When the number of random extraction increases the *fidelity* approaches 1.




FIG. 4: Figure shows the population evolution of the three states during the adiabatic transfer process for two different correlation time noise ($\tau_i = T/N_i$). For the first curve $N_1 = 10^4$ and for the second one $N_2 = 10^5$ for $T = 100$ (in arbitrary units).

a perturbation expansion in $\epsilon_k = \delta B_k^j/B_z$, $k = x, y$, we obtain

$$\begin{aligned} U_j &= \cos(B_z\tau) - i\sigma_z \sin(B_z\tau) \\ &- i(\sigma_x \frac{\delta B_x^j}{B_z} + \sigma_y \frac{\delta B_y^j}{B_z}) \sin(B_z\tau) + O\left(\epsilon_k^2\right) \end{aligned}$$

Note that the zero order terms are nothing but $U^0(\tau)$. We shall denote by δU_j the first order terms, i.e.,

$$\delta U_j \equiv i(\sigma_x \frac{\delta B_x^j}{B_z} + \sigma_y \frac{\delta B_y^j}{B_z}) \sin(B_z\tau)$$

Under the assumption that $\epsilon_k = \delta B_k^j/B_z \ll 1$ the operator $U(T)$ can be easily computed by first order perturbation theory: by keeping in (3) only the terms that are first order in ϵ_k we obtain

$$U(T) = U^0(T) - \sum_j \delta P^j \tag{14}$$

where $U^0(T) = \exp(-iTH_0)$ and

$$\delta P^j = \left(U^0(\tau)\right)^{j-1} \delta U_j \left(U^0(\tau)\right)^{N-j}$$

To compute the fidelity (7) for the (generic) initial state $|\psi(0)\rangle = \alpha|\uparrow\rangle + \beta|\downarrow\rangle$ we need to calculate the scalar product $\langle\psi^{(0)}(T)|\psi(T)\rangle = \langle U^0(T)\psi(0)|U(T)\psi(0)\rangle$. According to (14), this is given by

$$1 - \langle\psi(0)|(U^0(T))^\dagger(\sum_j \delta P^j)|\psi(0)\rangle$$

The matrix elements of δP^j in the logical basis are: $(\sum_j \delta P^j)_{mm} = 0$ and

$$(\sum_j \delta P^j)_{lm} = \frac{\sin(B_z\tau)}{B_z} \sum_j e^{iB_z\tau(N-2j+1)}(\delta B_x^j \pm i\delta B_y^j)\,.$$

By taking into account that $\sin(B_z\tau) \approx B_z\tau$ and $\exp(iB_z\tau) \approx 1$ for $\tau = T/N \ll 1$, we may finally evaluate the modulus of the scalar product and compute the fidelity. We obtain

$$\mathcal{F}_T = \left|1 - 2T\mathrm{Re}\left(\sum_j e^{i2B_z(T-\tau j)}(\frac{\delta B_x^j}{N} - i\frac{\delta B_y^j}{N})\alpha^*\beta\right)\right| \tag{15}$$

It is important to note that, also in this case, for $N \to \infty$ and $\tau \to 0$ (while keeping $N\tau = T$ of order 1), the fidelity approaches 1. This is so because δB_k^j are independent random variables with mean zero [18].

So, also in this case there is a noise cancellation effect. This effect is confirmed by the numerical simulations shown in Figure 3.

III. ADIABATIC EVOLUTION

The *adiabatic population transfer* is an important technique used in atoms optics to achieve population transfer between quantum states of atoms and molecules [19]. We first create coherence between the initial and final state (*population trapping*) and then produce an adiabatic evolution to transfer the population to the final state. This scheme has seen a great success and has been used in many different areas: chemical reaction [20], laser-induced cooling [21], atoms optics [22], cavity quantum electrodynamics [23, 24]. The wide range of application is due to many advantages of this scheme: it is easy to implement in different system, it has an high rate of population transfer and it is robust respect to variations of field parameters [25].

Consider two states $|1\rangle$ and $|2\rangle$ coupled to an excited state $|e\rangle$ by two lasers (i.e., a Λ system). The states $|1\rangle$ and $|2\rangle$ can be degenerate or quasi-degenerate but it is important that we can address separately both of




FIG. 5: Figure shows the *fidelity* and the population of exited state $|e\rangle$ as function of $T/\tau = N$.

them. The Hamiltonian in the rotating frame with resonant laser frequencies is

$$H = -(\Omega_1(t)|1\rangle\langle e| + \Omega_2(t)|2\rangle\langle e|) + h.c. \qquad (16)$$

where $\Omega_1(t)$ and $\Omega_2(t)$ are the time-dependent Rabi frequencies and depend on the parameters of the lasers (amplitude and phase). The diagonalization of (16) gives two eigenstates $|B_\pm\rangle = \frac{1}{\sqrt{2}\Omega}(\pm\Omega|e\rangle + \Omega_1|1\rangle + \Omega_2|2\rangle$ (called *bright states*) respectively with eigenvalues $\pm\Omega(t) = \pm\sqrt{\sum_{i=1}^{2}|\Omega_i|^2}$, and an eigenstate $|D\rangle = 1/\Omega(\Omega_2|1\rangle - \Omega_1|2\rangle)$ (called *dark state*) with zero eigenvalue. In the adiabatic evolution (i.e. when the Ω_i's change slowly and $\Omega T \gg 1$) it follows from the adiabatic theorem [26] that if the system starts at time $t = 0$ in an eigenstate of $H(0)$ (*dark* or *bright* state) during all the evolution it will remain in the eigenstate of $H(t)$ with the same eigenvalue.

Now we provide a simple example of the foregoing. Suppose that $\Omega_1(t)$ and $\Omega_2(t)$ are such that Ω is time-independent and $\Omega_1(0) = 0$ and $\Omega_2(0) = \Omega$. Moreover, suppose that the initial state is $|1\rangle$. Then *slowly* turn on the first Rabi frequency and turn off the second one. The system will always be in the *dark state* $|D\rangle$ and, at the end of the evolution (i.e., when $\Omega_1(T) = \Omega$ and $\Omega_2(T) = 0$), we will be in $|2\rangle$ state and have achieved population transfer from $|1\rangle$ to $|2\rangle$.

Consider now the case in which we have not a complete control of the laser field but the Rabi frequencies can fluctuate $\Omega_i \to \Omega_i + \delta\Omega_i$ (where the $\delta\Omega_i$ are independent and Gaussian distributed with zero mean and variance σ^2). This effect can produce errors in population transfer scheme for two reason: in general noisy perturbations may yield to significantly different output state and fast fluctuations could break the adiabatic approximation leading to transition to undesired (*bright*) states.

Numerical simulations show that, again, in the fast fluctuation regime the noise effects cancel out. In Figure 4 we present the population evolution of $|1\rangle$, $|2\rangle$ and $|e\rangle$ states subject to noisy evolution during the $|1\rangle \to |2\rangle$ *coherent adiabatic transfer* [28]. More precisely, we start from $|1\rangle$ and, during the evolution, the $|2\rangle$ is populated; at the end only the $|2\rangle$ is present. The $|e\rangle$ is never populated because of the high value of ΩT parameter. These simulations are done using $\Omega T = 1000$. Since we are in the adiabatic regime we are sure that the errors present are those induced by the perturbations. In Figure 4 we show the population evolution when the system is subject to noise with different correlation time τ (and different $N = T/\tau$). For $N = 10^4$ the transfer operation is not precise (i.e. the $|2\rangle$ state is not completely populated) and the $|e\rangle$ state is populated; this is a sign of the breaking of the adiabatic approximation due to noise. For $N = 10^5$ the evolution is much more similar to the ideal one: $|2\rangle$ is completely populated and $|e\rangle$ never appears during the evolution.

The above results can be explained calculating the amplitude transition from the *dark* to the *bright* state in presence of perturbations. The standard rule (see, e.g. [27]) to calculate the probability amplitude of a transition from the n state to one of the k state ($k \neq n$) at time t is $a_k = \sum'_n \int_{-\infty}^{t} \frac{a_n(t)}{\omega_{kn}} \frac{\partial V_{kn}}{\partial t} exp(i\int_0^t \omega_{kn}t')dt'$ where $\sum'$ indicates that the term $n = k$ is omitted, $\omega_{kn} = E_k - E_n$ and V_{kn} is the matrix element associate to the transition $n \to k$. In our case, the initial state is the *dark state* and the final states are the *bright states*. The eigenvalues $\pm\Omega$ are constant and then $\omega_{kn} = \Omega$. The perturbation Hamiltonian for time $j\tau \leq t \leq (j+1)\tau$ in the $|e\rangle$, $|1\rangle$, $|2\rangle$ basis is $\delta V^j = \sum_{i=1}^{2} \delta\Omega_i^j |i\rangle\langle e| + h.c.$ and in the new (*dark-bright states*) basis the relevant matrix element are

$$V^j_{DB_+} = -V^j_{DB_-} = \sum_j \frac{\delta\Omega_2^j\Omega_1 - \delta\Omega_1^j\Omega_2}{\sqrt{2}\Omega} S(j\tau,(j+1)\tau) \qquad (17)$$

To calculate the matrix element $\frac{\partial V_{kn}}{\partial t}$ we must take into account that $\partial S(j\tau,(j+1)\tau)/\partial t = \delta(t - j\tau) - \delta(t - (j+1)\tau)$. Let us focus our attention only on the first term in (17), by differentiating it we obtain $1/(\sqrt{2}\Omega)\sum_j \delta\Omega_2^j(\partial\Omega_1/\partial t)S(j\tau,(j+1)\tau) - \Omega_1\delta\Omega_2^j(\delta(t - j\tau) - \delta(t-(j+1)\tau)$. The terms $\partial\Omega_i/\partial t$ representing the adiabatic driven evolution are very small and can be neglected. Inserting this result in the expression for a_k and performing the integration we have

$$\begin{aligned} a_k &= \frac{1}{\sqrt{2}\Omega^2}\sum_j \delta\Omega_2^j[a_n(j\tau)\Omega_1(j\tau)e^{i\Omega j\tau} \\ &- a_n((j+1)\tau)\Omega_1((j+1)\tau)e^{i\Omega(j+1)\tau} + \\ &+ \text{terms with } (\Omega_1 \leftrightarrow \Omega_2) \end{aligned} \qquad (18)$$

If a_n and Ω_i change slowly i.e., $a_n((j+1)\tau) \approx a_n(j\tau)$ and $\Omega_i((j+1)\tau) \approx \Omega_1(j\tau)$. The exponential terms can be simplified to obtain $\exp(i\Omega\tau(j+1/2))\sin(\Omega\tau/2)/(2i)$

and eq. (18) for $\tau = T/N \ll 1$ gives

$$a_k = \frac{T}{2\sqrt{2}i\Omega} \sum_j e^{i\Omega j\tau} a_n(j\tau) \left[\Omega_1(j\tau) \frac{\delta\Omega_2^j}{N} - \Omega_2(j\tau) \frac{\delta\Omega_1^j}{N} \right] \tag{19}$$

The sums of $\frac{\delta\Omega_i^j}{N}$ converge to the mean of $\delta\Omega_i$ that is to zero; the other factors are bounded ($0 \leq |a_n| \leq 1$ and $0 \leq |\Omega_i| \leq \Omega$) and by consideration similar to those at the end of section II C (and [18]) we can conclude that $a_k \to 0$ for $N \to \infty$. In our case, $|n\rangle = |D\rangle$, $|k\rangle = |B_\pm\rangle$ and $|a_{B_\pm}|^2 \to 0$ as $N \to \infty$: the transition from *dark* to *bright states* is suppressed in the fast fluctuating regime and the evolution happen in the *dark space*.

To have a more detailed picture in Figure 5 we show also the trend of the *fidelity* (upper curve) and the relative average population of the $|e\rangle$ state (lower curve) as functions of T/τ. The trends of the two curves are correlated, which is suggesting that the main source of error in the operation is the population of excited state due to loss of the adiabatic approximation. As expected, because of the cancellation effect, for great T/τ the *fidelity* approaches 1 even in presence of strong noise and the $|e\rangle$ state is not populated.

These results not only can explain why the *adiabatic population transfer* scheme is robust against field fluctuation but can give information for the experimental set-up. In fact the adiabatic evolution is, in general, arbitrarily long; once the experimental parameters are fixed (as the laser with its proper noise correlation time τ) we can prolong the evolution time in order to increase T/τ and let the noise average out. As shown before, for every noise strength σ and correlation time τ we can find and evolution time T in order to obtain the desired *fidelity*: that is, to achieve the population transfer with arbitrary small error.

IV. CONCLUSION

We studied the effect of stochastic noise on several quantum systems. The noise is described by a Gaussian stochastic process with zero mean superposed to the ideal quantum evolution. For all of the systems we found that for fast fluctuating noise a *cancellation effect* appears: the noise fluctuations average out leading the system to a state near to the ideal one.

We showed by analytical and numerical calculation how this effect can appear in a two level system and propose an experiment to verify the presence of this cancellation regime. The experiment is based on the one performed in Ref. [14] and we think that, with a modification of the experimental set-up, it could be easily performed.

We applied the same model to another important technique in atoms physics : the *adiabatic population transfer*. We explained how this effect leads to the robustness of the adiabatic process against the perturbation noise. This can be important for the experiments using the *adiabatic population transfer*.

Acknowledgments

P.S. wishes to thank D. Kielpinski for useful comments.

[1] C.H. Bennett and G. Brassard *Proceedings of IEEE International Conference on Computers, Systems and Signal Processing* 175-179, IEEE, New York, 1984. C. H. Bennett *Phys. Rev. Lett.* **68**(21), 3121, 1992.

[2] C.H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wootters Phys. Rev. Lett. **70**, 1895, 1993.

[3] P.W. Shor *Proceeding of 35th Annual symposium on foundation of Computer Science*. (IEEE Computer Society Press, Los Alamitos, CA, 1994). L.K. Grover *Proc.* 28^{th} *Annual ACM Symposium on the Theory of Computation*, 212, ACM Press, New York, 1996.

[4] P. Zanardi and M. Rasetti, Phys. Rev. Lett. **79**, 3306 (1997).

[5] P.W. Shor Phys. Rev.A **52**, 2493 (1995); A.M. Steane Phys. Rev. Lett. **77**, 793 (1996); E. Knill, R. Laflamme, Phys. Rev.A **55**, 900 (1997) and references therein.

[6] L. Viola and S. Lloyd, Phys. Rev. A. **58**, 2733 (1998); L. Viola, E. Knill, and S. Lloyd Phys.Rev.Lett. **82**, 2417 (1999); D. Vitali, P. Tombesi, Phys. Rev. A **65**, 012305 (2002); P. Zanardi, Phys. Lett. A **258** 77 (1999).

[7] J.A. Jones *et al.* , Nature **403**, 869 (2000). G. Falci *et al.* , Nature **407**, 355 (2000). P. Zanardi and M. Rasetti, Phys. Lett.A **264**, 94 (1999). G. Falci *et al.* , Nature **407**, 355 (2000). R.G. Unanyan, B.W. Shore and K. Bergmann, Phys. Rev. A **59**, 2910 (1999). L.-M. Duan,J. I. Cirac and P. Zoller, Science **292**, 1695 (2001). L. Faoro, J. Siewert and R. Fazio, Phys. Rev. Lett. **90**, 028301 (2003). I. Fuentes-Guridi *et al.* Phys. Rev. A **66**, 022102 (2002). A. Recati *et al.* Phys. Rev. A **66**, 032309 (2002). P. Solinas *et al.* Phys. Rev. B **67**, 121307 (2003). P. Solinas *et al.* Phys. Rev. A **67**, 062315 (2003)

[8] WangXiang-Bin, M. Keiji Phys. Rev. Lett. **87**, 097901 (2001); WangXiang-Bin, M. Keiji Phys. Rev. Lett. **88**, 179901(E) (2002). X.-Q. Li *et al.* Phys. Rev. A **66**, 042320 (2002). S. L. Zhu, Z.D. Wang, Phys. Rev. Lett. **89**, 097902 (2002). P.Solinas *et al.* Phys. Rev. A **67**, 052309 (2003)

[9] A. Blais and A.-M. S. Tremblay Phys. Rev. A 67, 012308 (2003); A. Nazir, T. P. Spiller, and W. J. Munro, Phys. Rev. A 65, 042303 (2002); G. De Chiara, G.M. Palma, Phys. Rev. Lett. **91**, 090404 (2003); A. Carollo *et al*, Phys. Rev. Lett. 90, 160402 (2003); A. Carollo *et al* quant-ph/0306178; V.I. Kuvshinov, A.V. Kuzmin, Phys. Lett. A, **316**, 391 (2003); F. Gaitan, quant-ph/0312008.

[10] P. Solinas, P. Zanardi, N. Zanghì quant-ph/0312109.

[11] P. Facchi, S. Montangero, R. Fazio, S. Pascazio quant-ph/0407098.

[12] Our sampling set of the Bloch sphere (i.e. our inizial state space) is given by $\{\pm e_i\}_{i=x,y,z}, (\pm e_x \pm e_y)/\sqrt{2}, (\pm e_x \pm e_z)/\sqrt{2}, (\pm e_y \pm e_z)/\sqrt{2}$. Here e_i denotes the normalized vector of the i-th direction.

[13] We make simulations with 1000 different realizations and average the resulting *square fidelity*. A part from a noisier in the curve in the plots, the results are weakly affected by this parameters so that with 100 realizations we obtain the almost the same results.

[14] D. Kielpinski *et al.* Science **291**, 1013, (2001).

[15] M.A. Nielsen and I.L. Chuang *Quantum computation and information*, Cambridge University Press (2000).

[16] D.M Meekhof *et al.*, Phys. Rev. Lett. **76** 1796, 1996; S. Schneider and G.J. Milburn Phys. Rev. A **57**, 3748, (1998); M. Murao and P.L. Knight Phys. Rev. A **58**, 663 (1998).

[17] M.O. Scully and M.S. Zubairy *Quantum Optics*, Cambridge University Press (1997).

[18] This is an elementary instance of the law of large numbers: Suppose to have a sum written in the form $\sum_{j=0}^{N} f(\tau j)\frac{x_j}{N}$ where x_j are independent random variable with zero mean and variance σ^2, $\tau = T/N$ and $0 \leq j \leq T$. $Var(Y_N)$ (the variance of $Y_N = \frac{1}{N}\sum_{j=0}^{N} f(\tau j)x_j$) is $\frac{\sigma^2}{N^2}\sum_{j=0}^{N}(f(\tau j))^2$. Thus, if f is bounded in the interval $[0,T]$, $Var(Y_N) \to 0$ when $N \to \infty$ and therefore $Y_N \to 0$ since the x_j mean is zero. Since in our case $f(\tau j) = exp(-i2B_z\tau j)$ the second terms in eq. (15) goes to zero and $\mathcal{F} \to 1$ for $N \to \infty$ for every initial state (i.e. independently from α and β).

[19] K. Bergmann, H. Theuer, and B.W. Shore Rev. Mod. Phys. **70**, 1003 (1998).

[20] P. Dittmann *et al.* J. Chem. Phys. **97**, 9472-9475 (1992).

[21] S. Kulin *et al.* Phys. Rev. Lett. **78**, 4815-4188 (1997).

[22] M. Weitz, B. C. Young, and S. Chu Phys. Rev A **50**, 2438-2444, (1994).

[23] A.S. Parkins *et al.* Phys. Rev. Lett. **71**, 3095-3102, (1993)

[24] R. Walser, J.I. Cirac, and P. Zoller Phys. Rev. Lett. **77**, 2658-2661, (1996).

[25] L.P. Yatsenko, G. Guérin, and H.R. Jauslin quant-ph/01007065.

[26] A. Messiah *Mecanique Quantique*, Dunod Paris (1964).

[27] L.I. Schiff *Quantum Mechanics*, McGrawn-Hill (1968).

[28] We are interested in the $|1\rangle \to |2\rangle$ transfer then we start form $|1\rangle$ and do not sample the initial state space. The *fidelity* is still obtained averaging the results of 1000 different simulations.