

# Quantum Error Correction in Correlated Quantum Noise

Rochus Klesse and Sandra Frank

*Universität zu Köln, Institut für Theoretische Physik, Zùlpicher Str. 77, D-50937 Köln, Germany*

(Dated: May 20, 2005)

We consider quantum error correction of quantum noise that is created by a local interaction of qubits with a common bosonic bath. The possible exchange of bath bosons between qubits gives rise to spatial and temporal correlations in the noise. We find that these kind of noise correlations have a strong negative impact on quantum error-correction.

PACS numbers: 03.67.Pp, 03.65.Yz, 03.67.Hk, 03.67.Lx

The superiority of quantum computation over conventional computation relies on the fact that a quantum-bit (qubit) register can be in the superposition of a very large number of classical computational states. At the same time, maintaining coherence of this highly superpositional state poses also the main obstacle for the realization of a quantum computer.

For a small number of qubits this difficulty can be overcome by simply reducing the coupling to environmental degrees of freedom, as has been demonstrated by several groups for different physical implementations. With an increasing number of qubits, it will, however, become extremely difficult to reach the required coherence in that way [1]. It is therefore common opinion that a scalable implementation of a quantum computer must use some error correction scheme that recovers the quantum state after it has been distorted by external noise.

The existence of error correcting schemes for quantum states, which was shown independently by Shor [2] and Steane [3], is a remarkable fact and has been crucial for the development of the field. The key ideas presented in their work rapidly evolved to a beautiful theory of quantum error correcting codes and subsequently to the concept of fault tolerant quantum computation [4].

Quantum error correcting schemes are usually designed for the independent error model, which by definition does not exhibit correlations between noise of different times and locations. From a physical point of view, this requirement is rather annoying, since in general qubits do interact with a *common* environment which necessarily introduces some amount of correlations in the noise. To be more specific, in many if not all situations the qubits weakly interact with a common thermal bath of extended bosons (photons and/or phonons). The exchange of bosons between qubits will then cause spatial and temporal error correlations that violate the condition of error independence. Indeed, it has been shown [5] that these kind of processes can lead to drastically enlarged or reduced decoherence of certain states.

To which extent do such error correlations interfere with quantum error correction? We have analyzed this problem for optimal Calderbank-Shor-Steane (CSS) quantum error-correcting codes of variable length  $n$  (number of physical qubits) and size  $k$  (number of log-

ical qubits). As physical noise-model, we use a reduced spin-boson model consisting of  $n$  spins – describing an  $n$ -qubit register – coupled to a common bosonic bath [5, 6]. The amount of noise correlations can be controlled by the inter-spin distance  $r$ . Within this framework, we study how code states transform during spin-boson interaction and a subsequent error-correcting operation. The distance between the resulting code state and the initial one – in the sequel denoted as residual error  $\Delta$  [cf. Eq. (11)] – serves as a measure for the error-correcting performance of the code.

Our main finding is that quantum error correction with CSS codes is substantially hampered by the kind of noise-correlations captured in our model. This becomes evident by the fact that for any fixed information rate  $k/n > 0$  the residual error  $\Delta$  approaches a finite constant in the limit  $n \rightarrow \infty$ , unless the spin-boson coupling strictly vanishes or  $r$  is infinite (cf. Fig. 1). Using a simple scaling argument we conclude that for a wide range of model parameters CSS codes cannot provide the accuracy needed for large scale quantum computations.

In contrast to related studies [7], here the spin-boson coupling is treated in a non-perturbative manner, which we find to be indispensable in the large  $n$  limit. Substantial progress towards error correction beyond the independent error model has been made in very recent work [8, 9]. We will briefly comment our results in light of this new work at the end of this Letter. An extended discussion will follow in a future publication [10].

We begin with the physical model for the  $n$ -qubit register. It is defined by the Hamiltonian

$$H = \sum_{l=0}^{n-1} \frac{\epsilon}{2} \sigma_{z,l} + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^{\dagger} b_{\mathbf{k}} + \sum_{\mathbf{k},l} \sigma_{z,l} (g_{|\mathbf{k}|} e^{i\mathbf{k} \cdot \mathbf{r}_l} b_{\mathbf{k}}^{\dagger} + \text{H.c.})$$

of  $n$  spin- $\frac{1}{2}$  particles (qubits) at positions  $\mathbf{r}_l$  interacting with three-dimensional bosonic modes with creation (annihilation) operators  $b_{\mathbf{k}}^{\dagger}$  ( $b_{\mathbf{k}}$ ) and energies  $\omega_{\mathbf{k}} = |\mathbf{k}|$  [11].  $\sigma_{z,l}$  denotes the Pauli spin  $\sigma_z$  operator acting on the  $l$ th spin, and  $\epsilon$  is the Zeeman energy. The spin-boson couplings  $g_{|\mathbf{k}|}$  may be characterized as usual by a spectral function  $J(\omega) := \sum_{\mathbf{k}} \delta(\omega_{\mathbf{k}} - \omega) |g_{|\mathbf{k}|}|^2 \equiv A \omega^s e^{-\omega/\Omega}$ , with a cut-off frequency  $\Omega$ , a constant  $A$  of appropriate dimension, and a positive parameter  $s$  [12]. Note that the

spin-boson couplings do not lead to energy dissipation but to dephasing of the spin system.

The spin-boson model given by Hamiltonian  $H$  is quite suitable for our purposes because it shows full decoherence and is still analytically solvable [5, 6]. We assume that initially the system is in a product state  $\rho(0) \otimes \rho_b$  of a spin state  $\rho(0)$  and a thermal state  $\rho_b$  of the bosonic bath at temperature  $T$ . Following Refs. [5, 6] we determine the reduced density matrix  $\rho(\tau)$  of the spin system at some subsequent time  $\tau$  to be [10]

$$\rho(\tau) = \sum_{\eta\mu \in \mathbf{Z}_2^n} e^{-C_{\eta\mu}} P_\eta \rho(0) P_\mu, \quad (1)$$

where  $P_\eta$  for  $\eta \in \mathbf{Z}_2^n \equiv \{0, 1\}^n \equiv \{\uparrow, \downarrow\}^n$  is the projector on the state  $|\eta\rangle = |\eta_0\rangle \dots |\eta_{n-1}\rangle$ , and  $C_{\eta\mu}$  are real [13] coefficients given by

$$C_{\eta\mu} = \sum_{lm} (\eta_l - \mu_l)(\eta_m - \mu_m) \Gamma_{|\mathbf{r}_l - \mathbf{r}_m|}.$$

The distance-dependent decoherence-parameter [10]

$$\Gamma_r = A \int_0^\infty d\omega \omega^s \frac{1 - \cos \omega \tau}{\omega^2} \coth\left(\frac{\omega}{2T}\right) \frac{\sin \omega r}{\omega r} e^{-\omega/\Omega} \quad (2)$$

is a positive, monotonously decaying function of  $r$  if  $s < 2$ . Up to an singularity at  $r = \tau$ , the same holds for  $2 \leq s < 3$  if temperature is high,  $T \gg \tau^{-1}, r^{-1}$  [10]. We confine our considerations to  $s < 3$  and note that in quantum optical settings typically  $s = 1$ , the Ohmic case, where  $\Gamma_r(\tau)$  is linear in  $\tau$  for large  $\tau \gg 1/T$  [12].

To make the model manageable we simplify the coefficients  $C_{\eta\mu}$  by setting all distances  $|\mathbf{r}_l - \mathbf{r}_m|_{l \neq m}$  to the maximum distance  $r$ . Keeping in mind that this simplification lowers the effects of correlations, we obtain

$$C_{\eta\mu} = |\eta \oplus \mu|(\Gamma_0 - \Gamma_r) + (|\eta| - |\mu|)^2 \Gamma_r, \quad (3)$$

where  $|\eta|$  denotes the Hamming weight [14] of  $\eta$ , and  $\oplus$  means a bitwise addition modulo 2 in  $\mathbf{Z}_2^n$ .

Although representation Eq. (1) looks simple, it turns out to be rather cumbersome for our further calculations. A much better one can be given in terms of operators  $Z_\nu$  that are defined for  $\nu = (\nu_0 \dots \nu_{n-1}) \in \mathbf{Z}_2^n$  as product of exactly those  $\sigma_{z,l}$  where  $\nu_l = 1$ . By the structure of the interaction Hamiltonian it is evident that then

$$\rho(\tau) = \sum_{\nu, \nu' \in \mathbf{Z}_2^n} \alpha_{\nu\nu'}(\tau) Z_\nu \rho(0) Z_{\nu'}, \quad (4)$$

with suitable coefficients  $\alpha_{\nu\nu'}(\tau)$ . To determine them we let  $\rho(0) = |X\rangle\langle X|$  be the projector on the totally  $x$  polarized spin state and compare the expressions obtained in both representations. In this way we see that  $\alpha_{\nu\nu'}$  and  $e^{-C_{\eta\mu}}$  are related by a Fourier transformation in  $\mathbf{Z}_2^n \times \mathbf{Z}_2^n$ ,

$$\alpha_{\nu\nu'} = \frac{1}{4^n} \sum_{\eta\mu \in \mathbf{Z}_2^n} (-1)^{\nu \cdot \eta + \nu' \cdot \mu} e^{-C_{\eta\mu}} \quad (5)$$

( $\nu \cdot \eta$  denotes the standard inner product in  $\mathbf{Z}_2^n$ ). Later on we will only need the diagonal coefficients  $\alpha_{\nu\nu}$ , which solely depend on the Hamming weight  $|\nu|$ . For  $C_{\eta\mu}$  according to Eq. (3) the sum (5) can be easily carried out if we eliminate the squared term in the exponent by the identity  $e^{-(|\eta| - |\mu|)^2 \Gamma} = \frac{1}{\sqrt{\pi\Gamma}} \int dx e^{-\frac{x^2}{\Gamma} + 2ix|\eta| - 2ix|\mu|}$ . After some algebra we arrive at

$$\alpha_{\nu\nu} \equiv \beta_{|\nu|} = \int \frac{dx}{\sqrt{\pi\Gamma_r}} e^{-\frac{x^2}{\Gamma_r}} p_x^{|\nu|} (1 - p_x)^{n-|\nu|}, \quad (6)$$

where we introduced the  $x$ -dependent “probability”

$$p_x = (1 - e^{-\Gamma_0 + \Gamma_r} \cos 2x)/2 \in [0, 1]. \quad (7)$$

Thus, we have a convenient description of the decoherence process at hand and can now turn to CSS codes.

The key idea of quantum error-correction is to encode the information of  $k$  logical qubits in an appropriate subspace  $\mathcal{C}$  of the Hilbert space  $\mathcal{H}_n$  associated with  $n > k$  physical qubits.  $\mathcal{C} \subset \mathcal{H}_n$  is called a quantum code of length  $n$  and size  $k$ . Error operations that unitarily map  $\mathcal{C}$  on different cosets of  $\mathcal{C}$  can then be detected and corrected. A CSS quantum code is constructed on two linear classical codes [14] (i.e.  $\mathbf{Z}_2$ -linear spaces)  $C_2 \subset C_1 \subset \mathbf{Z}_2^n$  by

$$\mathcal{C} = \text{span}\{|Q\rangle\}_{Q \in C_2^\perp / C_1^\perp} \subset \mathcal{H}_n$$

where  $C_i^\perp$  is the orthogonal space of  $C_i$  with respect to the inner product in  $\mathbf{Z}_2^n$ , and the state vectors  $|Q\rangle$  are

$$|Q\rangle = \frac{1}{\sqrt{|C_1|}} \sum_{y \in C_1} Z_q |y\rangle, \quad q \in Q.$$

$\mathcal{C}$  encodes  $k = \log_2 |C_2^\perp / C_1^\perp| = \dim C_1 - \dim C_2$  logical qubits in  $n$  physical qubits [4, 15]. The error correcting capability of  $\mathcal{C}$  is determined by the minimum weights [14]  $d_1$  and  $d_1^\perp$  of  $C_1$  resp.  $C_1^\perp$ . An error correction scheme using  $\mathcal{C}$  can correct up to  $t = \lfloor \frac{d-1}{2} \rfloor$  universal qubit errors, where  $d = \min\{d_1, d_1^\perp\}$ . This characterizes  $\mathcal{C}$  as an  $[n, k, d]$  code. With  $P$  being the projector on  $\mathcal{C}$ , the error correcting operation  $\mathcal{R}$  associated with  $\mathcal{C}$  reads

$$\mathcal{R}(\rho) = \sum_{\nu, \mu \in \mathbf{Z}_2^n, |\nu|, |\mu| \leq t} P X_\mu Z_\nu \rho Z_\nu X_\mu P \quad (8)$$

( $X_\mu$  is analogously defined as  $Z_\nu$  with  $\sigma_{x,l}$  instead of  $\sigma_{z,l}$ ).

CSS codes have been used to demonstrate the existence of efficient (“good”) quantum error correcting codes [15], as specified by a theorem of Calderbank and Shor:

*Theorem (Calderbank and Shor):* For sufficiently large  $n$  there exists always an  $[n, k, d]$  CSS code satisfying

$$k/n > 1 - 2H_2(d/n) \equiv R_{css}(d/n). \quad (9)$$

( $H_2(x) = -x \log_2 x - (1-x) \log_2 (1-x)$  is the binary entropy function.) Provided that the error-correcting operation  $\mathcal{R}$  for such codes is error-free, this theorem can




FIG. 1: Residual error  $[\Delta]_t^n$  of  $[n, k, d = 2t + 1]$  CSS codes as function of  $t$  for  $\Gamma_0 = 0.01$ ,  $k/n = 0.062$ , and  $t/n = 0.05$ . The values of  $\Gamma_r$  are .01, .005, .0025, .00125 (squares, top to down), and  $\Gamma_r = 0$  (dots). The dashed lines are the asymptotic values of the residual errors.

be rephrased in a more pragmatic fashion: An  $n$ -qubit register that is perturbed in at most  $t$  qubits can be used to perfectly restore  $k = \lfloor n R_{css}(\frac{2t+1}{n}) \rfloor$  logical qubits that have been encoded in an appropriate  $[n, k, 2t + 1]$  CSS code.

Of course, a real and noisy physical  $n$ -qubit register is unlikely to guarantee the prerequisite of the theorem. It therefore does not necessarily provide a practical solution of the decoherence problem. One has to demonstrate that also under more realistic conditions the performance of the error-correcting code is still sufficient. This has been shown in refs. [15] for the independent error model. We will now investigate how CSS error correction performs when the errors are correlated.

To this end we consider an arbitrary state  $|\Psi\rangle_L = \sum_{\eta} \psi_{\eta} |\eta\rangle_L$  of an abstract  $k$ -qubit register with  $2^k$  orthonormal logical basis states  $|\eta\rangle_L$ ,  $\eta \in \mathbf{Z}_2^k$ . The logical state  $|\Psi\rangle_L$  is encoded in an  $[n, k, d]$  CSS code  $\mathcal{C}$  as a vector  $|\Psi\rangle = \sum_{\eta} \psi_{\eta} |Q_{\eta}\rangle \in \mathcal{C}$  according to some bijective mapping  $|\eta\rangle_L \mapsto |Q_{\eta}\rangle$ . The encoded state  $\rho_{\Psi} = |\Psi\rangle\langle\Psi|$  is first subjected to the noise operation defined by Eq. (4) and then corrected by  $\mathcal{R}$ . This results in a final state  $\rho'_{\Psi} = \mathcal{R}(\rho_{\Psi}(\tau))$  which we compare with the original state  $\rho_{\Psi}$  by the fidelity  $F(\rho_{\Psi}, \rho'_{\Psi})$ , which here is

$$\langle\Psi|\rho'_{\Psi}|\Psi\rangle = \sum_{|\mu|\leq t} \sum_{\nu\nu'} \alpha_{\nu\nu'} \langle\Psi|Z_{\mu+\nu}|\Psi\rangle \langle\Psi|Z_{\mu+\nu'}|\Psi\rangle. \quad (10)$$

Since the square root of  $1 - F$  defines a proper distance measure for density matrices [16], we call

$$\Delta_{\Psi} \equiv 1 - F(\rho_{\Psi}, \rho'_{\Psi}) \quad (11)$$

the residual error after CSS error correction.

Here we are not interested in specific codes but in the general properties of CSS codes. We therefore continue by taking the average over practically all good CSS codes of a given length  $n$  and a size  $k$ . This is technically possible because of the following

*Theorem: For any positive  $\varepsilon$  the  $[n, k, d]$  CSS code associated with randomly chosen classical codes  $C_2 \subset C_1 \subset$*

$\mathbf{Z}_2^n$  with  $\dim C_2 = \lfloor \frac{n-k}{2} \rfloor$  and  $\dim C_1 = \lfloor \frac{n+k}{2} \rfloor$  satisfies

$$k/n \geq (1 - \varepsilon) R_{css}(d/n)$$

with a probability larger than  $1 - 2^{-n(\alpha\varepsilon + \mathcal{O}(n^{-1}))}$ , where  $\alpha$  is a positive constant independent of  $n$  and  $k$ .

The situation is thus very much like the one in classical coding theory: randomly chosen subspaces of  $\mathbf{Z}_2^n$  yield codes that asymptotically reach the Gilbert-Varshamov bound [14]. The theorem can be proven [10] along the same lines as the proof in [17] for the classical case, plus application of MacWilliams' theorem [14].

Let an average over all CSS codes of length  $n$  and size  $k$  be defined via the uniform average over all pairs  $C_2 \subset C_1$  of linear subspaces in  $\mathbf{Z}_2^n$  with dimensions  $\dim C_{2/1} = \lfloor \frac{n+k}{2} \rfloor$ . According to the theorem, this average can be understood as an average over essentially all CSS codes of length  $n$  that asymptotically correct up to  $t$  qubit errors, where  $t = \lfloor \frac{d-1}{2} \rfloor$  is determined by  $k/n = R_{css}(d/n)$ . We denote this average by  $[\dots]_t^n$ .

Applying the average to Eq. (10) leads us finally to

$$[\Delta_{\Psi}]_t^n = \sum_{|\nu|>t} \alpha_{\nu\nu} = \sum_{w=t+1}^n \binom{n}{w} \beta_w, \quad (12)$$

where we used the completeness relation  $\sum_{\nu} \alpha_{\nu\nu} = 1$ , and suppressed terms of order  $2^{-|\mathcal{O}(n)|}$ . The physical interpretation of this expression is that more than  $t$  simultaneous qubit errors cannot be corrected, and therefore coefficients  $\alpha_{\nu\nu}$  with  $|\nu| > t$  contribute to the residual error. Terms with non-diagonal coefficients  $\alpha_{\nu\neq\nu'}$  turn out to be suppressed by a factor  $2^{-\frac{n+k}{2}}$  and therefore do not significantly contribute. Up to these exponentially small corrections the code averaged residual error is independent of the encoded logical state  $|\Psi\rangle_L$ .

In the following we consider  $[\Delta]_t^n$  for a fixed ratio  $q \equiv \frac{t+1}{n}$ . Inserting Eq. (6) into (12) immediately leads to

$$[\Delta]_{qn}^n = \int \frac{dx}{\sqrt{\pi\Gamma_r}} e^{-\frac{x^2}{\Gamma_r}} \sum_{w=qn}^n \binom{n}{w} p_x^w (1 - p_x)^{n-w}. \quad (13)$$

From this general formula for the residual error we can now draw conclusions on the performance of CSS codes.

Independent errors correspond to the case of diverging distance  $r$ , where the decoherence parameter  $\Gamma_r$  vanishes. In this limit the Gaussian in Eq. (13) shrinks to a normalized delta peak at  $x = 0$ . Hence,

$$[\Delta]_{qn}^n = \sum_{w=qn}^n \binom{n}{w} p_o^w (1 - p_o)^{n-w}, \quad (14)$$

where  $p_o = (1 - e^{-\Gamma_o})/2$  is the error probability of a single spin. This expression, formerly derived by Calderbank and Shor [15], implies a binomially distributed residual error that decays exponentially with  $n$  as long as  $p_o < q$ . The point is that the constraint on the single-spin decoherence  $\Gamma_0 \approx 2p_o < 2q$  is  $n$ -independent. In this

sense, CSS codes provide scalable quantum-error correction with an exponentially small residual error [15].

For finite  $\Gamma_r$  the asymptotic value of the residual error can be easily derived from Eq. (13) by the observation that in the limit  $n \rightarrow \infty$  the sum over  $w$  vanishes if  $p_x < q$  and equals unity if  $p_x > q$ . We obtain

$$\lim_{n \rightarrow \infty} [\Delta]_{qn}^n = \int_{p_x > q} \frac{dx}{\sqrt{\pi \Gamma_r}} e^{-\frac{x^2}{\Gamma_r}} \sim \operatorname{erfc} \sqrt{\frac{q}{\Gamma_r}}, \quad (15)$$

where the last approximation is good for  $\Gamma_r, \Gamma_0 \ll q$ . In sharp contrast to the independent errors discussed above for any finite  $\Gamma_r$  (i.e., for any finite  $r$ ) the residual error converges for large  $n$  to a finite constant (cf. Fig. 1).

According to quantum complexity theory [18] the total error in an  $m$  step quantum computation must be less than  $m^{-|\mathcal{O}(1)|}$  in order to produce useful results. If we therefore demand that the residual error must be limited by a  $\Delta_{max}(n) \sim b/n^\mu$  with some positive  $b$  and  $\mu$ , we deduce from (15) that  $\Gamma_r$  must obey

$$\Gamma_r < \Gamma_{max}(n) \sim q/(\text{const} + \mu \ln n). \quad (16)$$

This is no longer independent of  $n$  as for the uncorrelated model but approaches zero in the limit  $n \rightarrow \infty$ , although slowly with the inverse logarithm of  $n$ .

A further discussion must take into account that also  $\Gamma_r$  by Eq. (2) depends on  $n$  via the  $n$ -dependence of the maximum distance  $r$  and the observation time  $\tau$ . It is plausible to assume  $r \propto n^y$  with some exponent  $y \geq 1/3$ , and also  $\tau \propto n^y$ , because the observation time  $\tau$  must scale at least linearly with the time needed to transmit signals between qubits. We therefore set  $r = ar_0$ ,  $\tau = a\tau_0$ , where  $a = (n/n_0)^y$ , and write

$$\Gamma_r \equiv \Gamma(s, ar_0, a\tau_0, T, \Omega) = a^{1-s} \Gamma(s, r_0, \tau_0, aT, a\Omega).$$

The second equality is obtained by rescaling the integral (2). Inspection of Eq. (2) also reveals that for large effective temperatures  $aT \gg r_0^{-1}, \tau_0^{-1}$  the decoherence parameter is linear in temperature, and virtually independent on  $\Omega$ . For large  $n$  condition (16) is therefore equivalent to

$$\Gamma(s, r_0, \tau_0, T) < \frac{q}{\text{const} + \mu \ln n} \left(\frac{n}{n_0}\right)^{y(s-2)}. \quad (17)$$

The inequality is violated if  $s \leq 2$  and  $n$  is large. Consequently, in this regime the residual error exceeds  $\Delta_{max}(n)$ . On the other hand, for  $s > 2$  conditions (17) and (16) are satisfied for large  $n$ , meaning that here a sufficiently small residual error can be attained. We conclude that for interaction parameter  $s > 2$  scalable CSS error-correction is possible, while for  $s \leq 2$  it is not.

Apparently, noise correlations introduced by the spin-boson interaction are a strong hindrance for small values of  $s$ , whereas they are less harmful for larger  $s$ . An intuitive explanation is that with increasing  $s$  spectral weight in  $J(\omega)$  is shifted from lower to higher frequencies, and

therefore the larger the value of  $s$ , the faster the disturbing noise correlations decay with distance.

Note that the physical model we use is lacking dissipative couplings, which corresponds to a systematic underestimation of quantum noise. So, for a spin-boson model including dissipative couplings the residual error will be larger than the one calculated here. A second caveat is that we considered perfect quantum error-correction, while real systems will have to rely on *fault-tolerant* error-correcting schemes. It is not clear to what extent the results presented here do apply to those schemes. However, a naive guess might be based on the observation that fault-tolerant quantum computation is, in some sense, simulating perfect quantum error-correction with imperfect quantum gates. This suggests that fault-tolerant schemes cannot better perform than perfect error-correction investigated here. Following these arguments, it is important to note that our results are not at odds with a recently proven threshold theorem [9]. Here the spin-boson interaction is unbounded [8] and therefore does not belong to the class of interactions considered in Ref. [9].

We wish to thank H. Moraal for numerous valuable discussions and M. R. Zirnbauer for critical reading of the manuscript. The work is supported by the SFB TR/12.

- 
- [1] W. Unruh, Phys. Rev. A **51**, 992 (1995).
  - [2] P. W. Shor, Phys. Rev. A **52**, R2493 (1995).
  - [3] A. M. Steane, Phys. Rev. Lett. **77**, 793 (1996).
  - [4] see eg D. Aharonov, in *Ann. Rev. of Comp. Physics*, ed. by D. Stauffer (World Scientific, Singapore 1998); E. Knill et al. Proc. R. Soc. London A **454**, 365 (1998); J. Preskill, *ibid.*, 385 (1998); D. Aharonov, M. Ben-Or, arXiv e-print quant-ph/9906129; M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (University Press, Cambridge, 2000);
  - [5] G. M. Palma, K. A. Suominen, A. K. Ekert, Proc. R. Soc. London A **452**, 567 (1996).
  - [6] L.-M. Duan and G.-C. Guo, Phys. Rev. A **57**, 737 (1998); J. H. Reina et al., Phys. Rev. A **65**, 032326 (2002).
  - [7] L.-M. Duan, G.-C. Guo, Phys. Rev. A **59**, 4058 (1999); W. Y. Hwang et al., Phys. Rev. A **63**, 022303 (2001); D. V. Averin, R. Fazio, JETP letters **78**, 664 (2003); J. P. Clemens et al., Phys. Rev. A **69**, 062313 (2004).
  - [8] B. M. Terhal, G. Burkard, Phys. Rev. A **71**, 012336 (2005).
  - [9] P. Aliferis et al., preprint quant-ph/0504218 (2005).
  - [10] R. Klesse, S. Frank, to be published.
  - [11] We use units in which  $\hbar = 1$ ,  $k_B = 1$  and  $c = \omega_k/|k| = 1$ .
  - [12] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (University Press, Oxford, 2002); D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer-Verlag Berlin Heidelberg New York, 1994).
  - [13] We omit the imaginary parts. They correspond to a coherence preserving *unitary* transformation of the spin state [6], which is of minor interest for our discussion.
  - [14] F. J. MacWilliams, N. J. A. Sloane, *Theory of Error-Correcting Codes* (North-Holland, Amsterdam, 1977).

- [15] A. R. Calderbank and P. W. Shor, Phys. Rev. A **54**, 1098 (1996); A. Steane, Proc. R. Soc. Lond. A **452**, 2551 (1996).
- [16] A. Gilchrist et al., Phys. Rev. A **71**, 062310 (2005).
- [17] J. T. Coffey et al. Discrete Appl. Math. **33**, 43 (1991).
- [18] E. Bernstein et al., SIAM J. Comput. **26**, 1411 (1997).