

On the Significance of a Recent Experiment Demonstrating Quantum Interference in Time

Lawrence P. Horwitz*

School of Physics, University of Tel Aviv, Ramat Aviv 69978, Israel

and

Department of Physics, College of Judea and Samaria, Ariel, Israel

Abstract

I discuss the interpretation of a recent experiment showing quantum interference in time. It is pointed out that the standard nonrelativistic quantum theory, used by the authors in their analysis, does not have the property of coherence in time, and hence cannot account for the results found. Therefore, this experiment has fundamental importance beyond the technical advances it represents. Some theoretical structures which consider the time as an observable, and thus could, in principle, have the required coherence in time, are discussed briefly, and the application of Floquet theory and the manifestly covariant quantum theory of Stueckelberg are treated in some detail. In particular, the latter is shown to account for the results in a simple and consistent way.

The recent experiment of Lindner, *et al*¹, clearly shows the effect of quantum interference in time for the wave function of a particle. The results are discussed in that paper in terms of a very precise solution of the time-dependent nonrelativistic Schrödinger equation. In this note, I wish to point out that the nonrelativistic Schrödinger theory cannot be used to predict interference phenomena in time, and therefore the very striking results of this beautiful experiment have a fundamental importance which goes beyond the technical advances which they represent. They imply, in fact, that the time variable t must be adjoined to the set of standard quantum variables so that the standard ket $|x, t\rangle$ for the representation of the quantum state (in Dirac's terminology²) can be constructed. It is this structure for the wave function $\psi(x, t) \equiv \langle x, t | \psi \rangle$, where x and t are the spectra of self-adjoint operators, that provides the possibility of coherence in t , and therefore, interference phenomena. If the quantum theory is to remain symplectic in form, the variable E must also be adjoined.

The standard nonrelativistic quantum theory cannot be used to predict interference in time. For example, Ludwig³ has pointed out that the time variable cannot be a quantum observable, since there is no imprimitivity system (i.e., no operator exists that does not commute with t in the nonrelativistic theory) involving this variable. Note that the Hamiltonian of the standard theory evolves quantum states in time, but does not act as a shift operator since it commutes with t . Dirac² has argued that if t were an operator, then the resulting t, E commutation relation would imply that the energy of the system

* e-mail:larry@post.tau.ac.il.

is unbounded below, from which he concluded that the time cannot be an observable in the nonrelativistic quantum theory (note, however, that in a relativistic theory, negative energies correspond to antiparticle states, and are not excluded). Moreover, as the axiomatic treatment of Piron⁴ (see also, Jauch⁵) shows, the Hilbert space of the quantum theory is constructed of a set of wave functions satisfying a normalization condition based on integration over all space, e.g., for a single particle, $\int |\psi_t(x)|^2 d^3x \leq \infty$, for each value of the parameter t . There is therefore a distinct Hilbert space for each value of the parameter t . A simple argument based on the propagator for the Schrödinger equation demonstrates that interference in time cannot occur in the standard Schrödinger theory.

The propagator for wave functions in the standard Schrödinger treatment, is given by

$$\begin{aligned} \langle x|U(t)|x' \rangle &= \left(\frac{m}{2\pi i t}\right)^{\frac{3}{2}} e^{i\frac{m}{2t}(x-x')^2} \\ &= G(x-x', t), \end{aligned} \tag{1}$$

where x is here a three dimensional variable. The action of the propagator is

$$\psi_t(x) = \int dx' G(x-x', t-t') \psi_{t'}(x'). \tag{2}$$

The integration over x makes possible the description of the double slit experiment in space (by coherently adding up contributions from two or more locations in x at a given t'); there is, however, no integration over t' , and therefore no mechanism for constructing interference in time. This result, obvious from the form of (2), is a reflection of the arguments of Ludwig³ cited above, and is fundamental to the standard nonrelativistic quantum theory. This structure constitutes a formal proof that no interference effect in time is predicted by the standard nonrelativistic Schrödinger theory. Introducing two packets into the beam of an experiment at two different times t_1 and t_2 would result in the direct sum of the two packets at some later time, say t_3 , if one propagates the first from t_1 to t_3 , and the second from t_2 to t_3 . This would constitute, by construction, a mixed state, for which no interference would take place, just as the construction of a beam of $n+m$ particles by adding a set of n particles with definite spin up to another set of m particles with spin down results in a mixed state described by a diagonal density matrix with *a priori* probability $n/(n+m)$ for outcome spin up, and $m/(n+m)$ for outcome spin down. There is no coherent superposition which would result in some spin with certainty in *any* direction⁵.

Moreover, as pointed out by Wick, Wightmann and Wigner⁶, a Hilbert space decomposes into incoherent sectors if there is no observable that connects these sectors; hence, if there were a larger Hilbert space containing a representation for t , the absence of any observable that connects different values of t in the standard nonrelativistic physics would induce a decomposition of the the Hilbert space into a (continuous) direct sum of superselection sectors⁴. Therefore, no superposition of vectors for different values of t would be admissible. This would exclude the interpretation of the experiment given in ref.1 forming the basis of the analysis carried out by the authors involving the linear superposition of two parts of a particle wave function arriving at the detector simultaneously, but originating at two different times, in the framework of the standard nonrelativistic quantum theory.

The significance of the experiment of ref. 1 is that it demonstrates at least one class of phenomena actually seen to occur in nature at low energies (but high frequency), for which the standard nonrelativistic quantum theory does not provide an adequate description, and this fact demands the development of some new theoretical tools which are a proper generalization of the standard theory.

The situation for particles, in this respect, is very different from that of electromagnetic waves, for which the second order equations imply coherence in time as well as space (the coherence time for light waves is a commonly measured characteristic of light sources). It is clear from the (spatial) double slit interference of light, which travels at a fixed velocity, that the sections of a wave front passing through the two slits must pass at different times if they are to arrive simultaneously at the detection plane off-center. The arrival of pieces of a *particle* wave packet which have passed through two spatially separated slits simultaneously on a screen off center is made possible by the dispersion of momenta in the wave packet, permitting a range of velocities. If the two contributions to the linear superposition on the screen were not taken to be simultaneous at the two slits, they would not interfere, since they would have originated on wave packets at different values of time. If, indeed, such interference could take place, we would have to add up the contributions passing each slit for all times, and this would destroy the interference pattern (one can see in the standard calculation in every textbook that the two pieces of the wave packet that contribute to the interference are taken at equal time, i.e., from a single wave packet arriving at the slits). Although this example is given in every quantum mechanics text as an essential property of the quantum theory, the possibility of interference in time is never mentioned.

Moshinsky⁷, in 1952, raised the question of interference in time. His calculation, however, was concerned with the evolution of a single wave packet, passed through a spatial slit opened at time $t = 0$. The transient form of the wave function was then calculated; it has the appearance of a Fresnel interference pattern. Using semiclassical time of flight arguments, it was deduced that this behavior could be thought of as an interference in time. The actual superposition of wave functions at two different times was not considered.

If we were to assume, arbitrarily, that the waves from sources at two different times could be coherently added (note that Eq.(1) is only valid for the evolution of physical states up to an undetermined c-number phase), the propagator above could be expanded in a power series for small variations in the final time, and one would find some semblance of an interference pattern for a few maxima before distortion would set in. For ε the time between peaks of the laser beam, T the time between peaks on the predicted “interference pattern” on a screen at a distance L from the emission source, m the mass of the electron, one finds a crude estimate

$$\varepsilon T \sim \frac{\pi \hbar}{2} \sqrt{\frac{m}{2}} L (E_{kin}^e)^{-\frac{3}{2}},$$

where E_{kin}^e is the kinetic energy of the emitted electron, and L is the distance from emitter to the detector. Salières, *et al*⁸, remark that in the very nonlinear emission of an electron by a high energy laser beam, several hundred photons may be absorbed to be converted to kinetic energy. If we take the electron kinetic energy as equal to the energy $\hbar\omega$ of the

laser beam, for this experiment, approximately $1.46eV$, with a factor of 300, and $L \sim 1cm$, we obtain for the first several predicted peaks (before distortion due to nonconvergence of the power series expansion) a value of $\epsilon T \sim 9 \times 10^{-28}sec^2$, or, for $\epsilon \cong T$, a diffraction spacing of $T \sim 3 \times 10^{-14} = 30fs$. This calculation indicates how an exact solution of the time dependent Schrödinger equation could exhibit an interference-like pattern for several peaks on the detector plane in agreement with the experimental results.. The numerical predictions of this integration do not, however, provide reliable evidence that the procedure is consistent with the theoretical basis of the standard nonrelativistic quantum theory.

There appear to be several types of theories which accommodate time as an observable. These are, in a nonrelativistic framework, Floquet theory⁹, for which the evolution operator has the form $E + H$, where H is a standard Hamiltonian model (often t dependent) and E corresponds to the operator $i\partial/\partial t$, a formulation of quantum theory in which one of the space variables becomes the evolution parameter and the time variable becomes an observable¹⁰, the quantum Lax-Phillips theory for irreversible processes (also applicable to relativistic quantum theory)¹¹, and in a relativistic framework, the so-called constraint theories¹² in addition to the theory developed by Stueckelberg¹³. In the book edited by J.G. Muga, *et al*¹⁴, much discussion is devoted to the difficulties of treating time as an operator in the standard framework, and some constructions are given for which certain operator valued functions of the phase space variables q, p can be interpreted as “time operators”. The spectra of these operators do not, however, enter into a propagator in a way that can naturally generate interference effects of the type observed in Lindner, *et al*.

A time operator, corresponding to the variable t that we recognize as time in the laboratory, is constructed in the work of Hahne¹⁰, in which he permits a space coordinate, say, z to act as an evolution parameter, and x, y, t become operator valued. Although coherence in t can be achieved, the use of z as an evolution parameter for the flow of the wave packet makes it difficult to formulate a description of the Lindner, *et al*, experiment in these terms.

The quantum Lax-Phillips theory¹¹ provides a systematic and rigorous description of irreversible processes. In this theory, a unitary evolution by a parameter, say s , is introduced on a Hilbert space \bar{H} , which is foliated along the spectrum t of a “time” variable which is a self-adjoint operator on \bar{H} into a set of Hilbert spaces H_t , which may be identified with the Hilbert spaces of the ordinary quantum theory, but maintains its coherence in t . The existence of invariant subspaces delimited by time intervals makes it possible to construct semigroups in these subspaces. The simplest model for evolution in the nonrelativistic case is of the form $E + H$, coinciding, as we shall see below, with the Floquet construction. Although there are applications of Lax-Phillips theory to the description of the experiment of Lindner, *et al*, the discussion of the Floquet construction below will be adequate for our present purposes.

The constraint theories¹², primarily directed toward a description of relativistic dynamics, are constructed from a set of functions K_i , where $i = 1, 2, \dots, N$ for a system of N particles; each of the form $p_i^\mu p_{i\mu} + m_i^2 + \phi_i$, where the interaction terms ϕ are functions of all the momenta and coordinates of all of the particles (here, $\mu = 0, 1, 2, 3$). In the classical form of the theory, the K_i 's are taken to be zero on a constraint hypersurface, and satisfy the requirement that they be first class constraints (vanishing Poisson brackets).

The evolution of the system goes with some parameter, say s , according to the transformations induced in the phase space by some linear combination of the K_i 's. For the two particle case, for ϕ of the form $\phi_1(x_1 - x_2) = \phi_2(x_2 - x_1)$, it is easy to see that this theory is equivalent to the Stueckelberg theory¹³ we shall discuss below (the relative motion for which $x = x_1 - x_2$ appears as a one-body problem). We therefore will not discuss the constraint theory further here. In following, I discuss the Floquet theory and the Stueckelberg theory as somewhat categorical among the set of theories which may be used to describe the Lindner, *et al*, experiment. Furthermore, since, as we shall point out below, the non-relativistic limit of the Stueckelberg theory goes over to the Floquet form, it appears as the main candidate for a viable description. Although the Stueckelberg theory is essentially relativistic, and the energies of the macroscopic motions of the particles involved are low, the very high frequencies used to establish excitations and pulse rates involve high energy components of the wave packets, and thus the use of a relativistically covariant theory is appropriate.

Floquet theory⁹ was originally intended for the treatment of differential equations with periodic coefficients. It entered physics in an important way in solid state theory where the potential in a crystal is periodic in space. Utilizing the translation operator $U(\mathbf{a}) = e^{i\mathbf{p}\cdot\mathbf{a}}$, where \mathbf{a} is a crystal lattice vector, one can show that the solutions of the Schrödinger equation, as a representation of this translation group, take on the Bloch form.

The idea then arose that for a Hamiltonian periodic in time, the same method could be used. However, since the Hamiltonian commutes with t , to make the group action explicit, it was necessary to introduce a new variable E (the generator of translations in t). The evolution operator was then defined as

$$K = E + H,$$

where $E \equiv -i\hbar\partial_t$. Then, clearly,for

$$U(T) = e^{-iKT}$$

the operator $U(T)$ carries $t \rightarrow t - T$, translating functions of t to the right by T .

The introduction of this modification of the Hamiltonian was also suggested by Howland¹⁵ for both the classical and quantum theories for treating problems in which the Hamiltonian depends on time. For the classical theory, introducing a new parameter of evolution, say s , the Hamilton equations would then include the relations

$$\frac{dt}{ds} = \frac{\partial K}{\partial E}$$

and

$$\frac{dE}{ds} = -\frac{\partial K}{\partial t} = -\frac{\partial H}{\partial t},$$

thus providing some interpretation for E . Since then $\frac{dt}{ds} = 1$, by a change of variables, this formulation becomes completely equivalent to the standard form. However, in the quantum theory, the Hamilton equations, as operator equations, imply conditions on expectation

values; the variables t and s are then no longer equivalent. In this case, s is the parameter of the motion, and t is a quantum operator, an observable. The wave functions are then coherent in t , making possible, in principle, interference phenomena in t .

The resulting theory is very different from the standard Schrödinger theory. To see this, let us write the corresponding evolution equation in what I shall call “Floquet theory”, since it has the same structure for the Hilbert space, but I will not insist that H be periodic in t . The mathematical framework is independent of this periodicity (clearly the consequences of the theory, and the results one may obtain, can be very strong when H is periodic in t).

The evolution equation has the form

$$\begin{aligned} i\frac{\partial\psi_s}{\partial s} &= K\psi_s \\ &= (-i\hbar\partial_t + H)\psi_s \end{aligned} \quad (3)$$

where ψ_s is a function of x, t . The functions ψ_s have the property that

$$\|\psi_s\|^2 = \int d^3x dt |\psi_s(x, t)|^2 < \infty, \quad (4)$$

the condition that ψ_s belongs to a Hilbert space H_s (now labelled by s). As pointed out by Kulander and Lewenstein¹⁶, if H (or K) is periodic over some hundreds of cycles, it would be a good approximation to assume a “stationary state” in which the s derivative of ψ_s is replaced by an eigenvalue (their equation (72.31)). Such a state would be stationary in s , not t ; the idea is that the spacetime function $\psi_s(x, t)$ reaches a steady form and no longer changes, on the spacetime manifold, as a function of s (up to a phase determined by the eigenvalue). In this case, the solution of eq.(3) amounts formally to an integration of the time dependent Schrödinger equation over t , as carried out in the analysis of ref. 1, with the Hamiltonian shifted by the Floquet eigenvalue (possibly zero). The theory would then predict coherence in t for such a solution. This is not, however, a valid procedure for the conditions of the experiment of ref. 1, since this experiment involves essentially just a couple of cycles.

In the following I calculate the propagator for the Floquet equation (3) for the case of a *free* particle. I will show that even though interference in t is, in principle, possible in the Floquet framework, two narrow segments, in time, of a particle wave function will not interfere unless (a) the segments initially overlap, or (b) there is a nontrivial t -dependence (but not necessarily periodic) in H , the latter certainly providing an interesting possibility for application to the experiment we are discussing.

To obtain the form of the propagator, let us consider the x, t matrix elements of the unitary evolution $U(s)$ of ψ_s :

$$\langle x, t|U(s)|x', t'\rangle = \int dE' dE dp' dp'' \langle x, t|E' p'\rangle \langle E' p'|e^{-i(H-E)s}|E'' p''\rangle \langle E'' p''|x' t'\rangle \quad (5)$$

Here, the momenta and coordinates are three dimensional (the differentials are also $dp \equiv d^3p$).

I now assume that H has the free particle form $p^2/2m$ and therefore commutes with E . Then, (5) becomes

$$\langle x, t | U(s) | x', t' \rangle = \delta(t' - t + s) \left(\frac{m}{2\pi i s} \right)^{\frac{3}{2}} e^{i \frac{m}{2s} (x-x')^2}. \quad (6)$$

Let us now call the coefficient of the δ function $G(x - x', s)$. The propagation of $\psi_{s'}(x', t')$ to $\psi_s(x, t)$ is given by

$$\psi_s(x, t) = \int dx' dt' \delta(t' - t + (s - s')) G(x - x', s - s') \psi_{s'}(x', t') \quad (7),$$

clearly displaying the possibility of interference in t , i.e., there may be contributions at several different values of t' corresponding to the opening of time gates. However, there is no spreading, in this propagation, of the width of the time pulses, independently of the form of the wave packet $\psi_{s'}(x', t')$.

Consider the contribution of two gates at t_1 and t'_1 . In this case,

$$\begin{aligned} \psi_s(x, t) = & \int dx' G(x - x', s - s') \\ & \{ \delta((s - s') - (t - t_1)) \psi_{s'}(x', t_1) \Delta t_1 + \delta((s - s') - (t - t'_1)) \psi_{s'}(x', t'_1) \Delta t'_1 \}, \end{aligned} \quad (8)$$

where Δt_1 and $\Delta t'_1$ are the (narrow) widths of the gates.

One can evaluate $s - s'$ approximately through the Hamilton equations. Since $dx/ds = p/M$, it follows that $\Delta s \cong ML/p$, where L is the distance from source to detector. Since $d < t > / ds = 1$, the expectation value of $t(s)$ goes with s , so that $s - s' \cong t - t'$, the latter giving the time from the source gate to the time on the detector when the measurement is made. Due to the delta function constraint, we see that there can be no interference if the source pulses do not overlap. Alternatively, the delta functions would not appear if the Hamiltonian had an explicit t -dependence, and the result would depend on the particular model. Interference in the framework of the Floquet structure, therefore, although in principle possible, would not occur for narrow source pulses in the absence of explicit time dependence in the Hamiltonian.

The Floquet theory is, in fact, a nonrelativistic limit of Stueckelberg's relativistic quantum theory¹³, the second way of introducing coherence in time which I wish to discuss here. In 1976, Horwitz and Rabin¹⁷ pointed out that the relativistic quantum theory of Stueckelberg predicts interference in time. In this theory, t is treated as a quantum observable, since the Einstein variables x, t are considered, in relativity, as the nontrivial outcome of experiments measuring the place and time of occurrence of events. Their calculation will be briefly redone below for the parameters of the experiment of ref.1. In this theory, interference does not require initial overlap or an explicitly time dependent Hamiltonian. The estimate given below shows that the interference criteria are satisfied with numbers very close to the conditions and results of the experiment under discussion; the high frequencies required are due in this case to the large value of the velocity of light.

The Stueckelberg theory for the free particle introduces an equation quite similar to that of the Floquet equation, but with an evolution operator that is Lorentz invariant:

$$i\hbar \frac{\partial \psi_s}{\partial s} = \frac{p^2 - (\frac{E}{c})^2}{2M} \psi_s, \quad (9)$$

where $\psi_s(x, t)$ satisfies the same normalization condition as for the Floquet theory, on space and time, i.e., $\int |\psi_s(x, t)|^2 d^4x \leq \infty$, and M is the Galilean target mass (the so-called mass shell value for $m^2c^2 = (E/c)^2 - \mathbf{p}^2$). The propagator has a similar form to that of the Floquet propagator, but is Gaussian in all four variables:

$$\langle x|U(s)|x' \rangle = \left(\frac{M}{2\pi i s \hbar}\right)^2 e^{i\frac{M}{2s\hbar}(x-x')^2}, \quad (10)$$

where now $(x - x')^2$ is the invariant $(\mathbf{x} - \mathbf{x}')^2 - c^2(t - t')^2$; we write x for (x, t) . It is the quadratic term in $t - t'$ in the exponent which leads directly to interference in the same way as the double slit in space.

The diffraction formula, obtained from (10), using the Hamilton relations

$$\frac{dx}{ds} = \frac{p}{M}, \quad (11)$$

and (note that this relation allows for two pulses emitted at different times to arrive at a detector at the same time due to the spread in the spectrum of E)

$$\frac{dt}{ds} = \frac{E}{Mc^2}. \quad (12)$$

is¹⁷

$$\varepsilon T \cong \frac{2\pi\hbar L}{\langle p \rangle c^2}, \quad (13)$$

where ε is the gate spacing in time, and T is the time between diffraction peaks at a distance L .

For 850 nm light, as utilized in the experiment under discussion, as remarked above, $\hbar\omega$ is about $1.46eV$. Using the on-shell value for the electron mass, taking into account (as assumed above) that the electron may absorb about 300 photons during the emission, cp (for p in the beam direction) then has a value of $1.21 \times 10^3 eV$. With these values, one finds that ,

$$\varepsilon T \cong 6.9 \times 10^{-30} sec^2, \quad (14)$$

so that for $\varepsilon \sim T$, $T \sim 2.6 \times 10^{-15} sec$. This result, for the pulse rate and the observed diffraction pattern, is in good agreement with the results obtained in the experiment.

More precise estimates can be obtained by taking into account more details of the interaction, and the dependence on L can be used as a parameter to test the reliability of (13).

The relativistic model therefore seems to provide a simple and consistent description of the experimental results. At very low energies, the Stueckelberg theory, which carries a clear interpretation of t as an observable, reduces approximately to the Floquet form¹⁸. The estimate made above therefore includes the result that would be obtained approaching the low energy Floquet type as a limiting case.

I thank C. Piron for communications on the question of coherence, and Gerhard Paulus, one of the authors of ref. 1, for discussions of this experiment and the analyses

applied. I would also like to thank Igal Aharonovich for bringing the initial announcement of this experiment to my attention and Noam Erez and Jonathan Silman for useful discussions.

References

1. F. Lindner, M.G. Schätzel, H. Walther, A. Baltuska, E. Goulielmakis, F. Krausz, D.B. Milošević, D. Bauer, W. Becker and G.G. Paulus, quant-ph/0503165, Phys. Rev. Lett. **95**, 040401 (2005). See also, Pascal Szriftgiser, David Guéry-Odelin, Markus Arndt and Jean Dalibard, Phys. Rev. Lett. **77** 0031-9007 (1996); M. Wollenhaupt, A Assion, D. Liese, Ch. Sarpe-Tudoran, T. Baumert, S. Zamith, M.A. Bouchene, B. Girard, A Flettner, U. Weichmann and G. Gerber, Phys. Rev. Lett. **89**, 0031-9007 (2002).
2. P.A.M. Dirac, *Quantum Mechanics*, First edition, pp.34,36, Oxford Univ. Press, London (1930); *Lectures on Quantum Field Theory*, Academic Press, New York (1966). See also W. Pauli, *General Principles of Quantum Mechanics*, p.63, Springer-Verlag, Heidelberg (1980); E. Schrödinger Berl. Ber., p. 238 (1931); P. Carruthers and M.M. Nieto, Rev. Mod. Phys. **40**, 411 (1968), and references therein.
3. G. Ludwig, *Foundations of Quantum Mechanics I*, p.295, Springer-Verlag, New York (1983).
4. C.P. Piron, *Foundations of Quantum Physics*, W.A. Benjamin, Reading (1976). See also, C. Piron, physics/0204083 29 Apr. 2002; *Trends in Quantum Mechanics*, eds. H.-D. Doebner *et al*, p.270, World Scientific, Singapore (2000).
5. J.M. Jauch, *Foundations of Quantum Mechanics*, Addison-Wesley, Reading (1968).
6. G.C. Wick, A.S. Wightman and E. Wigner, Phys. Rev. **88**, 101 (1952). See also C. Piron, Helv. Phys. Acta **42**, 330 (1969).
7. M. Moshinsky, Phys. Rev. **88** 625 (1952).
8. P. Salières, B. Carrée, L. Le D’eroff, F. Grasbon, G.G. Paulus, H. Walther, R. Kopold, W. Becker, D.B. Milosevic, A. Sanpera, M. Lewenstein, Science **292**, 902 (2001).
9. For example, H.L. Cycon, R.G. Froese, W. Kirsch and B. Simon, *Schrödinger Operators with Application to Quantum Mechanics and Global Symmetry*, p. 146, Springer-Verlag, New York (1987); see also, R.M. Potvliege and R. Shakeshaft, *Atoms in Intense Laser Fields*, ed. M. Gavrilla, Academic Press, San Diego (1992); S.I. Chu, Adv. Chem. Phys. **73**, 739 (1989). Note that the definition of $H - i\hbar\partial_t$ as a self-adjoint operator is sufficient to define the structure of the Hilbert space and imply Eq. (5).
10. G.E. Hahne, Jour. Phys. A **367**, 144 (2003)
11. Y. Strauss, L.P. Horwitz and E. Eisenberg, Jour. Math. Phys. **41** 8050 (2000), Fermilab-Pub-97/227-T. The original form of the theory, intended for application to classical wave equations such as for optical or acoustic waves, is worked out in the book *Scattering Theory*, P.D. Lax and R.S. Phillips, Academic Press, New York (1967). See also, L.P. Horwitz and C. Piron, Helv. Phys. Acta **66**, 693 (1993).
12. F. Rohrlich, Phys. Rev. D **23**, 1305 (1980), and references therein to earlier work.
13. E.C.G. Stueckelberg, Hel. Phys. Acta **14**, 372, 585 (1941); **15**, 23 (1942); see also, L.P. Horwitz and C. Piron, Helv. Phys. Acta **46**, 316 (1973). *Proceedings of the Workshop on Constraints Theory and Relativistic Dynamics*, Arcetri, Firenze, Italy, 1986, ed. G. Longhi and L. Lusanna, World Scientific, Singapore (1987); L.P. Horwitz and

- F. Rohrlich, Phys. Rev. D**24**, 1528 (1984); K. Sundermeyer, *Constraint Dynamics*, Lecture Notes in Physics, vol.169, Springer, Berlin (1982).
14. *Time in Quantum Mechanics*, ed. J.Muga, R. Sala Mayato and I.L. Egusquiza, Lecture Notes in Physics, Springer, Berlin (2002).
 15. J.S. Howland, Indiana Math. Jour. **28**, 471 (1979).
 16. K.C. Kulander and M. Lewenstein, *Atomic, Molecular and Optical Physics Handbook* (G.W. Drake, ed.), p. 828, American Institute of Physics Press, Woodbury, N.Y. (1996).
 17. L.P. Horwitz and Y. Rabin, Lett. Nuovo Cimento **17**, 501 (1976).
 18. L.P. Horwitz, W.C. Schieve and C. Piron, Ann. Phys. **137**, 306(1981); L.P. Horwitz and F. Rotbart, Phys. Rev. D **24**, 2127 (1981).