

A nother Look at Quantum Teleportation

L. Vaidman,¹ N. Erez,^{1,2} and A. Retzker¹

¹ School of Physics and Astronomy
Raymond and Beverly Sackler Faculty of Exact Sciences
Tel-Aviv University, Tel-Aviv 69978, Israel

² Institute for Quantum Studies and Physics Dept.,
Texas A & M University, College Station, TX 77843-4242

A dialog with Asher Peres regarding the meaning of quantum teleportation is briefly reviewed. The Braunstein-Kimble method for teleportation of light is analyzed in the language of quantum wave functions. A pictorial example of continuous variable teleportation is presented using computer simulation.

I. INTRODUCTION

I, Lev Vaidman, knew Asher Peres since the beginning of my interest in the Foundations of Quantum Mechanics. Mostly, we were fighting to prove that the interpretations of quantum effects adopted by each of us were better. Asher was against the usage by Aharonov and myself of a quantum state evolving backwards in time [1, 2, 3, 4] and contesting our interpretation of the physical meaning of "weak measurements" [5, 6, 7]. He objected to the names I gave to my proposals like "cryptography with orthogonal states" [8, 9, 10] or "interaction-free measurements" [11]. Our disagreements did not make our interactions less fruitful: we agreed about physical facts and discussion of the interpretation only sharpened our (at least mine) understanding of various aspects of these effects.

Even the most basic disagreement, where Asher says: "Quantum mechanics needs no interpretation" [12], and I write that the many-worlds interpretation is by far the best way to view quantum mechanics [13], is also essentially a disagreement only about names. Asher explains his view describing Katchy, an experimental physicist who, after making a quantum experiment becomes a superposition of a lady who ate a cake and a lady who ate a fruit. Asher says that even at this stage, in principle, it is still possible to reverse the evolution and come back to the state which was before the quantum measurement. For me, this story is a gedanken test of the many-worlds interpretation. We completely agree on the facts: There is no such thing as the collapse of the quantum wave function. I interpret this story as splitting and reunion (with the help of super-technology) of two worlds, while Asher, to avoid paradoxes, considers this as an argument in favor of the approach according to which quantum mechanics should not be taken as a description of an objective reality.

The main part of this paper is devoted to the renewed analysis, performed together with two members of the quantum group of Tel-Aviv University, of the topic on which I and Asher, in a way, collaborated: this is the issue of teleportation. I was very pleased to hear from Asher that finally we came to an agreement. In the last e-mail I received from him two weeks before he left us, he recommended to the Jerusalem Report to interview me instead of him (due to his health condition) about teleportation. In his paper "What is actually teleported?" [14], Asher is joking about the suggestion of Charlie Bennett to cite the "weak measurements" of Aharonov and myself, but he mentions that in another work [15] there are seeds of the teleportation paper [16].

Indeed, I had the tools to find the solution for the teleportation problem. When I saw

the abstract of this seminal paper I immediately knew how to do it (in my way). But it took the genius of Asher and his collaborators to ask the question. Still, my way of teleportation was useful too. I proposed a method for two-way teleportation which is applicable also for continuous variables [17]. The importance of this work became clear only a few years later, when Braunstein and Kimble found a realistic way to implement the continuous variables teleportation experiment using squeezed light [18]. This experiment was successfully accomplished in 1998 [19] and recently improved [20].

Braunstein and Kimble [18] described their proposal in the language of Wigner functions, the common approach of the quantum optics community. Numerous analysis and further experiments since then mostly continued to use the Wigner function formalism. I believe that the language of quantum states has advantages in discussions of the Foundations of Quantum mechanics, so it is of interest to present the Braunstein-Kimble experiment in the language of quantum states. Section 3 is devoted to this purpose. In Section 4 the results of Section 4 are demonstrated on a particular example. But before this, I cannot resist the temptation to continue the interpretation dialogue with Asher.

II. WHAT IS ACTUALLY TELEPORTED?

In the framework of classical physics, teleportation, defined as e.g. "theoretical transportation of matter through space by converting it into energy and then reconverting it at the terminal point," the quotation from Asher's dictionary (Webster), is obviously a science fiction concept. Massive objects, "matter" cannot "jump" from one place to another. The definition in The Oxford Dictionary

teleportation. Psychics and Science Fiction. The conveyance of persons (esp. of oneself) or things by psychic power; also in futuristic description, apparently instantaneous transportation of persons, etc., across space by advanced technological means.

sounds equally impossible for implementation. However, quantum theory makes it more plausible. According to quantum theory, all elementary particles of the same kind are identical. There is no difference between the electrons in my body and the electrons in a rock on the moon. Thus, what defines a particular person is not a collection of the elementary particles he is made of, but the quantum state of these particles. If I want to move to the moon, I need not move my electrons, protons, etc. to the moon. It is enough to reconstruct the quantum state of the same particles there. From my point of view, I am the quantum state, so creation of this quantum state on the moon is my teleportation to the moon. Compare this view with Asher's reply when he was asked by a newsmen, whether it was possible to teleport not only the body but also the soul: "only the soul."

In this approach, teleportation sounds as trivial as a FAX machine, but it is not. There are two reasons why it seems impossible. First, it is impossible to measure (to scan) the quantum state. Second, the amount of information needed to specify a quantum state even of a small object is so huge that it is not feasible to transmit it in a reasonable time. The dual channel of quantum teleportation does the trick: the quantum state is teleported without being scanned. The quantum channel consists of entangled pairs of elementary particles, as many as we need for the object to be teleported. Originally, at the remote location there is a mixture of different states in which probability for any state is the same. Then,




FIG. 1: The Braunstein-Kimble Teleportation scheme

local joint measurement performed on the system to be teleported together with the local part of the quantum channel specifies the particular decomposition of the mixture in the remote location with relatively small number of states. Finally, the only information to be transmitted is the number of the "actual" state in the mixture. In my view, [21], the local measurement creates numerous worlds with the teleported quantum state which is deformed in various ways. The final stage of teleportation is the correction of the deformation such that in all worlds the final state of the remote system is the initial state of the local system.

Clearly, Asher would not join me considering myself as an (unknown) quantum state. For him, a quantum state is just the knowledge of the preparer [22]:

A state vector is not a property of a physical system (nor of an ensemble of systems). ... Rather, a state vector represents a procedure for preparing or testing one or more physical systems.

Then, the correction is really not that important: the preparer knows that I am teleported in a particular deformed way. The "deformed me" is, probably, not a living creature at all, so I tend not to accept Asher's approach. But, since we are both sure that now and in any foreseeable future, a realistic teleportation experiment with people is a science fiction story, this disagreement is irrelevant.

III. WAVE FUNCTION DESCRIPTION OF THE BRAUNSTEIN-KIMBLE SCHEME

In their seminal paper on the implementation of continuous teleportation with squeezed light [18], Braunstein and Kimble had used the Wigner representation. In this section, we explain their method using wave functions.

Figure 1 is a schematic representation of the experimental setup envisioned by Braunstein and Kimble. The (single mode) state of the beam incident on the "input" is to be teleported




FIG. 2: Beam Splitter

to the "out port". To this end, a highly squeezed two-mode state is used (which leaves the source marked "EPR"). One half of the "EPR-pair" is combined via a 50-50 beam splitter with the "in" beam and the two resulting beams are measured using homodyne detectors D_x and D_p measuring x and p appropriately. The results of these measurements are then used to implement corrections on the other (remote) half of the EPR pair which leave it in a state which closely approximates the input state.

For simplicity, we will consider the (asymmetric) 50-50 beam splitters (see Fig 2) which act on single photons in the following way:

$$j_i \mapsto \frac{\beta_i + j_i}{2}; \quad p_i \mapsto \frac{j_i - \beta_i}{2}. \quad (1)$$

If the incident beams in ports 1 and 2 are described by the quadrature-wave function $(x_1; x_2)$, then the beam splitter described by (1) leads to the transformation [23]:

$$(x_1; x_2) \mapsto \left(\frac{x_4 + x_3}{2}; \frac{x_4 - x_3}{2} \right). \quad (2)$$

We start with the initial state (x_1) for the quadrature-wave function of the input beam, $(x_2; x_5)$ for that of the (approximate) EPR-pair (the quantum channel) and initial "ready" states of two measuring devices:

$$j_i = \int_{-\infty}^{\infty} (x_1) j_{x_1} dx_1 \int_{-\infty}^{\infty} (x_2; x_5) j_{x_2} j_{x_5} dx_2 dx_5 j_{\text{ready}_1} j_{\text{ready}_2}. \quad (3)$$

Using Eq. (2), we have for the total state after the action of the beam splitter:

$$\int_{-\infty}^{\infty} \left(\frac{x_4 + x_3}{2}; \frac{x_4 - x_3}{2}; x_5 \right) j_{x_3} j_{x_4} j_{x_5} dx_3 dx_4 dx_5 j_{\text{ready}_1} j_{\text{ready}_2}. \quad (4)$$

At this stage, x_3 and p_4 are measured and the appropriate correction is applied to the state of the variable x_5 . The wave function in the x representation is shifted by $\frac{1}{2}x_3$, and the wave function in the p representation is shifted by $\frac{1}{2}p_4$. At this stage of our analysis we will not introduce the "collapse" of the quantum measurement, but continue to include the measuring device in the description of the total state. The shift in x leads to the following transformation:

$$\int_{-\infty}^{\infty} \left(\frac{x_4 + x_3}{2}; \frac{x_4 - x_3}{2}; x_5 + \frac{2x_3}{2} \right) j_{x_3} j_{x_4} j_{x_5} dx_3 dx_4 dx_5 j_{\text{ready}_1} j_{\text{ready}_2}. \quad (5)$$

This is just a convolution of the input function with a (real) Gaussian multiplied by an (imaginary) exponent. If the Gaussian is narrow and we can neglect the distortion due to the exponent, the convolution yields approximately the input wave function, i.e., we obtain teleportation with good fidelity.

The distortion due to the exponent depends on the value of p_4 . In order to estimate it, we can write an operator equation similar to those that appear in the original continuous teleportation paper [17]:

$$p_4 = \frac{p_1}{2} + \frac{p_a + p_b}{2}; \quad (15)$$

where p_1 is the momentum of the input mode and p_a and p_b are the momenta of the input modes of the EPR source. For our input state we have:

$$\langle p_a \rangle = 0; \quad p_a = 1/a; \quad p_b = 0; \quad (16)$$

Taking into account the rest of the requirements (10), we see that for probable outcomes of the measurement of p_4 , $|p_4| < \frac{1}{a}$. Since we consider a narrow Gaussian such that $\psi(x)$ is nearly constant on an interval of length a , the exponent does not lead to a significant distortion and we obtain:

$$\psi_{\text{tel}}(x_5) \approx \psi(x_5); \quad (17)$$

In the other case we had mentioned above, namely that only the squeezing in mode b is not ideal, the EPR state is given by:

$$\psi(x_2; x_5) = \psi(x_2, x_5) \frac{e^{-\frac{x_2 + x_5}{2b}}}{\sqrt{1 + \frac{1}{4b^2}}}; \quad (18)$$

The final state after the teleportation procedure is now (up to normalization):

$$\int e^{ix_3 p_4} e^{-\frac{x_5}{2b} p_4^2} \psi(x_5) \psi(x_3) \psi_{\text{EPR}}(p_4) \psi_{\text{EPR}}(x_5) dx_3 dp_4 dx_5; \quad (19)$$

Again, there is a partial entanglement between the system with the teleported state and the measuring devices. Let us look at a particular outcome of the measurement in mode 3, x_3 . This eliminates the entanglement. Then, the final teleported state (up to normalization) is:

$$\psi_{\text{tel}}(x_5) = \psi(x_5) e^{-\frac{x_5}{2b} p_4^2}; \quad (20)$$

For the distortion to be small, we need the Gaussian to be approximately constant over the interval where $|p_4|$ is significant. Let us denote the length of this interval by l which, according to our second choice in (10), is much smaller than b . The condition is then:

$$x_3 \approx \frac{1}{b}; \quad (21)$$

To estimate the range of the outcomes of measurement results of x_3 , we can write (compare with (15)):

$$x_3 = \frac{x_1}{2} + \frac{x_a + x_b}{2}; \quad (22)$$

In this case, the input state of the EPR source is characterized by

$$x_a = 0; \quad \langle x_b \rangle = 0; \quad x_b = b; \quad (23)$$

which together with (10), ensures that condition for good fidelity teleportation (21) is satisfied.

We have seen two cases where simple estimation of teleportation distortion was possible: in one case the operation was essentially convolution with a narrow Gaussian, and in the other case the operation was essentially multiplication by a wide Gaussian. Note, that we could consider the process of teleportation in the p representation and then we would have the same explanations, but multiplication in the first case and convolution in the second.

We have shown that either of the choices

$$a = 1; \quad b = 1 \quad \text{or} \quad a = 0; \quad b = 1 \quad (24)$$

ensures that the distortion shall be small. In a real experiment, both input beams for the EPR source exhibit, of course, finite squeezing (9). Then, we can see from (6) that the final state of the teleportation procedure (up to normalization) is:

$$\int_{-\infty}^{\infty} e^{ix_3 p_4} e^{-\left(\frac{x_5 - v}{2a}\right)^2} e^{-\frac{x_5 + v}{2b} \frac{p_4 x_3}{2}} e^{-\frac{p_4^2}{2}(v - x_5)p_4} (v) \langle x_3 \rangle \langle x_3 \rangle_{MD1} \langle p_4 \rangle \langle p_4 \rangle_{MD2} \langle x_5 \rangle dx_3 dp_4 dx_5 dv \quad (25)$$

Given particular outcomes of the measurements of p_4 and x_3 , the final teleported state (25) (up to normalization) is:

$$\rho_{tel}(x_5) = \int_{-\infty}^{\infty} e^{-\left(\frac{x_5 - v}{2a}\right)^2} e^{-\frac{x_5 + v}{2b} \frac{p_4 x_3}{2}} e^{-\frac{p_4^2}{2}(v - x_5)p_4} (v) dv; \quad (26)$$

The analysis of this expression is more complicated, but condition (10) is sufficient to ensure that the function will be teleported with little distortion, as will be seen in the computer simulation presented in the following section.

IV. A NUMERICAL SIMULATION

In this section we show numerically the impact of each parameter in the teleportation. We use as our input wave function the silhouette of a woman Fig.(3). We would like to stress that the choice of a human figure is intended to aid in the visual assessment of the distortion and is not, of course, supposed to suggest that teleportation of humans is feasible.

The characteristics of the spatial wave function are:

$$\langle x_1 \rangle = 50; \quad x_1 = 28; \quad l = 100; \quad (27)$$

where we have chosen units such that $h = 1$. Then, the computer calculations show that the characteristics of the wave function in momentum space are:


$$\langle p_1 \rangle = 0; \quad p_1 = 18; \quad (28)$$

First, we consider the case of infinite squeezing in port b and finite squeezing in port a. In Fig. 4a we show the results of the simulation for strong squeezing, $a = 1/180$ ($= 1/10 p_1$)

Silhouette



The Wave Function

FIG. 3: The original wave function $\psi(x_1)$.

and a probable outcome of the measurement of the momentum in mode 4, $p_4 = 180 (= 1\sigma_a)$. Fig. 4b shows the results for the same strong squeezing, but a rare large outcome of the measurement of the momentum, $p_4 = 1800 (= 10\sigma_a)$. Fig. 4c shows the results for weak squeezing, $\sigma_a = 1/5.4 (= 1/0.3 p_1)$ and probable outcome of the measurement of the momentum $p_4 = 2.7 (= 0.5\sigma_a)$. Finally, Fig. 4d shows the results for weak squeezing, $\sigma_a = 1/5.4 (= 1/0.3 p_1)$ and rare large outcome of the measurement of the momentum, $p_4 = 21.6 (= 2\sigma_a)$.

We see in Fig. 4a that with strong squeezing and not too large a value of p_4 , the wave function is teleported without significant distortion. Fig. 4b shows that for strong squeezing, but an improbable large value of p_4 leads to distortion of the regions which require large momenta. Fig. 4c shows that weak squeezing causes smoothing out of small details of the




FIG. 4: Teleported silhouette with finite squeezing in a and ideal squeezing in b




FIG . 5: The function $\psi(v)$ and (the real part of the) convolution kernel of Eq. (29) ($x_5 = 30$) corresponding to Fig. 4c. On the left is the scale of the wave function and on the right, that of the kernel.

wave function and large p_4 and weak squeezing lead to complete distortion of the teleported wave function.

Let us analyze, for example, something out of all details of the wave function which we have seen in Fig. 4c. The teleported wave function is given by the convolution (14) which, with the parameters $l = a = 5.4$ $p_4 = 2.7$, reads:

$$\psi_{\text{tel}}(x_5) = \int_{-\infty}^{\infty} e^{-i3.8(v - x_5)} e^{-\left(\frac{x_5 - v}{0.137}\right)^2} \psi(v) dv \quad (29)$$

In Fig. 5 we draw the terms of the convolution (29). The graph shows only the real part. The imaginary part is an order of magnitude smaller than the real part. The multiplication of the Gaussian by the exponent has small effect, since the width of the Gaussian is smaller than most details of the wave function $\psi(v)$. Thus, the result of the convolution is close to the original wave function, yet some details do disappear, like the mouth.

The second case is in finite squeezing in port a and finite squeezing in port b. In Fig.




FIG . 6: Teleported silhouette with finite squeezing in b and ideal squeezing in a




FIG. 7: The function $\psi(v)$ and the Gaussian which are the terms of the product (30) corresponding to Fig. 6a. On the left is the scale of the wave function and on the right, that of the Gaussian.

6a we show the results of the simulation for strong squeezing, $b = 280 (= 10 \times)$ and a probable outcome of the measurement of position, $x_3 = 280 (= b)$. Fig 6b shows the results for the same good squeezing, but rare large outcome of the measurement of position, $x_3 = 2800 (= 10 b)$. Fig. 6c shows the results for weak squeezing, $b = 8.4 (= 0.3 \times)$ and probable outcome of the measurement of position $x_3 = 4.2 (= 0.5 b)$. Fig. 6d shows the results for weak squeezing, $b = 8.4$ and rare large outcome of the measurement of the momentum, $x_3 = 33 (= 4 b)$.

We see in Fig. 6a that with strong squeezing and not too large a value of x_3 , the wave function is teleported without significant distortion. It is the product (20) of the original function with a wide Gaussian, which for the above parameters reads:

$$\psi_{\text{tel}}(x_5) = \psi(x_5) e^{-\left(\frac{x_5 + 400}{280}\right)^2} : \quad (30)$$

Fig. 7 shows the wave function and the Gaussian of the product (30). We can see that the Gaussian is nearly constant over the support of the wave function, thus causing little distortion.

Fig. 6b shows that strong squeezing, but improbable large value of x_3 lead to distortion of the relative amplitude. Fig. 6c shows that weak squeezing and small x_3 yield relatively faithful teleportation of a small part of the wave function and distortion (elimination) of




FIG. 8: Teleported silhouette with finite squeezing in a and b

other parts. Again, weak squeezing together with large x_3 lead to complete distortion of the teleported wave function.

Finally, we compute the teleported wave function when both input states are not ideal. Fig. 8b shows that strong squeezing $r_b = 280 (= 10 \times)$; $r_a = 180 (= 10 \times)$ and probable outcomes $p_4 = 180 (= 10 \times)$; $x_3 = 280 (= 10 \times)$ together lead to a very good fidelity, while large improbable outcomes $p_4 = 1800 (= 10 \times)$; $x_3 = 2800 (= 10 \times)$ lead to significant distortion, see Fig. 8c.

The discussion of the physical meaning of teleportation and the analysis of continuous variables teleportation experiment which we performed here provides an intuitive picture of the process and helps to understand the effect of various parameters on the fidelity of teleportation. We hope it will be useful for designing better teleportation experiments.

It is a pleasure to thank Shai Machnes for helpful discussions. This research was supported in part by grant 62/01 of the Israel Science Foundation. N.E. gratefully acknowledges support from the Robert A. Welch Foundation, grant no. A-1218.

-
- [1] Y. Aharonov and L. Vaidman, Phys. Rev. A 41, 11 (1990).
 - [2] A. Peres, Phys. Lett. A 194, 21 (1994).
 - [3] Y. Aharonov and L. Vaidman, Phys. Lett. A 203, 148 (1995).
 - [4] A. Peres, Phys. Lett. A, 203, 150 (1995).
 - [5] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. 60, 1351 (1988).
 - [6] A. Peres, Phys. Rev. Lett. 62, 2326 (1989).
 - [7] Y. Aharonov and L. Vaidman, Phys. Rev. Lett. 62, 2327 (1989).
 - [8] L. Goldenberg and L. Vaidman, Phys. Rev. Lett. 75, 1239 (1995).
 - [9] A. Peres, Phys. Rev. Lett. 77, 3264 (1996).
 - [10] L. Goldenberg and L. Vaidman, Phys. Rev. Lett. 77, 3265 (1996).
 - [11] Asker told me that the name should be "energy and momentum exchange-free measurements".
 - [12] C. A. Fuchs and A. Peres, Phys. Today, 53, 70 (2000).
 - [13] "The Many-Worlds Interpretation of Quantum Mechanics", L. Vaidman, in The Stanford Encyclopedia of Philosophy (Summer 2002 Edition), Edward N. Zalta (ed.), URL = <http://plato.stanford.edu/entries/qm-manyworlds/>.
 - [14] A. Peres, IBM J. Res. Dev. 48, 63 (2004).
 - [15] L. Vaidman, Y. Aharonov and D. Albert, Phys. Rev. Lett. 58, 1385 (1987).
 - [16] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. 70, 1895 (1993).
 - [17] L. Vaidman, Phys. Rev. A 49, 1473 (1994).
 - [18] S. L. Braunstein and H. J. Kimble, Phys. Rev. Lett. 80, 869 (1998).
 - [19] A. Furusawa, J. L. Sørensen, S. L. Braunstein, C. A. Fuchs, H. J. Kimble, and E. S. Polzik, Science, 282, no. 5389, 706 (1998).
 - [20] N. Takei, H. Yonezawa, T. Aoki, and A. Furusawa, Phys. Rev. Lett. 94, 220502 (2005).
 - [21] "On the Paradoxical Aspects of New Quantum Experiments", L. Vaidman, in Philosophy of Science Association 1994, pp. 211-217 (1994).
 - [22] A. Peres, Am. J. Phys. 52, 644 (1984).
 - [23] U. Leonhardt, Measuring the Quantum State of Light (Cambridge university press, 1997).