

Microscopic Theory of Spontaneous Decay in a Dielectric

Hao Fu, P. R. Berman

Michigan Center for Theoretical Physics,

FOCUS Center, and Physics Department,

University of Michigan, Ann Arbor, Michigan 48109-1040

(Dated: today; Received textdate; Revised textdate; Accepted textdate)

Abstract

The local field correction to the spontaneous decay rate of an impurity source atom embedded in a disordered dielectric is calculated to second order in the dielectric density. The result is found to differ from predictions associated with both "virtual" and "real" cavity models of this decay process. However, if the contributions from two dielectric atoms at the same position are included, the virtual cavity result is reproduced.

I. INTRODUCTION

The problem of spontaneous emission from an atom imbedded inside a dielectric has attracted considerable interest [1]. Most theoretical treatments of this problem follow a macroscopic approach [2]. Based on different models of the local environment of the imbedded atom, they give different types of local field corrections to the spontaneous decay rate Γ_0 of the impurity atom. The so-called virtual cavity model gives a decay rate, $\Gamma_{\text{virtual}} = \Gamma_0 \left(\frac{\epsilon - 2}{\epsilon + 2} \right)^2$, assuming that a virtual cavity surrounds the emitter, while the "real" cavity model gives a rate, $\Gamma_{\text{real}} = \Gamma_0 \left(\frac{\epsilon - 3}{\epsilon + 1} \right)^2$, assuming that an empty spherical cavity surrounds the emitter. The quantity ϵ is the permittivity of the dielectric, which is connected to the microscopic polarizability α by the Lorentz-Lorenz relation $\epsilon = 1 + \frac{4\pi N \alpha}{1 - \frac{1}{3} N \alpha}$, where N is the dielectric density. Expansions for the decay rates in powers of $N \alpha$ yield $\Gamma_{\text{real}} = \Gamma_0 \left[1 + \frac{7}{6} N \alpha + \frac{19}{72} (N \alpha)^2 + O(N \alpha)^3 \right]$ and $\Gamma_{\text{virtual}} = \Gamma_0 \left[1 + \frac{7}{6} N \alpha + \frac{17}{24} (N \alpha)^2 + O(N \alpha)^3 \right]$. To first order in $N \alpha$, the real and virtual cavity models give identical results, but they differ in higher order. To determine the validity range of these macroscopic models, calculations using a somewhat more fundamental approach are needed. Several attempts at such microscopic models involve (i) a polariton approach for crystals [3], (ii) a Green's function approach for crystals [4] and disordered dielectrics [5], and (iii) an amplitude approach for disordered dielectrics [6][7]. In the polariton method, the interaction between the vacuum radiation field and the crystal atom is solved exactly; the eigenmodes of this system are the polaritons. The source atom then decays by radiating polaritons. This polariton calculation agrees with the virtual cavity result [3]. In the Green's function approach, the modification of the decay rate results from scattering of radiation emitted from the source atom by the dielectric, calculated to all orders in the dielectric density. This calculation reproduces the virtual cavity result with the source atom at an interstitial position and the real cavity result with source atom at a substitutional position in the crystal [4]. For disordered dielectrics, the Green's function method gives the virtual cavity result [5]. The amplitude method represents a direct calculation of the modification of the decay rate as a perturbation series in $N \alpha$ [6][7]. To first order in $N \alpha$, the radiation emitted by the source atom is scattered back to the source atom by a single dielectric atom; the resultant decay rate agrees with the both virtual and real cavity models to first order in $N \alpha$ [6].

In this paper, the amplitude method is extended to second order by including scattering

events in which the radiation emitted by the source atom is scattered back to the source atom by a combined scattering from two dielectric atoms. It will be seen that the result differs from those of both the real and virtual cavity models; however, when contributions to the decay rate originating from scattering by two dielectric atoms located at the same physical point are included, the calculation reverts to the virtual cavity model.

II. CALCULATION OF SECOND ORDER CONTRIBUTION

The source atom located at $R = 0$, has a $J = 0$ ground state and a $J = 1$ excited state, the frequency separation of the ground and excited state denoted by ω_0 . The uniformly distributed dielectric atoms have $J = 0$ ground states and $J = 1$ excited states, the frequency separation of the ground and excited state denoted by ω_j . At $t = 0$, the source atom is excited to the $m = 0$ excited state sublevel, the dielectric atoms are all in their ground states, and there are no photons in the field. The process we consider is radiation emitted by the source atom that is scattered by two dielectric atoms back to the source atom. It is assumed that $\omega_j \ll \omega_0 \ll \omega_0 + \omega_j$ but that $\omega_j \ll \omega_0 \ll (\omega_0 + \omega_j) \ll 1$ [rotating wave approximation (RWA)].

We use a multipolar Hamiltonian [8]. The free part is

$$H_0 = \frac{\omega_0}{2} \sum_{j,m=1}^X X^{(j)} + \sum_{j,m=1}^X \frac{\omega_j}{2} z^{(j)}(m) + \sum_k \omega_k a_k^\dagger a_k; \quad (1)$$

where $z = (\sum_i \hbar^2 j_i - \sum_l \hbar^2 l_l)$, \sum_i and \sum_l are the $m = 0$ excited and $J = 0$ ground state eigenkets of the source atom, respectively, $z^{(j)}(m) = \sum_{j_1} \langle j_1 | z^{(j)} | m \rangle \sum_{j_2} \langle j_2 | z^{(j)} | m \rangle$ is the population difference operator between excited state $|j_1 = 1; m\rangle$ and ground state $|j_2 = 0; g\rangle$ of dielectric atom j , and a_k is the annihilation operator for a photon having momentum k and polarization λ . A summation convention is used, in which any repeated symbol on the right hand side of an equation is summed over, unless it also appears on the left-hand side of the equations.

The interaction part of the Hamiltonian is,

$$V = d_0 \frac{D(0)}{\omega_0} + \sum_j d_j \frac{D(R_j)}{\omega_j}; \quad (2)$$

where d_0 and d_j are the dipole operators of the source atom located at the origin and a dielectric atom located at position R_j respectively. The operator D is the displacement field

having positive frequency component

$$D_+(R) = i \sum_k \frac{r}{2\epsilon_0 V} \hat{e}_k^{(1)} a_k e^{ik \cdot R} \quad (3)$$

where V is the quantization volume and $\hat{e}_k^{(1)}$ is a unit polarization vector, with

$$\hat{e}_k^{(1)} = \cos \theta_k \cos \phi_k \hat{x} + \cos \theta_k \sin \phi_k \hat{y} - \sin \theta_k \hat{z} \quad (4)$$

$$\hat{e}_k^{(2)} = \sin \theta_k \hat{x} + \cos \theta_k \hat{y} \quad (5)$$

In the RW A, one can write

$$V = \sum_k \tilde{g}_k (a_k + a_k^\dagger) + \sum_{k; m} \tilde{g}_k^0 (m) + (j) (m) a_k e^{ik \cdot R} + g_k^0 (m) a_k^\dagger (j) (m) e^{-ik \cdot R} \quad (6)$$

$$g_k = i \frac{r}{2\epsilon_0 V} (\hat{e}_k^{(1)})_0 \quad (7)$$

$$g_k^0 = i \frac{r}{2\epsilon_0 V} (\hat{e}_k^{(1)})_m \quad (8)$$

where the \tilde{g}_k are raising and lowering operators for the source atom and $(j) (m)$ are raising and lowering operators between the excited state $|j=1; m\rangle$ and the ground state $|j=0; g\rangle$ of dielectric atom j , $(\hat{e}_k^{(1)})_0$ is the reduced matrix element of the dipole operator d_0 and $(\hat{e}_k^{(1)})_m$ is that of d_j between ground and excited state manifolds. The $(\hat{e}_k^{(1)})_1 = \frac{(\hat{e}_k^{(1)})_x - i(\hat{e}_k^{(1)})_y}{2}$, $(\hat{e}_k^{(1)})_0 = (\hat{e}_k^{(1)})_z$ are spherical components of the polarization vectors. The source atom interacts only with the z component of the radiation field.

The calculation proceeds as in Ref. [6], with the addition of terms that couple dielectric atoms to dielectric atoms via the radiation field. After eliminating intermediate states involving the radiation field, one arrives at

$$\dot{b}_2 = -\gamma_2 b_2 - i \sum_j G_{0; m_j} (R_j; !) b_{m_j} (t) \quad (9a)$$

$$\dot{b}_{m_j} = -\gamma_{m_j} b_{m_j} - i \sum_j G_{m_j; 0} (R_j; !) b_2 (t) - \sum_s G_{m_j; m_s} (R_j - R_s; !) b_{m_s} (t); \quad (9b)$$

where $\gamma_2 = 2 \Gamma_0^2 = 3\gamma^3$ and $\gamma_{m_j} = 2 \Gamma_0^2 = 3\gamma^3$ are (half) the excited state decay rate of the source and dielectric atoms, respectively, b_2 is the state amplitude for the source atom to be in state $|j=1; m=0\rangle$ and all dielectric atoms in their ground states, and b_{m_j} is the state amplitude for dielectric atom j to be in excited state $|j=1; m\rangle$ all other atoms in

their ground states. We have set $b_2(t) = b(t)$ and $b_{m_j}(t) = b_j(t)$ on the assumption that $R_0 = c; R_0 = c - 1$, where R_0 is the sample size. The quantity $G_{m_j, m_s^0}(R_j, R_s; !)$ is a propagator for scattering from a dielectric atom in sublevel m_j at position R_j to one in sublevel m_s at position R_s given by

$$G_{m_j, m_s^0}(R_j; !) = \frac{3}{8} \frac{1}{!^3} \int_0^Z dt \int_0^Z d_1 \int_0^Z d_k !^3 e^{i(!_k - !_1)t} \int_0^Z d_k \binom{!}{k}_{m_j} \binom{!}{k}_{m_s^0} e^{ik \cdot R_j}; \quad (10)$$

while $G_{m_j, 0}(R_j; !)$ is a propagator for scattering from the source atom to a dielectric atom in sublevel m_j at position R_j . In what follows we ignore the difference between $!_0$ and $!$, consistent with the RWA.

In order to solve Eqs. (9a,9b), we assume that b_2 varies slowly on the time scale $1 = \dots$. If $b_{m_j} = Y_{m_j} e^{-i t}$, Eqs.(9a,9b) are transformed to

$$b_2 = b_2 + \int_0^! G_{0, m_j}(R_j; !_0) Y_{m_j}(t) \quad (11a)$$

$$\int_0^! (i - \dots) Y_{m_j} = \int_0^! G_{m_j, 0}(R_j; !_0) b_2(t) + \int_0^! G_{m_j, m_s^0}(R_j, R_s; !_0) Y_{m_s^0}(t) \quad (11b)$$

The formal solution for b_2 is

$$b_2 = b_2 + \int_0^! G_{0, m_j}(R_j; !_0) \frac{1}{i + \dots} \int_0^! G_{m_s^0, 0}(R_s; !_0) b_2 \quad (12)$$

where G is an $3N \times 3N$ matrix having matrix elements $G_{m_j, m_s^0}(R_j, R_s; !_0)$. This can be expanded as a power series in N with $\dots = \frac{4}{\dots}$: To second order, one finds

$$b_{2,0} = b_{2,0} + \int_0^! \left(\frac{k_0^3}{6N} \right)^2 G_{0, m_j}(R_j; !_0) G_{m_j, 0}(R_j; !_0) + \int_0^! \left(\frac{k_0^3}{6N} \right)^2 G_{0, m_j}(R_j; !_0) G_{m_j, m_s^0}(R_j, R_s; !_0) G_{m_s^0, 0}(R_s; !_0) \quad (13)$$

The term linear in the density gives the first order local field correction $\dots^{(1)} = \frac{7}{6} N$ [6], which agrees with both the virtual and real cavity models to this order.

We now calculate the second order correction,

$$\dots^{(2)} = \int_0^! \left(\frac{k_0^3}{6N} \right)^2 G_{0, m_j}(R_j; !_0) G_{m_j, m_s^0}(R_j, R_s; !_0) G_{m_s^0, 0}(R_s; !_0): \quad (14)$$

The sum over R_j and R_s can be converted to integrals using $\dots = \int_0^! \dots dR$. In this manner, one finds

$$\dots^{(2)} = \int_0^! \left(\frac{k_0^3}{6} \right)^2 \int_0^! \int_0^! dR_2 dR_1 G_{0, m_j}(R_2; !_0) G_{m_j, m_s^0}(R; !_0) G_{m_s^0, 0}(R_1; !_0) \quad (15)$$

where $R = R_2 - R_1$: The next step is to evaluate the $G_{m_j m_s^0}(R; \hat{R})$. The details of the calculation are given in the Appendix and one obtains

$$G_{11} = \frac{p}{4} h_0(k_0 R) Y_{0,0}(\hat{R}) - \frac{1}{2} \frac{4}{r} h_2(k_0 R) Y_{2,0}(\hat{R}); \quad (16a)$$

$$G_{00} = \frac{p}{4} h_0(k_0 R) Y_{0,0}(\hat{R}) + \frac{4}{5} h_2(k_0 R) Y_{2,0}(\hat{R}); \quad (16b)$$

$$G_{1;1} = \frac{3}{2} \frac{8}{r} h_2(k_0 R) Y_{2;2}(\hat{R}); \quad (16c)$$

$$G_{-1;1} = \frac{3}{2} \frac{8}{r} h_2(k_0 R) Y_{2;2}(\hat{R}); \quad (16d)$$

$$G_{1;0} = \frac{3}{2} \frac{4}{r} h_2(k_0 R) Y_{2;1}(\hat{R}); \quad (16e)$$

$$G_{-1;0} = \frac{3}{2} \frac{4}{r} h_2(k_0 R) Y_{2;1}(\hat{R}); \quad (16f)$$

where $Y_{l,m}(\hat{R})$ is a spherical harmonic and $k_0 = \omega/c$. The remaining $G_{m_j m_s^0}$ s are obtained using $G_{-1;1} = G_{11}$, $G_{0;1} = G_{1;0}$, and $G_{0;1} = G_{-1;0}$: The spherical Hankel functions of the first kind, $h_0(k_0 R)$ and $h_2(k_0 R)$; conform to the appropriate boundary conditions in which only outgoing scattered waves are considered.

The calculation for $\frac{(2)}{1;1}$ is tedious, since it involves contributions from nine terms. We will show how to calculate one specific contribution, $m_j = 1, m_s^0 = 1$, and then give the final results for the other components. Substituting Eqs. (16a,16e) in Eq. (15), we find

$$\begin{aligned} \frac{(2)}{1;1} &= \frac{(N)^2 \left(\frac{k_0^3}{6}\right)^2 \frac{6}{5} \int_{\mathbb{Z}} \int_{\mathbb{Z}} dR_1 dR_2 h_2(k_0 R_2) Y_{2,1}(\hat{R}_2) h_0(k_0 R_{21}) Y_{0,0}(\hat{R}_{21}) h_2(k_0 R_1) Y_{2;1}(\hat{R}_1)}{(N)^2 \left(\frac{k_0^3}{6}\right)^2 \frac{3}{5} \int_{\mathbb{Z}} \int_{\mathbb{Z}} dR_1 dR_2 h_2(k_0 R_2) Y_{2,1}(\hat{R}_2) h_2(k_0 R_{21}) Y_{2,0}(\hat{R}_{21}) h_2(k_0 R_1) Y_{2;1}(\hat{R}_1)} \\ & \quad (17) \end{aligned}$$

To evaluate this, we expand the Hankel functions as [9]

$$\begin{aligned} h_1(k_0 R_{21}) Y_{1,m}(\hat{R}_{21}) &= i^{l_1+l_2-1} (-1)^{l_2+m} P \frac{1}{4(2l_1+1)(2l_2+1)} \\ & \quad h \begin{matrix} 0 & 1 & 0 & 1 \\ (R_2 & R_1) + (-1)^l (R_1 & R_2) @ \begin{matrix} l_1 & l_1 & l_2 & A \\ 0 & 0 & 0 & \end{matrix} @ \begin{matrix} l_1 & l_1 & l_2 & A \\ m_1 & m & m_2 & \end{matrix} \end{matrix} \\ & \quad h_{l_1}(k_0 R_{>}) j_{l_2}(k_0 R_{<}) Y_{l_1, m_1}(\hat{R}_{>}) Y_{l_2, m_2}(\hat{R}_{<}) \end{aligned} \quad (18)$$

where $\begin{matrix} \dots \\ \dots \\ \dots \end{matrix}$ is a 3-j symbol, $j_l(x)$ is a spherical Bessel function, and $R_{>} (R_{<})$ is the larger (smaller) of R_1 and R_2 . When this expansion is used in Eq. (17), the angular integration

selects out only $l=2;0$ and $m=1;1;0$ terms, such that

$$\frac{^{(2)}(1;1)}{7} = \frac{(N)^2}{7} \int_0^{Z_1} d_2 \int_0^{Z_2} d_1 \frac{1}{2} h_2(z_2) h_2(z_1) h_2(z_2) j_2(z_1) \quad (19)$$

with $z_1 = k_0 R_1$, $z_2 = k_0 R_2$. To evaluate the above integral, we add a convergence factor $e^{-\epsilon z}$, and eventually take the limit $\epsilon \rightarrow 0$. The imaginary part of the integral diverges as $\epsilon \rightarrow 0$, but the real part is finite and gives the local field correction to the decay rate. The result is

$$\frac{^{(2)}(1;1)}{7} = \frac{15}{112} (N)^2$$

and the corresponding results for the other terms are [10]

$$\begin{aligned} \frac{^{(2)}(1;1)}{7} &= \frac{^{(2)}(1;1)}{7} = \frac{15}{112} (N)^2; \\ \frac{^{(2)}(0;0)}{63} &= \frac{25}{63} (N)^2; \\ \frac{^{(2)}(0;1)}{28} &= \frac{^{(2)}(1;0)}{28} = \frac{^{(2)}(1;0)}{28} = \frac{^{(2)}(0;1)}{28} = \frac{3}{28} (N)^2; \\ \frac{^{(2)}(1;1)}{56} &= \frac{^{(2)}(1;1)}{56} = \frac{3}{56} (N)^2; \end{aligned}$$

The total second order correction to the decay rate is

$$\frac{^{(2)}}{7} = 2 \frac{^{(2)}(1;1)}{7} + \frac{^{(2)}(0;0)}{63} + 4 \frac{^{(2)}(0;1)}{28} + 2 \frac{^{(2)}(1;1)}{56} = \frac{71}{72} (N)^2; \quad (20)$$

This result differs from both the virtual $\frac{51}{72} (N)^2$ and real $\frac{19}{72} (N)^2$ cavity models.

Our result can be compared with Fleischhauer's [5]: The Fourier transform of $G_{m,m^0}(R; \omega)$ is the tensor field propagator $F^{(0)}(q; \omega)$ in his paper, differing only by prefactors. The integral (15) can then be done in either coordinate or momentum space. The momentum space integration gives a different result than our coordinate space calculation above. This surprising discrepancy can be explained by the way we expand $h_1(k_0 R_{21}) Y_{1,m}(\hat{R}_{21})$. The expansion we used is valid for $R_1 > R_2$ or $R_2 > R_1$, but is not defined for $R_1 = R_2$. For a wellbehaved integral this will not make any difference since $R_1 = R_2$ contributes a set of measure zero. In the present case, however, where the dipole-dipole interaction between dielectric atoms diverges when one atom is on top the other, i.e. when $R_1 = R_2$, the contribution from $R_1 = R_2$ can be finite.

It is not easy to estimate this contribution in the original form of the integral (17). Instead, it proves useful to Fourier transform just one of the G_{m,m^0} in the integrand. As an

example, we consider the integral in the second term of Eq. (17)

$$I = \int \int dR_2 dR_1 h_2(kR_2) Y_{2;1}(\hat{R}_2) h_2(kR_{21}) Y_{2;0}(\hat{R}_{21}) h_2(kR_1) Y_{2;-1}(\hat{R}_1) \quad (21)$$

We Fourier transform $h_2(kR_{21}) Y_{2;0}(\hat{R}_{21}) e^{-iR_{21}}$, using a convergence factor $e^{-R_{21}}$ that is physically connected with the boundary condition of outgoing spherical waves. Carrying out the Fourier transform in Eq. (21), we find

$$I = \frac{4}{k^3} \int \int dR_2 dR_1 h_2(kR_2) Y_{2;1}(\hat{R}_2) h_2(kR_1) Y_{2;-1}(\hat{R}_1) \int \frac{dp}{(2\pi)^3} \frac{p^2}{p^2 + i} Y_{2;0}(\hat{p}) e^{ip(R_2 - R_1)} \quad (22)$$

The angular integrations can be done by expanding e^{ipR_2} , e^{-ipR_1} in terms of spherical harmonics and Bessel functions. In this manner one obtains

$$I = \frac{1}{14} \frac{5(4\pi)^3 i}{k^3} \int \int dR_2 dR_1 R_2^2 R_1^2 h_2(kR_2) h_2(kR_1) \int \frac{dp}{(2\pi)^3} \frac{p^4}{p^2 + i} \quad (23)$$

We are interested only in the contribution in the region where $R_1 = R_2$. This contribution can be isolated by integrating R_2 from $R_1 - a$ to $R_1 + a$, and then integrating the resultant expression over p using the method of residues. In the limit that both a and ϵ tend to zero, one obtains the contribution I from the region $R_1 = R_2$ as

$$\text{Re}[I] = \text{Re} \left[\frac{1}{7} \frac{5 - 2i}{k^3} \int dR_2 R_2^2 h_2(kR_2) h_2(kR_2) \right] = \frac{5}{7k^6} \quad (24)$$

(the imaginary part of I diverges). The contribution from the sphere $R_1 = R_2$ is identical to that from $R_1 = R_2$ since all other points with $R_1 \neq R_2$ on the sphere are regular and contribute zero to the integral. The same calculation can be done for the first integral in the Eq. (17). For this term, there is no contribution from the region $R_1 = R_2$ (no delta function like term is found) since $h_0(kR_{21})$ has a lower order divergence at $R_{21} = 0$ than does $h_2(kR_{21})$.

Including contributions of the type (24), we find

$$\begin{aligned} \frac{(2)(0;0)}{3} &= \frac{1}{3} (N)^2; \\ \frac{(2)(1;1)}{48} &= \frac{(2)(1;-1)}{48} = \frac{7}{48} (N)^2; \\ \frac{(2)(0;1)}{12} &= \frac{(2)(1;0)}{12} = \frac{(2)(-1;0)}{12} = \frac{(2)(0;-1)}{12} = \frac{1}{12} (N)^2; \\ \frac{(2)(-1;1)}{8} &= \frac{(2)(1;-1)}{8} = \frac{1}{8} (N)^2; \end{aligned} \quad (25)$$

When these are summed the total $\frac{(2)}{24} = \frac{17}{24} (N)^2$ agrees with the virtual cavity result.

III. D I S C U S S I O N

The second order contribution to the modified spontaneous emission rate of an impurity atom in a disordered dielectric has been calculated using a microscopic theory. Depending on the manner in which overlapping atoms are treated, one arrives at different results. If the delta function contributions at $R_1 = R_2$ are included, the virtual cavity model is recovered, but if such terms are excluded, neither the real nor virtual cavity model results are found. It seems to us somewhat of an open question at this point as to whether or not such contributions can be uniquely calculated once Eq. (12) is expanded in a power series in the density. The reason for this is that the expansion parameter is not small as interatomic distances tend to zero. That the expansion can lead to divergences is already evident if the integrations are carried out using a different set of variables [10]. From physical considerations, however, the decay rate does not diverge, even for interparticle spacings much less than a wavelength. Actually, dielectric atoms within a sphere of radius $(\lambda/2)^{1/3}$ reradiate collectively; outside this radius, there is destructive interference resulting in some additional finite contribution to the decay rate. In dealing with a homogeneous dielectric, we have performed the ensemble average by integrating over all space assuming a constant density. This averaging process includes configurations where interparticle spacings are sufficiently small to invalidate the expansion (13). Nevertheless, the procedure has yielded finite results for the change in the decay parameter.

Different experiments support both the real and virtual cavity results [11]. The source atom in these experiments is usually an impurity ion in a protective molecular cage. No experiments of this nature have been carried out with impurity atomic radiators in a dielectric that consists of a dense atomic vapor. It may be possible to use an alkali metal atom as the source atom and rare gas atoms as the dielectric atoms. With such a system, one could not make the rotating wave approximation used in this paper, but the physics is not changed in any substantive manner. The key feature of the alkali metal - rare gas system is the extremely small quenching cross sections for rare gas collisions to inelastically change the electronic state of the alkali atom [12]. Any quenching cross sections would appear as a modification of the decay rate that would mask the sought after effect. For rare gas pressures on the order of 100 atmospheres, we estimate that a change in the decay rate of order of 3% could be observed. To increase the effect it is necessary to find radiator atoms whose first

excited state is radiatively coupled to the ground state and dielectric atoms whose lowest excited state is about 0.2eV above the energy of the excited state of the radiator. In this limit, quenching will be negligible, but the detuning is decreased from the alkali-rare gas system by a factor of 50. At the same time, it is necessary to achieve a high pressure for the dielectric atoms. A possible system would be Li radiators with a high density sodium dielectric; the energy mismatch of Li and Na is about 0.25eV, giving a correction factor to the lithium decay rate of $1:3 \times 10^{21}N$, where N is the sodium dielectric density in units of atom/cm³:

IV. ACKNOWLEDGMENTS

This research is supported by the National Science Foundation under Grants No. PHY-0244841 and the FOCUS Center Grant. PRB would like to thank Peter Milonni for helpful comments and Georg Raithel for a discussion concerning high pressure sodium lamps.

V. APPENDIX

In this appendix, we calculate explicitly $G_{1;1}(R;!)_1$ given in Eq. (10). The other $G_{m;m^0}(R;!)_1$ are calculated in a similar fashion. To carry out the angular integrations, one expands $e^{ik \cdot R}$ as

$$e^{ik \cdot R} = 4 \sum_{m=-1}^1 i^m Y_m(\hat{k}) Y_m(\hat{R}) j_1(kR); \quad (26)$$

uses the fact that $(\begin{smallmatrix} 1 \\ k \end{smallmatrix})_1 (\begin{smallmatrix} 1 \\ k \end{smallmatrix})_1 = \frac{1}{2} (1 + \cos^2 \theta) = \frac{p}{3} [2Y_{00}(\hat{k}) + \frac{1}{5} Y_{20}(\hat{k})]$, and the orthogonality of the spherical harmonics, to obtain

$$G_{1;1}(R;!)_1 = \frac{1}{2} \int_0^{\pi} d\theta \int_0^{2\pi} d\phi e^{i(l_k \cdot !)} [2Y_{00}(\hat{R}) j_0(kR) + \frac{1}{5} Y_{20}(\hat{R}) j_2(kR)] \quad (27)$$

The spherical Bessel function can be written in terms of spherical Hankel functions as $j_1(kR) = \frac{1}{2} [h_1(kR) + h_1(kR)]$, transforming Eq. (27) into

$$G_{1;1}(R;!)_1 = \frac{1}{2} \int_0^{\pi} d\theta \int_0^{2\pi} d\phi e^{i(l_k \cdot !)} [2Y_{00}(\hat{R}) [h_0(kR) + h_0(kR)] + \frac{1}{5} Y_{20}(\hat{R}) [h_2(kR) + h_2(kR)]];$$

In the calculation we always make the Wigner-Eisskopf approximation. Differences between $!;!_0$ and $!_k$ are neglected except they appear as exponential factors. In integrating

over k ; the $h_1(kR)$ terms give a contribution proportional to $(R=c+)$ while the $h_2(kR)$ terms give a contribution proportional to $(R=c)$. We retain only the $(R=c)$ contributions since they correspond to the retarded solution (outgoing spherical waves). As a consequence, we find

$$G_{1,1}(R;!) = \frac{P}{4} h_0(k_0 R) Y_{0,0}(\hat{R}) - \frac{1}{2} \frac{r}{5} h_2(k_0 R) Y_{2,0}(\hat{R}) \quad (28)$$

-
- [1] D. Toptygin, *J. Fluorenc.* 13, 201 (2003); A. Luks and V. Perinova, *Progress in Optics*, edited by E. Wolf (Elsevier, Amsterdam, 2002), vol. 43, pp. 295-431; A. Thranhardt, C. Ell, G. Khitrova, and H. M. Gibbs, *Phys. Rev. B* 65, 035327 (2002), and references therein.
- [2] S. Scheel, L. Knoll, D. G. Welsch, and S. M. Barnett, *Phys. Rev. A* 60, 1590 (1999), and references therein
- [3] J. Knoester and S. Mukamel, *Phys. Rev. A* 40, 7065
- [4] Pedro de Vries and Ad Lagendijk, *Phys. Rev. Lett.* 81, 1381 (1998) and reference therein
- [5] M. Fleischhauer *Phys. Rev. A* 60, 2534 (1999) and reference therein
- [6] P. R. Berman and P. W. Milonni *Phys. Rev. Lett.* 92, 053601 (2004)
- [7] M. E. Crenshaw and C. M. Bowden, *Phys. Rev. Lett.* 85, 1851 (2000)
- [8] E. A. Power *Introductory Quantum Electrodynamics* (Elsevier, New York, 1965) Chapter 8
- [9] M. Danos and L. C. Maximon *J. of Math. Phys.* 6, 766 (1965)
- [10] The contributions from other terms may involve factors such as $h_0(k_0 R_1) h_2(k_0 R_{21}) h_2(k_0 R_2)$. Direct expansion of $h_2(k_0 R_{21})$ gives a divergent result for the decay rate, when the contribution from $R_1 = R_2$ is excluded. Instead, we expand $h_0(k_0 R_1)$ in terms of R_{21} and R_2 and integrate over these two variables. This procedure leads to the finite results given in the paper. The fact that we get divergent results when directly expanding $h_2(k_0 R_{21})$ in terms of R_1 and R_2 can be traced to the fact that one finds different values for the integrals at $R_1 = R_2$ depending on whether the limit $R_1 = R_2$ is approached from above or below. If the contribution from $R_1 = R_2$ is included, it leads to a divergence that exactly cancels the divergence from the first integral and again leads to the same result given in the text.
- [11] Different experiments have lent support to the virtual-cavity formula [see, for instance, R. S. Meltzer, S. P. Feolov, B. Tissue, and H. B. Yuan, *Phys. Rev. B* 60, 14012 (1999)] as well as

the real-cavity formula [see, for instance, G . L . J . A . R i k k e n and Y . A . R . R . K e s s e n e r, Phys. Rev. Lett. 74, 880 (1995) or F . J . P . S c h u u m a n s, D . T . N . d e L a n g, G . H . W e g d a m , R . S p r i k, and A . L a g e n d i j k, Phys. Rev. Lett. 80, 5077 (1998)]

[12] See, for example, J . H . S t a m p e r, Phys. Rev. J. Chem Phys. 43, 759 (1965); M . G . E d w a r d s, J. Phys. B 2, 719 (1969); E . S p e l l e r, B . S t a u d e n m a y e r, and V . K e m p t e r, Z. Phys. A 291, 311 (1979).