Quantum Arnol'd di usion in a rippled waveguide

V.Ya.Dem ikhovskii¹, F.M. Izrailev² and A.I.M alyshev¹

¹N izhny N ovgorod State U niversity, 603950, N izhny N ovgorod, G agarin ave. 23, Russia ²Instituto de Fisica, U niversidad Autonom a de Puebla, Apdo. Postal J-48, Puebla 72570, M exico

(January 9, 2022)

We study the quantum A mol'd di usion for a particle moving in a quasi-1D waveguide bounded by a periodically rippled surface, in the presence of the time-periodic electric eld. It was found that in a deep sem iclassical region the di usion-like motion occurs for a particle in the region corresponding to a stochastic layer surrounding the coupling resonance. The rate of the quantum di usion turns out to be less than the corresponding classical one, thus indicating the in uence of quantum coherent elects. A nother result is that even in the case when such a di usion is possible, it term inates in time due to the mechanism similar to that of the dynamical localization. The quantum A mol'd di usion represents a new type of quantum dynamics, and may be experimentally observed in measurements of a conductivity of lowdimensional mesoscopic structures. As is well known, one of the mechanisms of the dynam ical chaos in the Ham iltonian systems is due to the interaction between nonlinear resonances [1]. When the interaction is strong, this leads to the so-called global chaos which is characterized by a chaotic region spanned over the whole phase space of a system, although large isolated islands of stability may persist. For a weak interaction, the chaotic motion occurs only in the vicinity of separatrices of the resonances, in accordance with the Kolm ogorov-Amol'd-Moser (KAM) theory (see, for example, Ref. [2]). In the case of two degrees of freedom (N = 2), the passage of a trajectory from one stochastic region to another is blocked by KAM surfaces.

The situation changes drastically in many-dimensional (N > 2) systems for which the KAM surfaces no longer separate stochastic regions surrounding di erent resonances, and chaotic layers of the destroyed separatrices form a stochastic web that can cover the whole phase space. Thus, if the trajectory starts inside the stochastic web, it can di use through-out the phase space. This weak di usion along stochastic webs was predicted by A mol'd in 1964 [3], and since that time it is known as a very peculiar phenom enon, how ever, universal form any-dimensional nonlinear H am iltonian systems (see, for example, review [4] and references therein).

Recently, much attention has been paid to the chaotic dynamics of a particle in a rippled channel (see, for example, Refs. [5{9]). The main interest was in the quantum -classical correspondence for the conditions of a strong chaos. Speci cally, in Ref. [5] the transport properties of the channel in a ballistic regime were under study. Energy band structure, the structure of eigenfunctions and density of states have been calculated in [6,7]. The quantum states in the channel with rough boundaries, as well as the phenom ena of quantum localization have been analyzed in [9]. The in uence of an external magnetic eld for narrow channels was investigated in [8]. These studies may have a direct relevance to the experiments with periodically modulated conducting channels. In this connection one can mention the investigation [10] of transport properties of a mesoscopic structure (sequence of quantum dots) with the periodic potential form ed by metallic gates.

In contrast with the previous studies, below we address the regime of a weak quantum

2

chaos which occurs along the nonlinear resonances in the presence of an external periodic electric eld. Our goal is to study the properties of the quantum A mol'd di usion which m ay be observed experimentally. The approach we use is based on the theory developed in Ref. [11] by m aking use of a simple m odel of two coupling nonlinear oscillators, one of which is driven by two-frequency external eld.

We study the A mold di usion in a periodic quasi-one dimensional waveguide with the upper prolegiven in dimensionless variables by the function $y = d + a \cos x$. Here x and y are the longitudinal and transverse coordinates, d is the average width, and a is the ripple amplitude. The low prole is assumed to be at, y = 0. The nonlinear resonances arise due to the coupling between two degrees of freedom, with the following resonance conditions,

$$= T_x = T_y = !_y = !_x$$
 (1)

Here T_y is the period of a transverse oscillation inside the channel, T_x is the time of ight of a particle over one period of the waveguide, $!_x$ and $!_y$ are the corresponding frequencies, and is the rational number.



FIG.1. Some of the coupling resonances for di erent values of , with the isoenergetic curve E = 160000 in the frequency plane. The locations of resonances are shown by dashed lines.

The mechanism of the classical A mol'd di usion in this system is illustrated in Fig. 1. Here some of the resonance lines for di erent values of are shown on the $!_x$, $!_y$ -plane. A lso, the curve of a constant energy is shown determ ined by the equation

$$!_{x}^{2} + \frac{!_{y}d}{!_{x}} = 2m E :$$
 (2)

Here E and m are the dimensionless kinetic energy and particle mass, respectively, which we set to unity in what follows. The neighboring coupling resonances are isolated one from another by the KAM -surfaces, therefore, for a weak perturbation the transition between their stochastic layers is forbidden. Such a transition could occur in the case of the resonance overlap, i.e. in the case of the global chaos only. In the absence of an external eld, the passage of a trajectory along any stochastic layer (this direction is shown at Fig. 1 by two arrows) is also in possible because of the energy conservation. However, the external timeperiodic eld removes the latter restriction, and a slow di usion along stochastic layers becomes possible.

An external electric eld with the corresponding potential V (y;t) = $f_0y(\cos_1t + \cos_2t)$ gives rise to two main resonances with the frequencies $!_y = _1$ and $!_y = _2$. Their locations are shown in Fig.1 by the dashed lines. In order to calculate the di usion rate, we consider a part of the A mol'd stochastic web created by three resonances, namely, by the coupling resonance and two guiding resonances with frequencies $_1$ and $_2$. Therefore, we chose the initial conditions inside the stochastic layer of the coupling resonance. To avoid the overlapping of the resonances, we assume the relation $f_0=a = 1000$ is full led.

For the further analysis it is convenient to pass to the curvilinear coordinates x_i° in which both boundaries are at [12]. The covariant coordinate representation of the Schrödinger equation has the following form,

$$\frac{1}{2^{\mathbf{p}}\overline{g}}\frac{\varrho}{\varrho\mathbf{x}_{i}^{\circ}}\mathbf{p}\overline{g}g_{ij}\frac{\varrho}{\varrho\mathbf{x}_{j}^{\circ}} = E$$
(3)

where g_{ik} is the metric tensor, g det (g_j) . Here we use the units in which the Plank's constant and elective mass are equal to unity. As a result, the new coordinates are

$$x^{0} = x; \qquad y^{0} = \frac{y}{1 + \cos x}$$

$$\tag{4}$$

where = a=d. In these coordinates the boundary conditions are $(x^0; 0) = (x^0; d) = 0$ and the metric tensor is

$$g_{ij} = \begin{pmatrix} 0 & 1 & \frac{x^{0} \sin x^{0}}{1 + \cos x^{0}} & 1 \\ \frac{x^{0} \sin x^{0}}{1 + \cos x^{0}} & \frac{1 + \frac{2}{x} x^{02} \sin^{2} x^{0}}{(1 + \cos x^{0})^{2}} & A \end{pmatrix};$$
(5)

with the orthonorm ality condition,

$$\sum_{i=j}^{2} p_{\overline{g}} dS^{0} = ij:$$
 (6)

If the ripple amplitude a is small compared to the channel width d, than keeping only the rst-order terms in in the Schrödinger equation (3), we obtain the following H am iltonian [12],

$$\hat{\mathbf{H}} = \hat{\mathbf{H}}_{0} (\mathbf{x}; \mathbf{y}) + \hat{\mathbf{U}} (\mathbf{x}; \mathbf{y}); \tag{7}$$

where

$$\hat{H}_{0} = \frac{1}{2} \frac{\theta^{2}}{\theta x^{2}} + \frac{\theta^{2}}{\theta y^{2}}^{!}$$
(8)

and

$$\hat{\mathbf{U}} = \frac{1}{2} 2 \cos x \frac{\theta^2}{\theta y^2} \quad 2y \sin x \frac{\theta^2}{\theta x \theta y} \quad y \cos x \frac{\theta}{\theta y} \quad \frac{1}{2} \cos x \quad \sin x \frac{\theta}{\theta x} \stackrel{!}{:} \tag{9}$$

Here and below we om itted primes in coordinates x^0 and y^0 .

Since the H am iltonian is periodic in the longitudinal coordinate x, the eigenstates are B loch states. This allows us to write the solution of the Schrodinger equation in the form $(x;y) = \exp(ikx)_k (x;y)$ where $_k (x + 2;y) = _k (x;y)$. For an in nite periodic channel the B loch wave vector k has continuous values, in particulary, 1=2 k 1=2 in the rst B rillouin zone.

By expanding $_{k}(x;y)$ in the double Fourier series the eigenstates can be written as

$$^{k}(x;y) = e^{ikx} \int_{n,m}^{X} c^{k}_{nm} \int_{nm}^{0} (x;y)$$
(10)

where

$$\int_{nm}^{0} (\mathbf{x}; \mathbf{y}) = \frac{s}{Ld} e^{in \mathbf{x}} \sin \frac{m \mathbf{y}}{d}$$
(11)

are the eigenstates of the unperturbed H am iltonian $\hat{H}_0(x;y)$ with the corresponding eigenvalues

$$E_{nm}^{0} = \frac{1}{2} (n+k)^{2} + \frac{2m^{2}}{d^{2}}$$
(12)

for the considered case with L 2. Now we proceed to solve the system of algebraic equations for the coe cients c_{nm}^{k} ,

$$E (k)c_{nm}^{k} = \frac{1}{2} (n+k)^{2} + \frac{2m^{2}}{d^{2}} c_{nm}^{k} + \sum_{n^{0}m^{0}} U_{(k+n)m;(k+n^{0})m^{0}} c_{nm}^{k} :$$
(13)

Here the matrix elements are

Following Refs. [11], we analyze the dynamics in the vicinity of the main coupling resonance

= 1 which is determined by the condition $!_{n_0} = !_{m_0}$ where $!_{n_0} = E_{n_0+1}(k)$ $E_{n_0}(k) = k + n_0 + 1 = 2$ and $!_{m_0} = E_{m_0+1}$ $E_{m_0} = ^2(2m_0 + 1) = 2d^2$. In a deep sem iclassical region where n_0 1 and m_0 1, one can write $!_{n_0}$ n_0 and $!_{m_0}$ $^2m_0 = d^2$. It should be noted that the similar resonance condition can be satisfied for negative n_0 also in the case when

 $n_0 = 2^n - 2^n + 1$, which corresponds to the particles moving in the opposite direction. Below we use the fact that when $j_0 j = 1$ the two resonances (two sets of states), corresponding to $n_0 > 0$ and $n_0 < 0$ are not coupled for all k apart from the center and edges of the B rillouin zone. This fact is due to the anti-unitary symmetry of the H am iltonian for all values of k, apart from the indicated above. The properties of the energy spectra and eigenstates for speci c values k = 1=2 and k = 0 will be discussed elsewhere.

In the vicinity of the resonance it is convenient to introduce new indexes r = n n_0 and $p = r + (m m_0)$. Then, instead of the system (13) we obtain,

$$E(k)c_{rp}^{k} = p!_{m_{0}}c_{rp}^{k} + \frac{1}{2}r^{2} + \frac{2}{d^{2}}(p r)^{2}c_{rp}^{k} + \frac{X}{r^{0}p^{0}}U_{k+n_{0}}r_{p}r_{m_{0}}r_{k+n_{0}}r^{0}p^{0}r^{0} + m_{0}c_{r}^{k}c_{p}^{k}; (15)$$

Here we count energy E (k) from the level $E_{n_0m_0}^0$ (k).

Num erical calculation of the solution of these equation gave the following results. First, as in Ref. [11], the energy spectrum consists of a number of Mathieu-like groups corresponding to the coupling resonance. These groups are separated one from another by the energy $!_{n_0}$. The structure of energy spectrum in each group is typical for a quantum nonlinear resonance. Inside the resonance, the lowest levels are practically equidistant, the accumulation point corresponds to the classical separatrix and all states are non-degenerate. The states above the separatrix are quasi-degenerate due to to the rotation in opposite directions.

In accordance with the spectrum structure it is convenient to characterize the states at coupling resonance by two indexes: group number q and s | level number inside the group. Correspondingly, the energy of each group can be written as follows,

$$E_{q;s}(k) = !_{n_0}(k)q + E_{q;s}^{M}(k);$$
(16)

where E_{qs}^{M} is the M athieu-like spectrum for one group. The indexes q and s correspond to fast and slow variables characterizing the motion inside the classical coupling resonance.

Let us consider now the dynamics of a charged particle in the rippled channel in the presence of the time dependent electric eld described by the potential V (y;t) = $f_0y(\cos_1t + \cos_2t)$. We assume that the frequencies $_1$ and $_2$ are chosen to fulll the condition $!_{n_0} = (_1 + _2)=2$ in order to provide equal driving forces for a particle inside the stochastic layer of the separatrix under consideration. Speci cally, we take, $!_{n_0} = 400; _1 = 350; _2 = 450$, therefore, the period T of the perturbation is, T = 7 - 2 = 1 = 9 - 2 = 2 - 0.126.

Since the total H am iltonian is periodic in time, one can write the solution of the nonstationary Schrödinger equation as $(x;y;t) = \exp((i_0 t)u_0 (x;y;t))$, where $u_0 (x;y;t)$ is the quasienergy (QE) function and $_q$ is the quasienergy. As is known, the QE functions are the eigenfunctions of the evolution operator $\hat{U}(T)$ of the system for one period of the perturbation. The procedure to determ ine this operator was described in details in R ef. [11]. The matrix elements $U_{q;s;q^0;s^0}$ (T) of the evolution operator can be calculated by means of the num erical solution of the non-stationary Schrödinger equation. Then, the evolution matrix $U_{q;s;q^0;s^0}$ (NT) for N periods can be easily obtained.

Our goal is to analyze the dynam ics of a particle placed inside the separatrix under the condition that the coupling and two driving resonances do not overlap. The evolution of any initial state can be computed using the evolution matrix as follows,

$$C_{q;s}(N T) = \sum_{q^{0};s^{0}}^{X} U_{q;s;q^{0};s^{0}}(N T) C_{q^{0};s^{0}}(0) :$$
(17)



FIG.2. Time dependence of the variance $_q$ for di erent initial states at coupling resonance for the group with q = 0: (a) the lowest level with s = 0, (b) the above-separatrix level with s = 45, (c) the near separatrix level with s = 22. Here a = 0.01 and $f_0 = 10$.

In Fig. 2 typical dependencies of the variance $H^2 = \frac{1}{n_0} \frac{1}{q}$ of the energy are shown versus the time measured in the number N of periods of the external perturbation, for di erent initial conditions. The quantity $\frac{1}{q}$ is de ned as follows,

$$_{q} = \int_{q}^{X} (q \ q)^{2} \int_{s}^{X} f_{q;s} f_{s} = \int_{q}^{X} (q \ q)^{2} \int_{s}^{X} f_{q;s} f_{s}$$
(18)

The data clearly demonstrate a di erent character of the evolution of the system in dependence on the initial state. For the state taken from the center of the coupling resonance, as well as above the separatrix, the variance oscillates in time, in contrast with the state taken from inside the separatrix. In the latter case, after a short time the variance of the energy increases linearly in time, thus manifesting a di usion-like spread of the wave packet.

In order to characterize the speed of the di usion, we have calculated the classical di usion coe cient D_{cl} for the coupling resonance = 1, in the comparison with the quantum one, D_q, for di erent go er am plitude and initial wave number k \notin 0; 1=2, see Fig. 3. It was found that the quantum A mol'd di usion roughly corresponds to the classical one. However, one can see a system atic deviation which indicates that the quantum di usion is weaker than the classical A mol'd di usion. This deviation is due to the in uence of coherence quantum e ects which can be very strong even in a deep sem iclassical region. A s was shown in Ref. [13], these stabilizing quantum e ects are enhanced for the motion inside narrow stochastic layers surrounding the nonlinear resonances. This e ect is due a relatively sm all num ber M_s of quantum states belonging to the region of the separatrix, the fact which is crucial in the study of the quantum -classical correspondence for the system s with a weak chaos in the classical limit.



FIG.3. Classical D $_{cl}$ versus quantum D $_{q}$ di usion coe cients in the dependence on the am plitude a of the rippled prole.

In this connection it is instructive to estimate the number of the energy states that occupy the separatrix layer in our calculations. We have found that for a = 0.01 of the am – plitude characterizing the upper prole of the waveguide, the number of stationary states in the separatrix chaotic layer is about M_s 10, therefore, one can treat our results sem iclassically. Our additional calculation has shown that with the decrease of the am plitude a, the number M_s decreases, and for $1 = \frac{p}{a}$ 20 this number is of the order one. For this reason the last right point in Fig. 3 corresponds to the situation when the chaotic motion along the coupling resonance is completely suppressed by quantum elects. This elect is known as the \Shuryak border" [13]) which establishes the conditions for a complete suppression of classical chaos.

Since the di usion along the coupling resonance is e ectively one-dimensional, one can expect an Anderson-like localization which is known to occur in the presence of any weak disorder in low-dimensional structures. Indeed, the variance of the QE eigenstates of the evolution operator is nite in the q-space. This means that eigenstates are localized, and the wave packet dynamics in this direction has to reveal the saturation of the di usion. More speci cally, one can expect that the linear increase of the variance of the energy ceases after som e characteristic time.

In order to observe the dynam ical localization in ourm odel (along the coupling resonance inside the separatrix layer), one needs to analyze a long-time dynam ics of wave packets. O ur num erical study for large times (see curve (c) in Fig. 2) have revealed that after some time t 200T, the di usion-like evolution term inates for all range of the amplitude a. Speci cally, on a large time scale the variance starts to oscillate around a mean value. This e ect is of the same origin as the so-called dynam ical localization which was discovered in the kicked rotorm odel [14,15]. O ne should note that the dynam ical localization is, in principle, di erent from the Anderson localization, since the latter occurs for the models with random potentials. In contrast, our model is the dynam ical one (without any random ness), and a kind of pseudo-random ness is due to the quantum chaos mechanism.

In conclusion, we have studied the quantum A mold di usion for a particle in the quasi-

10

1D rippled waveguide under the time-periodic electric eld. This di usion is known to occur in the corresponding classical systems, below the threshold of an overlap of nonlinear resonances resulting in a strong chaos. As is known, in this case the chaotic motion is possible only for the trajectories inside narrow stochastic layers surrounding the nonlinear resonances. The classical A mol'd di usion is exponentially weak, however, it leads to the unbounded motion inside the resonance web created by the resonances of di erent orders.

The problem under study was to nd out whether the A mol'd di usion is feasible in quantum system s describing the electron motion in quasi-1D waveguides. Our results clearly demonstrate the peculiarities of the quantum A mol'd di usion and establish the conditions under which it can be observed experimentally. One of the most important results is that the quantum di usion is typically weaker than the corresponding classical one. The analysis has shown that the reason is the in uence of quantum coherent elects which are essentially strong for the motion in the vicinity of the nonlinear resonances, in the case when these resonances do not overlap. Moreover, we found that with a decrease of the perturbation responsible for the creation of stochastic layers, the quantum A mol'd di usion can be completely suppressed, thus leading to the absence of any chaos in the system.

A nother in portant result is the observation of the dynam ical localization of the A mol'd di usion, which is manifested by the term ination of the di usion on a large time scale. We have brie y discussed the mechanism of this phenomenon, relating it with the fam ous A nderson localization known to occur in low-dimensional disordered structures. Thus, in addition to the Shuryak border, the dynam ical localization is a new mechanism destroying the quantum A mol'd di usion.

Recently, the weak electron di usion was observed experimentally [16] in superlattices with stationary electric and magnetic elds. At certain voltage, the unbounded electron motion along a stochastic web changes the conductivity of the system and results in a large increase of the current ow through the superlattice. Similarly, one can expect that electron A mol'd di usion inside resonance stochastic layers can increase the high frequency conductivity of a rippled channel, and can be detected experimentally.

11

This work was supported by the program D evelopm ent of scientic potential of high school" of Russian M inistry of Education and Science and partially by the CONACYT (M exico) grant No 43730. A JM. acknow ledges the support of non-prot foundation Dy-nasty".

- B.V.Chirikov, Atom naya Energia 6,630 (1959) Engl. Transl. J.Nucl. Energy Part C: Plasm a Phys. 1, 253 (1960)].
- [2] A J.Lichtenberg and M.A.Lieberm an, Regular and Chaotic D ynam ics, Springer-Verlag, New York, 1992.
- [3] V J. A mol'd, DAN USSR 156 (1964) 9 (in Russian).
- [4] B.V. Chirikov, Phys. Rep. 52 (1979) 263.
- [5] G A. Luna-Acosta, A A. Krokhin, M A. Rodriguez, P H. Hemandez-Tejeda, Phys. Rev. B 54 (1996) 11410.
- [6] G A.Luna-Acosta, K.Na, LE.Reichl, A A.Krokhin, Phys. Rev. E 53 (1996) 3271.
- [7] G A. Luna-Acosta, J.A. Mendez-Bermudez, F.M. Izrailev, Phys. Rev. E 64 (2001) 036206.
- [8] C.S.Lent, M.Leng, J.Appl. Phys. 70 (1991) 3157.
- [9] F. M. Izrailev, J.A. M. endez-Berm udez, and G.A. Luna-Acosta, Phys. Rev. E 68 (2003) 066201; F. M. Izrailev, N. M. M. akarov, M. Rendon, arX iv cond-m at/0411739 (2004).
- [10] LP.Kouwenhoven et al, Phys.Rev.Lett. 65 (1990) 361.
- [11] V.Ya. Dem ikhovskii, F.M. Izrailev, and A.I. Malyshev, Phys. Rev. Lett. 88 (2002) 154101;
 V.Ya. Dem ikhovskii, F.M. Izrailev, and A.I. Malyshev, Phys. Rev. E 66 (2002) 036211.
- [12] V.Ya.Dem ikhovskii, S.Yu.Potapenko, A.M. Satanin, Fiz.Tekh.Poluprovodn.17 (1983) 213

[Sov.Phys.Sem icond.17 (1983) 137].

- [13] E.V. Shuryak, Zh. Eksp. Teor. Fiz. 71 (1976) 2039 (in Russian).
- [14] G. Casati, B.V. Chirikov, F.M. Izrailev, and J. Ford, Lect. Notes Phys., 93 (1979) 334.
- [15] B.V. Chirikov, F.M. Izrailev, and D.L. Shepelyansky, Sov. Sci. Rev. 2 (1981) 209.
- [16] T M .From hold et al., Nature 428 (2004) 726.