

Limitations of Quantum Simulation Examined by Simulating a Pairing Hamiltonian using Nuclear Magnetic Resonance

Kenneth R. Brown, Robert J. Clark, and Isaac L. Chuang

Center for Bits and Atoms, Research Laboratory of Electronics, & Department of Physics
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

(dated: January 20, 2002)

Quantum simulation uses a well-known quantum system to predict the behavior of another quantum system. Certain limitations in this technique arise, however, when applied to specific problems, as we demonstrate with a theoretical and experimental study of an algorithm to find the low-lying spectrum of a Hamiltonian. While the number of elementary quantum gates required does scale polynomially with the size of the system, it increases inversely to the desired error bound. Making such simulations robust to decoherence using fault-tolerance constructs requires an additional factor of $1/\epsilon$ gates. These constraints are illustrated by using a three qubit nuclear magnetic resonance system to simulate a pairing Hamiltonian, following the algorithm proposed by Wu, Byrd, and Lidar [1].

The unknown properties and dynamics of a given quantum system can often be studied by using a well-known and controllable quantum system to mimic the behavior of the original system. This technique of quantum simulation is one of the fundamental motivations for the study of quantum computation [2,4], and is particularly of interest because a quantum simulation may be performed using space and time resources comparable to the original system. Such "efficient" scaling is dramatically better than the exponentially large resource requirements to simulate any general quantum system with a classical computer, as Feynman originally observed [2].

Recent work has continued to arouse great interest in quantum simulation, because it offers the possibility of solving computationally hard problems without requiring the resources necessary for algorithms such as factoring [5] and searching [6]. Experimental results have demonstrated simulations of a truncated oscillator and of a three-body interaction Hamiltonian, using a nuclear magnetic resonance (NMR) quantum computer [7, 8], and explored various solid-state models on two qubit systems [9,12]. Interest has also extended to simulating complex condensed matter systems with quantum optical systems [13], demonstrated vividly by the observation of a superfluid to Mott insulator transition in a Bose-Einstein condensate [14].

Often overlooked in the discussion of quantum simulations, however, is the question of desired precision (or error) in the final measurement results. Current quantum simulation techniques generally scale poorly with desired precision; they demand an amount of space or time which increases as $1/\epsilon$, broadly translating into a number of quantum gates which grow exponentially with the desired number of bits in the final answer. Why is this scaling behavior so poor, and what is its physical origin?

Consider as a specific example the problem of calculating the energy gap between the ground state $|\psi_0\rangle$ and the first excited state $|\psi_1\rangle$, of a Hamiltonian H . It can be found using the following steps: 1) map the Hilbert

space of the system to be simulated to n qubits, 2) prepare the computer in the state $|\psi_0\rangle = \alpha_0 |\psi_0\rangle + \alpha_1 |\psi_1\rangle$, 3) evolve under the Hamiltonian for times t_k , 4) extract the phase difference as a function of time between the evolution of the ground and first excited state.

Two methods for calculating the phase difference, and thus ϵ , are as follows. The first method uses the phase estimation algorithm [15, 16]. This method relies on the quantum Fourier transform (QFT) and requires simulating the Hamiltonian for times $t_k = 2^k t_0$, for integer k from 0 to q . Since the input state is a superposition of $|\psi_0\rangle$ and $|\psi_1\rangle$, the measured phase will either be $E_0 t_0$ or $(E_0 + \epsilon) t_0$, where E_0 is the ground state energy. One needs to run the algorithm on average $1/\epsilon^2$ times to get both values and thus measure ϵ .

The second method does not use the QFT, and instead simulates the Hamiltonian for times $t_k = k t_0$, for integer k from 0 to Q , and then measures any operator M such that $\langle \psi_1 | M | \psi_1 \rangle \neq 0$. Typically, any operator that does not commute with the Hamiltonian suffices. After calculating $\langle M(t_k) \rangle$, one classically Fourier transforms (FT) over the averaged values yielding a spectrum $M(\omega)$ with peaks at ω_0 and $\omega_0 + \epsilon$.

For fixed precision, obtaining up to error ϵ for fixed ϵ , both methods can be "efficient," in that the number of elementary steps (or quantum gates) required increases only polynomially with the number of qubits n , if the Hamiltonian can be efficiently simulated and the initial states efficiently prepared. A d -qubit Hamiltonian can be simulated with a number of gates of order $O(n^d)$ assuming two qubit interactions between any qubits [4]. If one assumes only nearest neighbor two qubit gates, it scales as $O(n^{d+1})$. Most physical systems of interest are described by two-body interactions which can be described by four qubit Hamiltonians.

Consequently, the challenge of designing efficient quantum simulations is choosing a property that can be efficiently extracted. However, no general measurement method is known which allows ϵ to be measured efficiently.

ciently with respect to the precision using such quantum simulations. For error ϵ , the number of digits of precision in the result is $\log(1/\epsilon)$, and both of the above methods require $1/\epsilon$ steps (or gates) to obtain this precision. In contrast, an efficient algorithm would only require a number of steps polynomial in $\log(1/\epsilon)$. The origin of this limitation lies not only in the inability to design efficient measurements, but also in the accumulation of errors which occurs in the course of performing a quantum simulation.

Here, we consider these limitations on the precision of results obtained by quantum simulations in the context of a specific algorithm for the simulation of pairing models, as proposed by Wu, Byrd and Lidar (WBL) [1], which follows the framework of the two methods described above. We present a study of the errors in its discrete time step implementation, and experimental results from a realization using a 3 qubit nuclear magnetic resonance (NMR) quantum computer, answering three questions: 1) What are the theoretical bounds on the precision of the quantum simulation? 2) How do faulty controls affect the accuracy of a simulation? 3) Can the theoretical bounds on precision be saturated by an NMR implementation?

The WBL algorithm uses the classical FT algorithm described above to solve the question of the low-lying energy gap in pairing Hamiltonians. Pairing Hamiltonians are used to describe both nuclear dynamics and superconductivity [17][19] and are usually written in terms of Fermionic creation and annihilation operators c^\dagger and c as

$$H_{\text{pair}} = \sum_{m=1}^n \frac{m}{2} (c_m^\dagger c_m + c_m^\dagger c_{m+1} + c_m c_{m+1}^\dagger) + \sum_{m,l=1}^n V_{m,l} (c_m^\dagger c_l^\dagger c_l c_m);$$

where n is the total number of modes, m is the onsite energy of a pair in mode m , and $V_{m,l}$ are the coupling constants between modes.

WBL map the pairing Hamiltonian onto the qubit Hamiltonian

$$H_{\text{pair}} = \sum_{m=1}^n \frac{m}{2} (Z_m) + \sum_{m < l} \frac{V_{m,l}}{2} (X_m X_l + Y_m Y_l);$$

where X_m ; Y_m ; and Z_m are the Pauli operators on the m th qubit and $m = m + V_{ll}$ (dropping an unimportant global energy shift, and using the standard convention $Z|j\rangle = (-1)^j|j\rangle$). The number of modes that can be simulated equals the number of qubits n , and the number of pairs equals the total number of qubits in the state $|j\rangle$. WBL show that for a specific number of pairs, one can approximately prepare the state $|j\rangle$ by quasiadiabatic evolution. Since H_{pair} is a 2-body Hamiltonian, the system's evolution can be efficiently simulated on a quantum computer for any number of qubits [3]. WBL propose implementation of their algorithm using an NMR quantum computer, in which the operator M is simply Z

for a single spin. An advantage of the ensemble nature of NMR is that a single measurement for a simulated time t yields $M(t)$. Fixing a maximum energy width and desired precision makes the FT independent of the number of qubits.

Let us begin by addressing the first question posed above, regarding theoretical bounds on the precision of this quantum simulation: how does the number of gates scale with the error? The WBL method requires constructing an operator that approximates the simulated Hamiltonian for time t_k . The classical FT then yields an error of $2^{-E_{\text{max}} t_k} = \epsilon$ where E_{max} is the largest detectable energy $1/t_0$ (~ 1). In the case of using phase estimation and the QFT, setting $2^q = Q$ yields the same precision. How long does it take to implement the Hamiltonian for a time t_0 compared to implementing a Hamiltonian for time t_0 ? In general, the operator is assumed to be constructed of repetitions of the basic time step and requires Q more gates or time. This leads to the number of gates scaling inversely with the error. A similar problem faced in quantum factoring is overcome in Shor's algorithm by a clever way to perform the modular exponentiation [5].

A second bound on the number of gates required arises in calculating the time required to perform the algorithm. Quantum simulations typically employ a Trotter formula to approximate a Hamiltonian from combinations of non-commuting Hamiltonians [16]. For example, given the ability to evolve under Hamiltonians H_A and H_B , one can approximate evolution under $H_A + H_B$ with bounded error. To lowest order, $\exp(-i t(H_A + H_B)) = (\exp(-i t H_A) \exp(-i t H_B))^k + \dots$, where for $k|H_A|H_B|kt^2 \ll 1$, the error is $O(t^2/k)$. Higher-order techniques can yield an error $O(t^{m+1}/k^m)$ at the cost of needing $O(2^m)$ more gates [20].

This approximation method leads to a subtle but important difficulty in reducing the gate count for simulations. It is apparent that the Trotter formula demands an exponential increase in the number of discrete gates for an exponential decrease in the error. However, from a Hamiltonian control perspective, this conclusion seems unfair, because the total time required can be small even if the gate count is high. Specifically, the gate $U_A(t=k) = \exp(-i t H_A = k)$ requires $1/k$ the time needed to implement $U_A(t)$. Therefore, the simple Trotter method given above requires only time $2t$, independent of k . This implies that "Trotterization" errors involved in approximating desired Hamiltonians can be reduced efficiently with respect to the time cost.

Unfortunately, this optimistic observation is incompatible with fault tolerant error correction [21, 22], which will likely be needed to extend simulation times beyond limits imposed by qubit decoherence times. This is because the fault-tolerant implementation of $U_A(t=k)$ takes approximately the same amount of time as the gate $U_A(t)$, whether using teleportation [23] or the Solovay-Kitaev approximation [24].

Consequently, fault-tolerant simulations using the Trotter formula and the FT/QFT require a number of gates and amount of time that scales as $1/\epsilon^2$. Circumventing this problem would require removing the inefficiency of Trotterization, and constructing methods to approximate $U_H(t)$ with error using $\text{poly}(\log(1/\epsilon))$ gates. However, such methods would imply that the approximation of $U_H(qt)$ could take only $\text{poly}(\log(q))$ more gates than the simulation of $U_H(t)$. Such a dramatic simplification may hold for specific $H(t)$, but is unlikely to be possible for general $H(t)$. The WBL algorithm studied here unfortunately does not scale efficiently when made fault-tolerant.

These theoretical bounds establish that present quantum simulations such as the WBL algorithm, using the QFT or the FT, require a number of gates which scales inversely with the desired answer precision for two reasons: fault-tolerant gate construction and the precision of a finite FT. Therefore, the time required for a d -qubit quantum simulation is $O(n^{d+r})$, where $r \geq 1$ varies depending on the approximation methods employed, and $r = 1$ when quantum error correction and fault-tolerant gates are not used.

We turn now to the second question, which concerns the impact of faulty controls in a real physical implementation of the WBL algorithm. Recall that the foundation of the WBL algorithm is approximation of the unitary evolution under H_{pair} , $U_{\text{pair}}(qt_0) = \exp(-iH_{\text{pair}}qt_0)$. An ideal NMR implementation accomplishes this by a repeatable pulse sequence $V_{\text{pair}}(t_0)$, where $U_{\text{pair}}(qt_0) = (V_{\text{pair}}(t_0))^q$. H_{pair} contains three noncommuting parts: $H_0 = \sum_{m=1}^M \frac{\omega_m}{2} (Z_m)$, $H_{XX} = \sum_{m < l} \frac{J_{ml}}{2} X_m X_l$, and $H_{YY} = \sum_{m < l} \frac{J_{ml}}{2} Y_m Y_l$. Assuming that the corresponding unitary operators $U_0(t)$, $U_{XX}(t)$, and $U_{YY}(t)$ can be implemented, $V_{\text{pair}}(t_0)$ can be constructed using the third order Trotter-Suzuki formula [20, 25]

$$V_{\text{pair}}(t_0) = [U_0(t_0=2k)U_{XX}(t_0=2k)U_{YY}(t_0=k)U_{XX}(t_0=2k)U_0(t_0=2k)]^k;$$

yielding an expected error $kU_{\text{pair}}(t_0) - V_{\text{pair}}(t_0)k = O(t_0^3/k^2)$.

However, this ideal procedure is not actually achieved in a real experiment because the unitaries are not direct implementations of the Hamiltonians but instead composed from a series of pulses. These pulses depend on assumptions about the system Hamiltonian that become unreasonable for short simulated times. The reason is that all real systems have small, often unknown, energy shifts that are averaged away for large simulated times. When not using a fault-tolerant construction these shifts can lead to faulty controls. In atomic physics, for example, undesired Stark shifts need to be carefully accounted for in order to get exact rotations [26].

Control errors in NMR quantum computation arise, for example, since single qubit gates require finite time




FIG. 1: Three qubit quantum circuits for the unitaries U_{XX} (top) and U_{YY} (bottom), implemented using method W1. These are depicted for simulating Hamiltonian H_2 (see text). For H_1 , $\frac{\pi}{2}$ pulses are applied to qubit c in parallel with those on a and b, and the decoupling X pulse is omitted.

and unwanted two qubit coupling occurs during this time. In a static magnetic field $B_0 \hat{z}$, the unitary evolution of a typical used spin system [11, 27] in the rotating frame is given by $U_{ZZ}(t) = \exp(-i \sum_{i,j} \frac{J_{ij}}{2} Z_i Z_j t)$, where the J_{ij} are the scalar coupling constants. The time t required for a radiofrequency (RF) pulse to rotate individual spins by radians is much smaller than the typical delay times t_d during which no RF is applied, $t \ll t_d \implies 1/J_{ij}$. Thus, it ordinarily suffices to approximate the RF pulses as δ -functions in time, implementing perfect single qubit rotations $R^i(\theta) = \exp(\frac{i}{2}(X_i \cos \theta + Y_i \sin \theta))$. However, this approximation breaks down as t_d becomes comparable to t , causing the expected evolution to be best described not by discrete one and two-qubit gates, but instead by the piecewise continuous time-dependent Hamiltonian $H_{\text{nmr}}(t) = \sum_i g_i(t)X_i + \sum_i f_i(t)Y_i + \sum_{i < j} \frac{J_{ij}}{2} Z_i Z_j$. This discrepancy leads to additional errors in implementations of quantum algorithms and simulations, which, for a small number of qubits, can be mitigated using optimal control techniques [10, 28].

The impact of such control errors in an NMR implementation of the WBL algorithm can be studied by comparing a baseline realization with no control error compensation (denoted W1) versus another with simple error compensation (denoted W2). The baseline W1 realization implements U_0 using composite pulses to create rotations about the \hat{z} axis, $U_0 = \prod_m R_{\hat{z}}^m(\pi/2) R_0^m(\pi) \prod_m R_{\hat{z}}^m(\pi/2)$; an equivalent method, used elsewhere [1, 10] temporarily shifts the rotating frame. Control errors arise in the simulation of U_{XX} and U_{YY} , which are generated by applying single qubit pulses to rotate the scalar coupling from the \hat{z} axis to the \hat{x} and \hat{y} axis, using the quantum circuit in Fig. 1. Control errors in this baseline realization are thus small only when delays needed to generate U_{XX} are long compared to the time required to perform single qubit gates, but also short enough that the Trotter error is small.

A simple, scalable compensation technique for control




FIG. 2: Frequency-domain spectra of Hamiltonian H_2 obtained using methods W1 (marked by circles) and W2 (diamonds). Dots are experimental data, and solid lines are Fourier transforms of an exponentially decaying sinusoid (with four free parameters) fit to the time dependent data. The width of the exact curve is taken to be the dephasing rate ($1/\tau$) of the ^{13}C nucleus.

errors provides a contrasting realization of the WBL algorithm for comparison. This W2 realization accounts for unwanted two qubit coupling during single qubit gates by reducing delay times during which coupling is desired. Specifically, every instance of $R_1(1)U_{ZZ}(t)R_2(2)$ is replaced with $R_1(1)U_{ZZ}(t)R_2(2)$, where $t = \frac{t}{2}(1 + \epsilon)$. This technique was critical in the successful implementation of Shor's algorithm with NMR [29]; here, it is used with care, since many Hamiltonians have the same ϵ as the pairing Hamiltonian of interest, and it is possible to tune ϵ to get the right ϵ for the wrong reasons.

Numerical simulations comparing W1 and W2 show that the effect of such control errors on the WBL algorithm is a shift in the estimated gap value from the expected value. This shift can be quite significant, as shown in Fig. 2, and indeed can dominate errors due to other imperfections, such as the Fourier transform. Compensating for unwanted scalar couplings in NMR implementations of quantum simulations is thus vital for obtaining correct results; implementations with other physical systems will similarly have to deal with faulty controls.

Finally, we consider the third question: saturation of the predicted precision bounds with an NMR implementation of the WBL algorithm. The WBL algorithm is parameterized by the number of qubits, n , the simulated time step t_0 , the number of steps Q , the degree of Trotterization k , the adiabatic time step t_{ad} , and the number of adiabatic steps S . An NMR system is characterized by a characteristic decoherence time τ , and following the discussion above, it is convenient to work with a small number of qubits for times shorter than 3τ , such that quantum error correction is unnecessary. We implement

two specific instances of the pairing Hamiltonian H_{pair} involving three modes ($n = 3$) and two pairs, leading to a 3-dimensional Hilbert space spanned by $|j_0 1 i\rangle$, $|j_1 0 i\rangle$, and $|j_1 1 i\rangle$. The simulation is started the ground state of the two spin up subspace of H_0 , $|j_1 1 i\rangle$, prepared using temporal labeling [30].

WBL estimated the expected size of the system that could be simulated without error correction by choosing $k=t_0 = 0.1$ and $F_T = 1$. They found that the number of gates required scales as $3n^4 = F_T$, including the necessary decoupling pulses. The gate time, t_g is assumed equal to 10^{-5} and for up to $n = 10$, can be found to precision 10^{-3} . Here, we need to precision 10^{-4} , and the number of qubits is $n = 3$, consistent with the WBL bound $n \leq 4$ for these parameter choices.

The first stage of the WBL algorithm is to quasidiabatically evolve into the ground state $|j_1 i\rangle$ of H_{pair} , with discrete changes in the simulated Hamiltonian, using a procedure previously demonstrated [31]. The Hamiltonian used at each discrete step s is $H_{ad}(s) = (1 - s/S)H_0 + (s/S)H_{\text{pair}}$, where S is the maximum number of steps. Unitary evolution U_{ad} at each step for time t_{ad} is then approximated using the above pulse sequences, as $U_{ad} = \prod_{s=0}^S V_{ad}(s; t_{ad})$. Preparation of the state $|j_1 i\rangle$ requires evolving at a rate faster than that for adiabatic evolution, thereby exciting the state $|j_1 i\rangle$. This quasidiabatic evolution is accomplished by reducing S or t_{ad} compared to the adiabatic case [1]. Higher-energy states will also be excited, but S and t_{ad} can be adjusted to minimize this. Quasidiabatic evolution in this experiment was attained with $S = 4$ steps and $t_{ad} = 1/700$ s. Note that for $kH_{XX} + H_{YY} \rightarrow kH_{0k}$ there can be a phase transition as s is changed [32]; as the gap goes to zero at the phase transition this can be problematic, since the number of steps required for successful quasidiabatic evolution grows inversely with the gap.

The second stage of the algorithm is evolution of the state $|j_1 i\rangle$ under the pairing Hamiltonian for Q time steps of duration t_0 with $k = 2$. These parameters are chosen such that $1/(Q t_0) = 100$, $Q t_0 < \tau$, and $k=t_0 > 0.1$. Note that many Q and t_0 yield the same F_T ; this is used to our advantage below.

We performed our experiments using a 500 MHz Varian UNITY INOVA spectrometer and ^{13}C -labeled CHFB r_2 , with coupling strengths $J_{HC} = 224$ Hz, $J_{HF} = 50$ Hz, and $J_{CF} = 311$ Hz. The two pairing Hamiltonians simulated were H_1 , the "natural" Hamiltonian, in which $V_{12} = J_{HC}$, $V_{13} = J_{HF}$, and $V_{23} = J_{CF}$, and a harder case, an artificially constructed Hamiltonian H_2 , in which $V_{ab} = J_{HC}$ and $V_{ac} = V_{bc} = 0$. For both Hamiltonians, $\nu_1 = 150$ Hz, $\nu_2 = 100$ Hz, and $\nu_3 = 50$ Hz. Implementation of H_2 required an additional decoupling pulse compared to H_1 , following the circuit in Fig. 1.

Each Hamiltonian was simulated for times t_0 to $Q t_0$, using the W1 and W2 methods, and an NMR spectrum

Model	ω /Hz	Method	$\omega_{\text{exp}}/2$ Hz	τ_e /ms	t_0 /ms	Q
H_1	218.2	W1	227.2	180	1	400
		W2	220.2	250	2	200
H_2	452.2	W1	554.10	30	.5	200
		W2	440.5	80	.5	200

TABLE I: Experimental results for gaps found for Hamiltonians H_1 and H_2 . Estimated gaps (ω_{exp}) and effective coherence times (τ_e) for given time steps t_0 and number of steps Q are obtained by least-squares fitting of the time-dependent NMR peaks to an exponentially-decaying sinusoid.

was acquired for each time duration. A classical discrete FT of the NMR peak intensities of one spin (hydrogen) over the t_k yielded four spectra of H_{pair} . The experimental result ω_{exp} was determined by a least-squares fit of the highest-signal NMR peak to a damped sinusoidal function with frequency ω and decay rate τ_e .

Ideally, the result should find $\omega = 218.2$ Hz for H_1 and $\omega = 452.2$ Hz for H_2 , as determined by direct diagonalization. Note that for H_2 , ω is the energy difference between $|E_1\rangle$ and $|E_2\rangle$, since $|E_1\rangle$ is not connected by usual adiabatic evolution; the larger gap requires that $t_0 = k$ be smaller when simulating H_2 . For the experimental result, we expect that $\omega_{\text{exp}} = \omega + \delta\omega_{\text{sys}} + \delta\omega_{\text{FT}}$ where $\delta\omega_{\text{sys}}$ is an offset due to Trotterization and/or faulty controls. Q and t_0 determine the theoretical bound on the precision, in the absence of control errors, and the experiment should saturate this bound when the ω_{exp} is ω_{FT} from the actual value.

Experimental results for the spectra of H_2 are shown in Fig. 2; experimental parameters Q and t_0 and numerical results from the analysis for ω_{exp} and τ_e for each experiment are summarized in Table I.

The impact of systematic and random errors was investigated by simulating H_1 with W1 (no control error compensation) for $\omega_{\text{FT}} = 2.5 \times 2$ Hz at two different simulation times, $t_0 = 1$ ms and $t_0 = 2$ ms. As expected, the random error for both cases is ω_{FT} . Note that the systematic error increases with smaller t_0 . This signals that the error due to unwanted scalar coupling becomes larger than the errors due to the Trotter approximation. Consequently, a slightly longer t_0 yields a systematic error that is within ω_{FT} of the exact answer, saturating the predicted theoretical bounds on precision.

Convergence to the correct result is another important issue for all discrete time simulations. For this 3-qubit system, we performed a detailed numerical simulation to determine that $t_0 = 2$ was optimal. For a large system this is no longer possible, and convergence tests would need to be used to verify the answer. The procedure would reduce t_0 (or increase k) until the change in ω_{exp} was smaller than the desired precision.

While the results for Hamiltonian H_1 were good even

without control error compensation, the effects of control errors were very evident in the results for H_2 . Hamiltonian H_2 was implemented with W1 (no error compensation) and W2 (simple error compensation) for $\omega_{\text{FT}} = 10 \times 2$ Hz and $t_0 = 0.5$ ms. The shorter time step was necessary because the larger ω made the simulation more sensitive to Trotter errors. Comparing the W2 and W1 results shows that with no control error compensation, a gap is found that is $\omega_{\text{FT}} = 5$ away from the actual value. In contrast, with simple error compensation ω_{exp} is ω_{FT} from the actual value, saturating the theoretical bound. Future implementations should certainly strive to detect and bound control errors; this could be done by verifying that ω_{exp} scales as t_0^3 for small values of t_0 , as theoretically expected.

In conclusion, we have studied the theoretical and empirical bounds on the precision of results obtained with quantum simulations, in the context of the pairing Hamiltonian algorithm proposed by Wu, Byrd, and Lidar. We have implemented the smallest problem instance that requires quasiadiabatic evolution, verifying that the algorithm computes the gap to within the precision of the method. We also find, however, that simulations of this type are particularly sensitive to systematic errors in the applied Hamiltonian and that fault-tolerant implementations are inefficient with respect to precision using current Trotter approximation methods.

Nevertheless, in practice, when only limited precision is desired and for a sufficiently large system, quantum simulations may still outperform classical numerical simulation, as demonstrated for molecular energies [33]. A voiding the cost of precision is desirable, and can be done by designing quantum simulations to explore questions that are insensitive to the microscopic details of the Hamiltonian [34]. How to develop quantum simulations for faulty small scale (10-20 qubit) quantum computers that can outperform classical computations remains an open question.

-
- [1] L. A. Wu, M. S. Byrd, and D. A. Lidar, Phys. Rev. Lett. 89, 057904 (2002).
 - [2] R. P. Feynman, Int. J. Theor. Phys. 21, 467 (1982).
 - [3] S. Lloyd, Science 273, 1073 (1996).
 - [4] D. S. Abrams and S. Lloyd, Phys. Rev. Lett. 83, 5162 (1999).
 - [5] P. W. Shor, SIAM J. on Comp. 26, 1484 (1997).
 - [6] L. K. Grover, Phys. Rev. Lett. 79, 325 (1997).
 - [7] S. Somaroo, C. H. T seng, T. F. Havel, R. Laamme, and D. G. Cory, Phys. Rev. Lett. 82, 5381 (1999).
 - [8] D. Cory, R. Laamme, E. Knill, L. Viola, T. Havel, N. Boulant, G. Boutis, E. Fortunato, S. Lloyd, R. Martinez, et al., Fortschr. Phys. 48, 875 (2000).
 - [9] X. Peng, J. De, and D. Suter, Phys. Rev. A 71, 012307 (2004).
 - [10] C. Negrevergne, R. Somma, G. Ortiz, E. Knill, and

- R. Laamme, Phys. Rev. A 71, 032344 (2004).
- [11] Z. Chen, J. Yezpez, and D. G. Cory (2004), the LANL preprint quant-ph/0410198.
- [12] X.-D. Yang, A. M. Wang, F. Xu, and J. Du (2004), the LANL preprint quant-ph/0410143.
- [13] E. Jane, G. Vidal, W. Dur, P. Zoller, and J. Cirac, Quant. Inf. and Comp. 3, 15 (2003).
- [14] M. Greiner, O. Mandel, T. Esslinger, T. W. Hansch, and I. Bloch, Nature 415, 39 (2005).
- [15] R. Cleve, A. Ekert, C. Macchiavello and M. Mosca, Proc. Roy. Soc. London Ser. A 454, 339 (1998).
- [16] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge University Press, Cambridge, UK, 2000).
- [17] J. Ketterson and S. Song, Superconductivity (Cambridge University Press, Cambridge, United Kingdom, 1999).
- [18] G. D. Mahan, Many-Particle Physics (Kluwer, New York, 2000), 3rd ed.
- [19] P. Ring and P. Schuck, The nuclear many-body problem (Springer, New York, 1980).
- [20] M. Suzuki, Phys. Lett. A 165, 387 (1992).
- [21] J. Preskill, Proc. Roy. Soc. London Ser. A 454, 385 (1998).
- [22] D. Gottesman, Phys. Rev. A 57, 127 (1997).
- [23] D. Gottesman and I. L. Chuang, Nature 402, 390 (1999).
- [24] A. Y. Kitaev, A. H. Shen, and M. N. Vyalii, Classical and Quantum Computation, vol. 47 of Graduate Studies in Mathematics (American Mathematical Society, Providence, 2002).
- [25] H. Trotter, Pacific Journal of Math 8, 887 (1958).
- [26] D. J. Wineland, C. Monroe, W. M. Itano, D. Leibfried, B. E. King, and D. M. Meekhof, J. of Res. of the National Inst. of Standards and Technology 103, 259 (1998), <http://nvl.nist.gov/pub/nistpubs/jres/103/3/cnt103-3.htm>.
- [27] L. M. K. Vandersypen and I. L. Chuang, Rev. Mod. Phys. 76, 1037 (2004).
- [28] N. Khaneja, S. J. Glaser, and R. Brockett, Physical Review A 65:032301 (2002).
- [29] L. Vandersypen, M. Steen, G. Breyta, C. Yannoni, R. Cleve, and I. Chuang, Nature 76, 646 (2000).
- [30] L. M. K. Vandersypen, M. Steen, M. H. Sherwood, C. S. Yannoni, G. Breyta, and I. L. Chuang, Appl. Phys. Lett. 76, 646 (1999).
- [31] M. Steen, W. van Dam, T. Hogg, G. Breyta, and I. Chuang, Phys. Rev. Lett. 90, 067903 (2003).
- [32] R. H. McKenzie, Phys. Rev. Lett. 77, 4804 (1996).
- [33] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, Science 309, 1704 (2005).
- [34] J. Barjaktarevic, G. M. Ilum, and R. H. McKenzie, Phys. Rev. A 71, 012335 (2005).