nam ics.

Decoherence induced by a phase-dam ping reservoir

Renato M. Angelo

Departamento de F sica, Universidade Federal do Parana, Caixa Postal 19044, CEP 81531-990, Curitiba, Parana, Brazil

E.S.Cardoso and K.Furuya

Instituto de F sica 'G leb W ataghin', U niversidade E stadual de Campinas, Caixa Postal 6165, CEP 13083-970, Campinas, Sao Paulo, Brazil

A phase dam ping reservoir com posed by N -bosons coupled to a system of interest through a cross-K err interaction is proposed and its e ects on quantum superpositions are investigated. By m eans of analytical calculations we show that: i-) the reservoir induces a G aussian decay of quantum coherences, and ii-) the inherent incom m ensurate character of the spectral distribution yields irreversibility. A state-independent decoherence tim e and a master equation are both derived analytically. These results, which have been extended for the therm odynam ic lim it, show that nondissipative decoherence can be suitably contemplated within the EID approach. Finally, it is shown that the sam e mechanism yielding decoherence are also responsible for inducing dynam ical disentanglem ent.

PACS num bers: 03.65.Y z, 03.67 M n, 03.65.Ta

I. IN TRODUCTION

Environment-induced decoherence (ED) has been widely accepted as the mechanism responsible for the de-

struction of quantum superpositions. A great am ount of

research in the last two decades has put in evidence -

theoretically [1] and experimentally [2] - the potentiality

of this e ect in explaining challenging issues in quantum mechanics, as for instance the nonexistence of macroscopic superpositions and the measurement process. The

EID approach is based on the coupling of a system S with a many degrees of freedom environment, which is theoretically described in terms of microscopic quantum particles in equilibrium at temperature T. The coupling with the reservoir forces the system to evolve through a

nonunitary, irreversible and, in general, dissipative dy-

O ne of the conceptual di culties of the E ID program, as it has been pointed out by som e authors, concerns the irreversible energy transfer from the system to the reserknown phase-destroying master equation [8, 9]:

$$\frac{\partial b}{\partial t} = \frac{[H^{0};b]}{\{\sim \quad \sim^{2} [H^{0}; [H^{0};b]];}$$
(1)

being a speci c parameter in each theory. By means of straightforward calculations in the eigenbasis f E_n ig of \dot{P} one may prove that the above master equation admits the following solution:

$$b(t) = \sum_{\substack{n,n^{\circ} \in 0 \\ n;n^{\circ} = 0}}^{X^{i}} \frac{(t) (E_{n} - E_{n}^{\circ})^{t}}{(E_{n} - E_{n}^{\circ})^{2}} \quad f_{n} ih E_{n^{\circ}} ; (2)$$

where $nn^{0} = hE_{n} \frac{1}{2}(0) \pm \frac{1}{n}i$ and $_=$. The purity, P = Tr[²], then reads

$$P(t) = \int_{n = 0}^{X^{1}} j_{n n^{0}} j^{2} \exp \left(\frac{2(t) (E_{n} - E_{n}^{0})^{2}}{2} \right)$$
(3)

W hen is constant, for example, one can observe an exponential decay with time for the o -diagonal term s of b, in plying in a decrease of the purity (P(0) > P(1)), even though the energy of the system remain unchanged, i.e., Tr[P(t)] = Tr[P(b(0)]. Clearly, the master equation (1) suitably describes nondissipative phase decoherence. Recently, how ever, the relevance of SID as a physically m eaningful process has been questioned [10].

In this work we show that it is also possible to obtain phase decoherence within the EID scenario, without any additional assumption to the standard quantum form alism if we consider a diagonal interaction. Our approach is based on a many-boson model in which the interaction between the components is given by the wellknown cross-Kerr coupling $\mathbf{b}_{j}^{i}\mathbf{b}_{j}\mathbf{b}_{j}^{j}\mathbf{b}_{j}$. This assumption is grounded in the fact that this kind of interaction is

voir, which prohibits the application of this approach to conservative system s. Several alternative m ethods have been proposed to settle this problem . In general, these approaches defend the so term ed self-induced decoherence (SID), whose cause is not attributed to external degrees of freedom (environm ent). Am ong them one m ay quote, for instance, proposals of a generalized Schrodinger equation [3], assumptions about uctuations of classical param eters or internal degrees of freedom yielding decoherence [4], and nondissipative decoherence as a result of the quantum state averaging over a therm all distribution of velocities [5]. A lso, references [6, 7] contribute to this list.

A recurrent result in the SID program is the reduction of the system dynamics to the following form of the wellphysically conceivable, i.e., two hybrid bosonic degrees of freedom such as ionic vibrationalm ode and the quantized eld m ay interact e ectively in this way as has been dem onstrated recently [11]. Besides, several inform ation transfer processes have been developed based on cross-Kerr interactions [12] in such hybrid systems, and several sources of decoherence without transition between the states of di erent quantum num bers - thus destroying coherence without causing energy dissipation - have also been discussed in [13]. These facts lead us to think that decoherence under this kind of phase channel will be a real problem to be considered. One can also do engineering of a bath of N identical ion traps [14] each one coupled to the same eld, such that they satisfy the following conditions of Ref. [11]: (i) the detuning of the atom with the eld is di erent from them ultiples of the frequency of the vibrationalm ode (\in k, k integer) thus avoiding to excite the vibrational degree of freedom of each ion; (ii) in the large detuning limit (g ;q being the ion-eld coupling) in order to achieve a dispersive regime; and (iii) if the non-Lamb-Dicke regime is desirable. A lthough usually the energy dissipating sources of decoherence m ay also be present (via typical oscillator-oscillator position coupling), in the above engineered bath case, a nondissipative kind of decoherence is expected. By means of analytical calculations we show that the model indeed produces nondissipative decoherence, transform ing quantum superpositions into m ixed states through a Gaussian decay.

In addition, we show that bipartite entanglement can be dynamically destroyed by such a reservoir. Dodd and Halliwell showed how open system dynamics destroys entanglement by means of the same mechanism that destroys interference [15]. Here, analytical formulas allow us to make the same identication rigorously.

This paper is outlined as follows. In Sec. II the reservoir is modeled as a system of N harmonic oscillators in a therm al equilibrium and the coupling with the system of interest is assumed to be given e ectively by a cross-K err interaction. Exact results dem onstrating the capability of the model in yielding decoherence for arbitrary quantum states are presented in Sec.III, where: (A) analytical results dem onstrating the G aussian decay of quantum coherences and an estim ate for the decoherence time are obtained for short times; (B) the equilibrium regime is analyzed; (C) the therm odynam ic lim it is im plem ented; (D) two exam ples of decohering dynam ics for some special quantum states, such as the cat state and a superposition of Fock states, are given, and (E) a phase-destroying master equation is derived in the regime of weak coupling. In Sec.IV we dem onstrate the dynam ical disentanglem ent induced by the same mechanisms responsible for phase decoherence. Finally, in Sec.V we relate our concluding rem arks.

II. THE MODEL

In ref.[11], Sem iso and Barranco have derived an effective H am iltonian for a model of two-level trapped ion coupled to the quantized eld inside a cavity. Under certain conditions, which allow for the rotating wave approxim ation and assumptions on both a large detuning between the ion and the eld (dispersive lim it) and a sm all value for the Lam b-D icke parameter, the authors nd the follow ing e ective H am iltonian

$$\mathbf{H}_{eff}^{p} = \mathbf{h}_{1}^{y} \mathbf{h}_{1} + \mathbf{h}_{2}^{y} \mathbf{h}_{2} + \mathbf{h}_{1}^{y} \mathbf{h}_{1} \mathbf{h}_{2}^{y} \mathbf{h}_{2} : \qquad (4)$$

This K err-type coupling produces only phase interaction, not being able to introduce energy exchanges between the bosons. For this reason, this kind of phase coupling turns out to be a potential candidate for modeling nondissipative reservoirs.

Having been inspired by (4), we consider the following many-bosons model $% \left(\frac{1}{2} \right) = 0$

$$\mathbf{P} = \sum_{\substack{i=0\\j=0}}^{X^{N}} \sim !_{i}\mathbf{b}_{i} + \sum_{\substack{i,j=0\\i>j}}^{X^{N}} \sim g_{ij}\mathbf{b}_{i}\mathbf{b}_{j};$$
(5)

where \mathbf{b}_i denotes the number operator of the i-th boson. The coupling is clearly incapable to produce energy exchanges between any two degrees of freedom.

Now we focus on the dynam ics of a single boson, say the one with index 0", in order to investigate the features of such an environm ent. For simplicity, we also assume that there is no interaction between the reservoir components. The Ham iltonian (5) is then replaced by

$$\dot{\mathbf{H}} = \dot{\mathbf{H}}_{\mathrm{S}} + \dot{\mathbf{H}}_{\mathrm{R}} + \dot{\mathbf{H}}_{\mathrm{I}}; \qquad (6)$$

where

$$\mathbf{P}_{\mathrm{S}} = \sim \mathbf{P}_{\mathrm{O}} \mathbf{b}_{\mathrm{O}}; \qquad (7a)$$

$$\mathbf{b}_{\mathrm{R}} = \sum_{k=1}^{X^{\mathrm{N}}} \sim \mathbf{b}_{k}; \qquad (7b)$$

$$\dot{P}_{I} = \sum_{k=1}^{X^{N}} \sim g_{0k} b_{0} b_{k}$$
: (7c)

Initially, the joint quantum state is assumed to be

$$b(0) = b_{S}(0) - \frac{e_{h}^{\dot{P}_{R}}}{Tr e^{\dot{P}_{R}}};$$
 (8)

which indicates that all oscillators of the reservoir are in thermal equilibrium at a temperature (k_B) 1 . The system of interest is initially in an arbitrary pure state b_S (0).

As it has been shown by Ford et al [16], the most general quantum Langevin equation, describing macroscopically the dissipative dynamics of a quantum system, can be realized with an independent oscillator (IO) model. A lso, these authors have dem onstrated that several traditionalm odels of reservoirs inducing dissipative decoherence can be mapped on an IO model, which, accordingly, m ay be regarded as a fundam ental m odel for the E ${\rm I\!D}$ program .

Our reservoir, however, may be regarded as being grounded on a type of generalized IO model which im plem ents nonlinear couplings. In fact, the originalm odel for the ions, from which the e ective cross-K err coupling given in Eq.(4) has been derived in ref.[11], contains the fundam ental ingredients of an IO model, including energy exchanges.

III. PHASE DECOHERENCE

The rather simple form of the coupling allows one to solve analytically the dynam ics by means of straightforward calculations, with no prior need for weak coupling or specic statistical assumptions. The exact result for the system density matrix in the \mathbb{P}_0 eigenbasis is written as

$$b_{\rm S} (t) = \sum_{\substack{m \ \text{,} m \ ^0 = 0}}^{X^{\rm I}} {}^{\rm M} C_{m \ \text{,} m \ ^0} (t) e^{\{(m \ m \ ^0) \ ! \ _0 t} jn \ ihm \ ^0 j; (9)$$

where $b_s = Tr_R [b]_{n m m^0} = hm jb_s (0) jm^0 i$ and

$$C_{m,m} \circ (t) = \frac{\dot{Y}^{i}}{\sum_{k=1}^{k=1}} \frac{1 e^{-i k}}{1 e^{-i k} e^{i (m m - 0) g_{0k} t}} : (10)$$

The purity of the system state, $P_s = Tr_s [b_s^2]$, is given by

$$P_{S}(t) = \int_{m \ m}^{X^{2}} j_{m \ m} \circ j^{2} j_{m \ m} \circ (t) j^{2}; \qquad (11)$$

with

$$jc_{m,m}\circ(t)j^{2} = \frac{\gamma^{N}}{\sum_{k=1}^{k} 1 + \frac{\sin^{2}\frac{(m-m^{0})g_{0k}t}{2} = 1}{\sinh^{2}\frac{-\frac{1}{2}t}{2}}; \quad (12)$$

Notice in Eq.(11) that all the time and temperature dependences com e from the function $C_{m,m}$. Also, in (9) we see that C_m , o is the only function depending on the coupling param eters between the system and the reservoir. Thus, it is natural to expect that this function carries all the dynam ical inform ations about the correlations of the system with the environment and, therefore, about the decoherence of the quantum state.

The analytical results (9)-(12) provide a rather interesting explanation for the well-known requirement of a spectral distribution for the coupling parameters q_{0k} . For resonant ($g_{0k} = g$) or com m ensurate ($g_{0k} = \frac{1}{k}g$, $\frac{1}{k}$ integer) couplings, the time dependence in Eq.(10) behaves as e ^{Mgt}, with M integer. Then, for all instants satisfying $t_n = 2 \text{ n=g}$, with n 2 N, one has $C_{m,m^0} = 1$,

3

for all m and m^0 , indicating that the quantum state has just recovered its initial condition of purity, namely P_{S} (t_n) = P_{S} (0) = 1. This certainly is an unexpected recurrence for a supposedly irreversible reservoir. How ever, far apart from this very especial conditions, incommensurate values of coupling inherent to any physically acceptable spectral distribution will set irreversibility just by avoiding simultaneous recurrences in C_{m m}. This is indeed ensured by the product in (10). These statem ents are dem onstrated in Fig.1, where a typical behavior for $\mathfrak{T}_{m,m} \circ \mathfrak{f}$ is num erically simulated. Notice that all ar-




FIG.1: Log-linear plot for the function $\mathcal{C}_{m,m} \circ \mathcal{J}$ as a function of $(m m^{0})$ tfor N = 100 and ~! = 1. Two cases of couplings were simulated, namely: (a) a resonant spectrum, $g_{0k} = I N$, and (b) a G aussian spectral distribution, $g_{0k} = I N$ = N exp $\frac{(k k_0)^2}{2 N^2}$. In (a) = 0:1 and in (b) = $g_{0k}^2 = 0.01$ (see Eq.(14)) and 0:1125. In both cases, G^2 $k_0 = 50$. Notice the long-time recurrences in (a).

qum ents given in this section rem ain valid even for the simplest case of resonant harm onic frequencies $(!_k = !)$. Since this assumption does not change signi cantly the m ain features of the m odel, it will be considered hereafter for sim plicity.

A. The short-time regime

In Fig.(1) one may clearly realize the presence of a short-time regime in which the collapse of the characteristic function occurs. Actually, in this section we shall show analytically that this is the very time scale of decoherence.

Let us consider the regime of short times, namely (m m 0) g_{0k} t 1. In this lim it Eq.(12) can be replaced with

$$\mathcal{T}_{m,m} \circ (t) \mathcal{J}^{2} = e^{(m,m,0)^{2} t^{2} = \mathcal{J}^{2}}; \qquad (13)$$

where the decoherence time $_{\rm D}$ is given by

$$_{\rm D} = 2 \,{\rm G}^{-1} \, \sinh \left(\sim ! = 2 \right) :$$
 (14)

Note, by Eq.(13), that the decoherence time has been de ned regardless of the di erence m $\,$ m 0, even though the instant of decay is strongly dependent on this di erence. This choice attributes to $_{\rm D}$ the role of an upper bound for surviving coherences, i.e., it corresponds to the time scale beyond which decoherence has indeed a ected all elements of the matrix $b_{\rm S}$.

A nother in portant characteristic of our model is the G aussian decay given by Eq.(13), which has already been found in other approaches [4, 5]. This behavior implies in a sudden decrease in the purity of the (arbitrary) quantum state for times longer than $_{\rm D}$. In fact, well beyond this time scale the quantum state will have reached its equilibrium con guration. In this regime, since C_{m m} $_{\circ}$ (t) = $j_{\rm Cm m} _{\rm m} _{\circ}$ (t) $j_{\rm e}^{i}$ (t), one may write

$$C_{m,m} \circ (t > D) = m,m \circ; \qquad (15)$$

indicating that the dynam ics will transform

$$b_{S}(0) = \int_{m \ m^{0}=0}^{X^{1}} ihm^{0} jn \ ihm^{0} jr \qquad (16)$$

with

into

$$b_{\rm S} (t > {}_{\rm D}) = \sum_{\substack{m \neq m \\ m = 0}}^{\chi^2} j_{\rm m} j_{\rm m} j_{\rm m} j_{\rm m}$$
(18)

with

P (t > D) =
$$\begin{array}{c} X \\ m_{\pi} m \end{array}$$
 (19)

Note that the structure of the coupling given by Eq.(7c) autom atically determ ines the pointer basis, i.e., the basis in which the density m atrix becom es diagonal. This allow sus to conjecture that this reservoir will fatally destroy the entanglem ent for a state like jnijn $i + j^0 i j n^0 i$ (this will be dem onstrated in Sec.IV).

A nother rem arkable result derives from equations (17) and (19): P (0) is always greater than P (t > $_{\rm D}$), indicating the occurrence of a base-independent decoherence. Furtherm ore, since no assumption was made about the pure initial state, the decohering process is also state-independent, $_{\rm D}$ being the general upper bound for the time scale of such a process.

This generality associated to $_{\rm D}$ attests that this time scale is in fact an intrinsic feature of the reservoir model and not of the particular quantum state of the system . This is perhaps the most im portant di erence regarding the usual tim e scales found in literature, which in general exhibit a dependence with dⁿ, d being the distance in the conguration space separating the two components of the particular superposition state under investigation (for exam ples of such a state-dependent tim e scales see references [1, 2], for the free particle problem, and [17, 18], for oscillators m odels). In our m odel this dependence m ight be introduced by incorporating the \distance" m m° in the de nition given by Eq.(14). However, we have opted to attribute a general character to $_{\rm D}$. Concerning the dependence with other physical parameters, such as characteristic frequencies, tem perature and ~, our tim e scale agrees qualitatively with those of the references quoted above. For instance, all time scales predict a decrease in the decoherence time as the temperature increases. A precise quantitative agreem ent is not expected in principle since we are working, in contrast to those approaches, with nonlinear interactions.

B. The long-time regime

Now we analyze the equilibrium situation (t ! 1). In this case, a direct inspection of Eq.(12) allows us to calculate a lower bound, which for any m and m⁰, reads

$$f_{m m}^{2} \circ f_{LB}^{2} = \frac{\sinh^{2}(\sim !=2)}{1+\sinh^{2}(\sim !=2)}^{N} : \quad (20)$$

It is easy to see, for N large but nite, that the lower bound given by Eq.(20) can range from arbitrarily small values, in the regime of high temperatures (~! 1), to values arbitrarily close to the unit, in the regime of low temperatures (~! 1). The latter regime, for which $\mathfrak{f}_{m\,m^0}\mathfrak{f}_{LB}$! 1, corresponds to the one in which the dynamics is practically unitary, without decoherence. On the other hand, the former, for which $\mathfrak{f}_{m\,m^0}\mathfrak{f}_{LB}$! 0, corresponds to the regime in which the ext of decoherence are accentuated. In Fig.2 these regimes can be identied clearly.

The abrupt transition between these two limits (central region of Fig 2) corresponds to a regime in which decoherences still occurs, but only partially, since a nite value for $\mathfrak{f}_{m,m} \circ \mathfrak{f}_{B}$ indicates the existence of surviving o -diagonal terms in the quantum state. The critical tem perature, i.e., the one beyond which the e ects of decoherence become accentuated, may be estimated by imposing that $\mathfrak{f}_{m,m} \circ \mathfrak{f}_{B} = 1=2$ in Eq.(20). The result is a function of N :

$$_{\rm crit}$$
 (N) = $\frac{-!}{2}$ arcsinh $\frac{1}{2^{1-N}}$: (21)

For tem peratures in the vicinity of $(k_B _{crit})^{-1}$ the quantum state must contain o -diagonal terms contributing




FIG.2: Lowerbound C_{LB}^{2} as a function of the dimensionless parameter ~! for N = 10^{2} , 10^{3} , and 10^{4} (from left to right). The critical values $_{\rm crit}$ ~!, estimated from Eq.(21) for each N, are given by the vertical dashed lines.

to an increase in the value of the purity in the regime of long times.

C. The therm odynam ic lim it

In the previous sections we have discussed the general characteristic of our cross-K err reservoir for an arbitrary value of N . Now, we investigate the therm odynam ic lim it, which better describes the physical reality of a reservoir.

R igorously, the therm odynam ic lim it is in plemented by taking N ! 1 and V ! 1, with (N=V) ! 1. Before considering this lim it in ourmodel, however, it would be necessary to replace the coupling g_{0k} by its original physical form $g_{0k} = V$, which derives from the process of quantization of the electromagnetic eld [19]. W ith this correction, the frequency G de ned in Eq.(14) would assume the form $G = \frac{p_{0k}}{k} \frac{q_{0k}^2 = V}{N + k}$. Then, in the resonance $g_{0k} = r$, for instance, we would have $G = \frac{p_{0k} = V}{N + k}$, which possesses a well de ned thermodynam ic lim it

This simple example suggests that we can implement the therm odynamic limit in our formulas just by requiring that the coupling g_{0k} be quadratically normalized, i.e., that $G^2 = \int_k g_{0k}^2$ be nite. For instance, for the resonant case mentioned above, the assumption that $g_{0k} = = N$ will automatically set G =. In a less trivial case, in which the coupling is given by a G aussian spectral distribution, the therm odynamic limit can be ensured just by assuming that

$$g_{0k} = p_{\frac{1}{N}} \exp \left(\frac{(k - k_0)^2}{2N^2} \right)$$
 (22)

In this case, one m ay verify num erically that the relation G $\,<\,$ is always obeyed for any N . For instance, for

 $k_0=50,$ one may verify that G $\,!\,$ 0:706 $\,$ as N $\,!\,$ 1 . A lso in this case, it may be veried that G 6 0:889 , this upper bound occurring for N $\,=\,2k_0=\,100.$

W ith these considerations we can reinterpret the results given by E sq.(13)-(21) within the fram ework of a reasonable therm odynam ic reservoir possessing the following characteristics:

- i-) Gaussian decay for the short-tim e regim e;
- ii-) Decoherence time given by $_{D} = 2G^{1}$ sinh (~!);
- iii-) Characteristic frequency given by $G^2 = \prod_{k=1}^{P} g_{0k}^2$;
- iv-) Irreversibility induced by a quadratically norm alizable spectral distribution (g_{0k});
- v-) Lowerbound jC $f_{LB} = 0$ and $T_{crit} = 0$.

Item v-), which can be directly obtained by taking N !1 in Eq.(20) and Eq.(21), tell us that decoherence will occur in the therm odynam ic lim it even for arbitrarily low tem peratures.

D. Exam ples

Any initially pure state for the system of interest may be expanded as

$$y_{0}^{i} = p(n) j_{1};$$
 (23)

where p(n)f is the photon distribution de ning the quantum state. The corresponding density operator is then written as $b_s(0) = j_0$ in $_0$ j and its matrix elements in Fock basis reads $m_{\pi} \circ = p(m)p(m^0)$. Thus, Eq.(11) becomes

$$P_{S}(t) = \sum_{m = 0 m}^{X^{1}} \sum_{n = 0}^{X^{1}} \dot{p}(m) \dot{j} \dot{p}(m) \dot{j} \dot{j} m^{0} \dot{j} \dot{j} m_{\pi} \circ (t) \dot{j} : (24)$$

Now we apply our model to investigate num erically the decoherence process for: (i) the cat state

$$j_{0}i = \frac{j_{1}i + j_{1}i}{2 + e^{2j_{1}j^{2}}}$$
(25)

and (iii) the superposition of Fock states

$$j_{0}i = \frac{jn_{1}i + jn_{2}i}{p_{\overline{2}}};$$
 (26)

with $m_1 \notin m_2$. For this Fock superposition, for which an analytical calculation yields $P_S(t) = 1 + jC_{m_1,m_2}(t)j^2 = 2$, it is easy to verify the classicalization just by applying formulas (16) and (18). Indeed, the equilibrium value for the purity, $P_S(1) = 1=2$, is directly associated to the statistical mixture $b_S(1) =$ (jn 1 ilm 1 j+ jn 2 ilm 2 j)=2. For the cat state, on the other hand, at the equilibrium time we have a mixed state with $0 < P_S(1) < 1$. In Fig.3 we show numerical simulations for the purity of these states. The decoherence time is also plotted, attesting both the validity and universality of the estimate (14). Notice that for ' 10⁶ s¹ we have p' 10⁸ s.




FIG .3: Log-linear plot for the purity of (i) a cat state (dashed curve), with = 2:0, and (ii) the superposition $(jli+j2i)=\frac{1}{2}$ (solid curve). For these simulations, N = 100 and ~! = 0:01. Here we have used the spectral distribution employed in Fig.1-(b). The vertical line stands for the decoherence time $_{\rm D}$ = 1:13 10⁻², estimated from Eq.(14).

E. Master Equation

The interaction H am iltonian given by Eq.(7c) can be written as

$$H_{I}^{p} = \Psi_{S} \quad \Psi_{R}; \qquad (27)$$

where we have de ned: $\oint_S = \mathbf{b}_0$ and $\oint_R = \int_k g_{0k} \mathbf{b}_k \cdot \mathbf{B} \mathbf{y}$ m eans of relatively simple calculations we then deduce a master equation for our model in the weak coupling approximation. The result, in the interaction picture, reads

$$\frac{\mathfrak{G}\mathbf{b}_{\mathrm{S}}^{\mathrm{I}}}{\mathfrak{G}\mathbf{t}} = \frac{\overset{\mathrm{h}}{\mathfrak{P}_{\mathrm{S}}}; \overset{\mathrm{I}}{\mathfrak{b}_{\mathrm{S}}^{\mathrm{I}}}}{\{\sim} \quad \frac{\mathsf{t}}{\sim^{2}} \overset{\mathrm{h}}{\mathfrak{P}_{\mathrm{S}}}; \overset{\mathrm{h}}{\mathfrak{P}_{\mathrm{S}}}; \overset{\mathrm{ii}}{\mathfrak{b}_{\mathrm{S}}^{\mathrm{I}}}; \qquad (28)$$

where $\vartheta_{\rm S}$ $\vartheta_{\rm R}^{n}$ $i\vartheta_{\rm S}^{n}$ and ${}_{\rm h}$ $\vartheta_{\rm R}^{2} i=\vartheta_{\rm R}^{0}$ i^{2} . The reservoir expectations, $\vartheta_{\rm R}^{n} i = {\rm Tr} \ b_{\rm R} (0) \vartheta_{\rm R}^{n}$, are given exactly by

$$h \mathcal{P}_{R} i = \frac{X^{N}}{e^{-!} + 1} \frac{g_{0k}}{e^{-!} + 1};$$
 (29a)

$$h \mathcal{P}_{R}^{2} i = \sum_{k=1}^{N} g_{0k}^{2} \frac{e^{-1} + 1}{(e^{-1} - 1)^{2}} + h \mathcal{P}_{R} i^{2}$$
: (29b)

U sing the spectral distribution given by Eq.(22) one can rew rite these form ulas as

$$h \mathcal{P}_{R} i = \frac{p \frac{p}{2N}}{e^{-!} 1} A_{2} (N);$$
 (30a)

$$h P_R^2 i = \frac{e^{-!} + 1}{(e^{-!} - 1)^2} A_1 (N) + h P_R i^2;$$
 (30b)

where

$$A_n (N) = \frac{X^N}{p n N^2} \exp \frac{(k k_0)^2}{n N^2} :$$
 (31)

O ne can verify numerically that $A_n (N)$ is a real constant obeying $A_n (N)$ 6 1 for all N. Now we can implement the them odynamic limit just by taking N ! 1, with some additional care. For the high-temperature limit (~! ! 0), we see that these expectation values explode. However, this regime is supposed not to be suitably described by the master equation, which presumes weak coupling. For the low-temperature limit (~! ! 1) on the other hand, a solution for the formulas (30) is allowed to exist for N ! 1, in agreement with the analysis carried out in Sec.IIIC. Finally, we note that in this case we have $M_R^2 i$! $M_R^2 i^2$, indicating the disappearance of some types of correlations.

The structure of equation (28) may be promptly identi ed to the phase-destroying master equation (1) com – monly used in SID approaches. However, a rather in portant conceptual di erence must be noted in the (nonunitary) di usion term, which is in fact the one responsible for phase decoherence. In SID approach, \dot{P} denotes the Ham iltonian of the system, whereas in EID approach the decoherence of the system is induced by its interaction with the reservoir. If on one hand these approaches share the sam e m athem atical structure, on the other they cannot be conciliated physically.

The dependence of Eq.(28) on t occurs due to the nature of the coupling employed to model the reservoir. In fact, this is expected for models of reservoir inducing G aussian decay in the quantum coherences, as can be concluded from the form ulas (1)-(3). A natural extension of our model in order to contemplate the exponential decay commonly observed in several contexts may be tried by assuming an adequate temporal dependence for the coupling g_{0k} . R essarch on this topic is in progress.

IV. D ISENTANGLEMENT INDUCED BY PHASE DECOHERENCE

Consider now the case in which the system of interest is composed by two coupled oscillators. The H am iltonian of the system is now considered to be

$$\mathbb{H}_{S} = -!_{a} \mathbf{b}_{a} + -!_{b} \mathbf{b}_{b} + -g_{ab} \mathbf{b}_{a} \mathbf{b}_{b}$$
(32)

A dopting the same algebraic procedure of Sec.III, we obtain the characteristic function $f(x) = \frac{1}{2} \int_{-\infty}^{\infty} \frac{1}{2} \int_$

$$C_{\substack{m_{a};m_{0}\\m_{b};m_{0}}}(t) = \bigotimes_{k=1}^{Q} \frac{1 e^{-\frac{1}{k}}}{1 e^{-\frac{1}{k}} e^{-\frac{1}{(m_{a}-m_{0}^{0})}g_{ak} + (m_{b}-m_{0}^{0})}g_{bk}};$$
(33)

which is associated to the two-boson state

$$b_{S}(t) = \sum_{\substack{m_{a};m_{a}^{0} = 0 \\ m_{b};m_{b}^{0} = 0 \\ m_{b};m_{b}^{0} = 0 \\ e^{\{(m_{b}, m_{b}^{0})\} \cdot b^{t}} jm_{a}m_{b} jm_{a}^{0}m_{b}^{0}j j} (34)$$

where

$$m_{a}; m_{b}^{0} = lm_{a}m_{b}j_{ab}(0) jn_{a}^{0}m_{b}^{0}i:$$
(35)

A lso, it is possible to show that

The analysis realized in the precedent sections concerning the short and long-time regimes can be also applied here with minimal changes. For the short-time limit we obtain

$$t^{c}_{m_{a};m_{b}^{a}}(t) t^{2} = e^{t^{2} = \frac{2}{ab}};$$
(37)

where the decoherence time is now given by

$${}_{ab}^{2} = \frac{X^{N}}{{}_{k=1}} \frac{g_{ak} (m_{a} m_{a}^{0}) + g_{bk} (m_{b} m_{b}^{0})}{2 \sinh(- ! = 2)}^{2} : (38)$$

These expressions show how the decoherence process transform s

$$b_{S}(0) = \bigvee_{\substack{m_{a};m_{a}^{0} = 0 \\ m_{b};m_{b}^{0} = 0}}^{X^{1}} j_{m_{a};m_{b}^{0}} j_{m_{a};m_{b};m_{a}^{0}} j_{m_{a};m_{b};m_{b}^{0}} j_{m_{b};m_{b}^{0}} j_{m_{b};m_{b};m_{b}^{0}} j_{m_{b};m_{b};m_{b}^{0}} j_{m_{b};m$$

into

$$b_{\rm S} (t > {}_{\rm ab}) = \sum_{\substack{m_{\rm a}, m_{\rm b} = 0}}^{M_{\rm a}, m_{\rm a}, m_{\rm a}} jn_{\rm a}m_{\rm b}im_{\rm a}m_{\rm b}j (40)$$

which has no o -diagonal term s in the Fock basis. Here again, additional care is required for the analysis of the critical temperature regime (for N nite), in which jC f may assume nite values in the equilibrium, indicating the presence of surviving entanglement. Also, as it occurred in the precedent sections, for extremely low temperatures the degree of entanglement may remain practically unchanged.

Let us now consider the simplest entangled initial state for the system of two oscillators, namely

$$j_0 i = \frac{1}{\frac{p}{2}} \quad jnni + jmmi; \quad (41)$$

with n \in m. Since the interaction does not couple different number states, the above vector plays the role of a two-qubit state coupled to the reservoir. In this case, we may follow the procedure adopted in ref.[20], in which entanglement has been evaluated by means of two-qubits measures, as for instance, the negativity. This quantity has the following de nition [21, 22]:

$$N [b(t)] = 2m ax f0; _{neq}(t)g;$$
 (42)

where n_{eg} (t) is the negative eigenvalue of the partially transposed density matrix in the joint space E_{ab} . Follow – ing [20] we obtain analytically:

In the high-tem perature and short-tim e regim e we obtain

neg (t) =
$$\frac{1}{2}e^{(n m)^2 t^2 = \frac{2}{ab}}$$
; (44)

where

$$_{ab} = 2G_{ab}^{1} \sinh(\sim !=2);$$
 (45)

with G_{ab}^2 $P_k (g_{ak} + g_{bk})^2 = 2$. Equations (44) and (42) show how entanglement disappears as the dynam ics takes place, provided the tem perature is higher than the critical value given by Eq.(21) for the case of N nite. Also, Eq.(44) makes explicit that the disentanglement is induced by the same mechanism that produces decoherence, namely, the G aussian decay in the characteristic function C (t).

V. CONCLUDING REMARKS

Inspired by a recent work [11] we propose and theoretically investigate the properties of a reservoir composed by harmonic oscillators coupled to the system of interest by means of a cross-K err interaction. We have shown that this type of phase coupling is su cient to guarantee decoherence without dissipation.

Since the analysis have been carried out analytically in virtue of the sim plicity of the model, the main features of our cross-K err reservoir could be clearly identied. For the regime of short times and nite temperatures, a state-independent decoherence time has been estimated and a G aussian decay for quantum coherences has been demonstrated through the analysis of the characteristic function $C_{m,m} \circ (t)$.

Concerning the coupling, we have concluded that the precise pro le of the spectral distribution is relevant only in order to guarantee the irreversibility of the dynam ics, not being important to decide whether the decoherence m ay occur or not. In fact, the mechanism responsible for the decoherence has been shown to be the G aussian decay (derived from the features of both the coupling and the therm alstate), which is always present in the characteristic function for short times independently either on the regime of tem perature or the num ber of oscillators in the reservoir.

By analyzing the long-time dynamics for nite reservoirs we have detected regimes of partial decoherence, in which the occurrence of surviving o -diagonal terms in the quantum state is expected. In this case, it was possible to identify a critical temperature beyond which decoherence is expected to be total. In the therm odynamic limit (N ! 1), however, the regime of partial decoherence no longer exists since $T_{\rm crit}$! 0.

W e have also derived a phase-destroying master equation without statistical assumptions in the weak coupling approximation. The therm odynamic limit has been derived for the regime of low temperatures, indicating the

- [L] W H.Zurek, Phys. Rev. D 26, 1862 (1982); Rev. M od. Phys. 75, 715 (2003).
- [2] E. Joos, H. D. Zeh, C. K iefer, D. G iulini, J. Kupsch and I.O. Stam atescu, Decoherence and the appearance of a Classical W orld in Quantum Theory, 2nd ed. (Springer, New York, 2003).
- [3] G J.M ilburn, Phys. Rev. A 44, 5401 (1991).
- [4] R.Bonifacio, S.O livares, P.Tom besiand D.V itali, Phys. Rev. A 61, 053802 (2000).
- [5] G W .Ford and R F.O 'C onnell, Phys. Lett. A 286, 87 (2001).
- [6] M. Frasca, Phys. Lett. A 283, 271 (2001); Phys. Lett. A 306, 184 (2002) - Erratum; Phys. Lett. A 308, 135 (2003).
- [7] M. Castagnino and R. Laura, Phys. Rev. A 56, 108 (1997); Phys. Rev. A 62, 022107 (2000).
- [8] D F.W alls and G J.M ilbum, Quantum Optics (Springer, Berlin, 1996).
- [9] C W .Gardiner, Quantum Noise (Springer, Berlin, 1991).
- [10] Maximiliam Schlosshauer, Phys. Rev. A 72, 012109 (2005).
- [11] F L. Sem iso and A Vidiella-Barranco, Phys. Rev. A 72, 064305 (2005).

Last but not least, we have veri ed a G aussian decay of the entanglem ent associated to the a bipartite system coupled to our cross-K err reservoir. The result, which has been obtained for arbitrary T and N, illustrates the role of disentanglem ent being induced by the sam em echanism responsible for the decoherence.

A cknow ledgm ents

The authors would like to thank M H Y. M oussa and F L. Sem iso for helpful suggestions and fruitful discussions. KF and ESC acknowledge support from CNPq under projects # 300651=85 6 (KF - partial) and # 140243=2001 1 (ESC).

- [12] J. Lee, M. Patemostro, C. Ogden, et al., New J. Phys. 8, 23 (2006).
- [13] M.Murao and PL.Knight, Phys.Rev.A 58, 663 (1998).
- [14] C J. M yatt, B E. King, Q A. Turchette, et al., Nature (London) 403, 256 (2000); Q A. Turchette, C J. M yatt, B E. King, et al., Phys. Rev. A 62, 053807 (2000).
- [15] P.J. D odd and J.J. Halliwell, Phys. Rev. A 69, 052105 (2004).
- [16] G W . Ford, J.T. Lew is and R F. O 'Connell, Phys. Rev. A 37, 4419 (1988).
- [17] R F. O'Connell and J. Zuo, Phys. Rev. A 67, 062107 (2003).
- [18] G.W. Ford and R.F.O 'Connell, Acta Phys. Hung. B Quantum Electronics B 20, 91 (2004).
- [19] See e.g., M.O. Scully and M.S. Zubairy, Quantum Optics, C am bridge University Press, C am bridge, 1999.
- [20] E S. Cardoso, M C. de O liveira and K. Furuya, Phys. Rev. A 72, 042320 (2005).
- [21] Tzu-Chieh W ei, et al., Phys. Rev. A 67, 022110 (2003).
- [22] F. Verstraete, K. Audenaert and B. D e M oor, Phys. Rev. A 64, 012316 (2001).