A ccurate quantum state estimation via \K eeping the experimentalist honest"

Robin Blume-Kohout

Institute for Quantum Information, Caltech 107-81, Pasadena, CA 91125 USA

Patrick Hayden

School of Computer Science, McGill University, Montreal, QC H3A 2A7 Canaday

In this article, we derive a unique procedure for quantum state estimation from a simple, self-evident principle: an experimentalist's estimate of the quantum state generated by an apparatus should be constrained by honesty. A skeptical observer should subject the estimate to a test that guarantees that a self-interested experimentalist will report the true state as accurately as possible. We also not a non-asymptotic, operational interpretation of the quantum relative entropy function.

Consider a source of quantum states such as a laser, or an ion trap with a preparation procedure. Quantum state estimation is the problem of deducing what state it emits by analyzing the outcomes of measurements on many instances. The usual procedure for state estimation is quantum state tom ography [1, 2], together with some variant of maximum—likelihood estimation [3, 4] to ensure positivity. The obvious goal is an estimate \close" to the true state. Dierent metrics, such as delity [5], relative entropy [6], trace norm, or Hilbert-Schmidt norm [7], will favor dierent estimation procedures.

Here, we derive an optimal state-estimation procedure by rst identifying quantum relative entropy as a unique metric for characterizing an estimate's \goodness". Our procedure is broadly adaptable to (1) arbitrary prior knowledge (or ignorance) and (2) arbitrary measurement procedures.

K eeping the Experim entalist H onest: Implicit in the idea of state estimation is the assumption that some estimates are better than others. Suppose that is an estimate of the true state , and that f(:) is a measure of how \good " an estimate is. We propose that this measure should obey three principles:

- 1. The best estim ate of is itself. If f(:) m easures how well estimates , then f(:) > f(:) for all $\frac{6}{}$.
- 2.f(:) should correspond to some operational test, as the payo or cost of some experimental procedure.
- 3. The \reward" for correctly predicting an event should depend only on the predicted probability for that event. This is a version of the likelihood principle (see [8]).

Remarkably, these simple assumptions single out one measure: the relative entropy between and , or S (k) = Tr(ln ln). It arises as the expected

payo in a type of gam e between a cash-strapped experim entalist and her employer.

A lice, an ambitious scientist attempting to build a quantum computer, produces states that she believes are described by the density operator. She inform sheremployer, Bob, that she has produced the state. Bob, a conscientious scientic administrator, would like to ensure that A lice does not lie { that = . He will periodically visit A lice's lab and measure one of her states, in a way that may depend on her estimate. Her future funding will depend on the outcomes of these measurements. What measurement should Bob perform, and how should he pay A lice, so that she has no incentive to deceive him?

We propose that Bob should measure in a basis $ff_iig_{i=1}^n$ that diagonalizes = $_is_if_iinf_ij$. Upon getting outcome i, he should pay A lice $R_i = C + D$ in s_i dollars, where C and D are non-negative constants. We denote this as the \honest experimentalist reward scheme," or HERS.

HERS motivates honesty: Bob's measurement yields outcome iwith probability $p_i = Tr(j_i h f_i)$. Alice's expected reward is

$$\overline{R} = \sum_{i=1}^{X^{n}} p_{i}R_{i} = C + D \sum_{i=1}^{X^{n}} p_{i} \ln s_{i};$$
 (1)

Rewriting the last term as ${}^{p}_{i} p_{i} \ln s_{i} = \text{Tr ln}$ yields

$$\overline{R}$$
(:) = C + D Tr ln
= C + D [Tr ln (Tr ln Tr ln)]
= C D [H () + S (k)] (2)

Since C, D, and are xed, A lice maxim izes her expected reward by reporting a that minim izes the relative entropy S (k). This constrains to be itself [9]. A lice is thereby motivated to be honest. She is also motivated to produce pure states { but not to lie about how pure the actual state is.

HERS is unique: Unless Bob can do non-projective POVMs [31], this turns out to be the only veri cation procedure that satis es our three criteria.

In classical statistics, a reward scheme for a probabilistic forecast is a scoring rule. It assigns an average reward \overline{R} (P:Q) to a forecast Q when events are distributed according to P. A reward that is uniquely maximized by an

E lectronic address: robin@blum ekohout.com

 $^{{}^{}y}$ E lectronic address: patrick@ cs.m cgill.ca

honest forecast is a strictly proper scoring rule or SPSR (see review [10]). Given some P, the maximum reward under such a rule is P's value, G(P) \overline{R} (P:P). Savage showed that for every SPSR, G(P) is strictly convex [11].

To consider the quantum case, we observe that a measurem ent transforms a state into a probability distribution fP $_{i}$ g over outcomes, to which we can apply a scoring rule. We represent a projective measurement of basis B as a quantum channel . For any state , let []= P.P is a diagonal matrix of probabilities, and simply annihilates o -diagonal elements. Let G() be the maximum of G([]) over all . Since the eigenvalues of majorize those of [][12], and G is convex, this maximum is achieved when []= . Thus G()= G(fig), where fig are the eigenvalues of .

Lem m a 1. Given a physical state and an estimate, Bob can ensure A lice's honesty by applying a SPSR to the probabilities for a measurement of basis B if and only if is diagonal in B.

Proof: Represent Bob's -dependent measurement as a CP-map that annihilates o-diagonal elements in basis B. Let the SPSR yield a value G.

1. Suppose that B diagonalizes , so [] = . Then A lice's expected reward is

The inequalities are simultaneously saturated if and only if =, in which case A lice earns the full value G () of her state. When \oplus , one or both of the inequalities is strict, so A lice earns strictly less than G (). Thus, A lice maxim izes her reward uniquely by reporting =.

2. Suppose that there exists a so that $[] \in$. Then let = []. Since $^2 =$, [] = = []. A lice's expected reward is

$$\overline{R}$$
 ([]: []) = \overline{R} (;) = G (): (4)

whereas if she (truthfully) reports , she can expect

$$\overline{R}$$
 ([]: []) = G ([]) G (); (5)

where the inequality holds because 's eigenvalues majorize those of [] (by Schur's theorem [12]), and because G is convex. A lice expects the same reward for reporting or \uplies , so her honesty is not ensured. \square

So far we have dem anded only that our scoring rule be strictly proper. We now dem and that A lice's reward depend only on her predicted probability for the observed event. In other words, $R_i(fs_jg) = R_i(s_i)$. This rejects the Likelihood P rinciple [13]: all the relevant information in an event is contained in the likelihood of the hypothesis (here, p(ij)). How the experiment was performed is irrelevant. In particular, this avoids any argument between A lice and B ob about how to describe the outcome [s] that did not occur.

A remarkable theorem by Aczel (see also [8]) then restricts the form of the reward function $R_i\left(s_i\right)$.

Theorem 1 (Aczel [14]). Let n 3. The inequality

$$X^n$$
 X^n X^n $p_i R_i (q_i)$ $p_i R_i (p_i)$ (6)
$$i=1 i=1$$

is satis ed for all n-point probability distributions $(p_1 ::: p_n)$ and $(q_1 ::: q_n)$ if and only if there exist constants $C_1 ::: C_n$ and D such that for all i 2 [1:::n],

$$R_{i}(p) = D \ln p + C_{i}$$
: (7)

In the scenario we consider, there is even less freedom . A czel's theorem allows the constants C_i to depend on i. In the quantum setting, all the C_i must be equal to a xed C independent of i (see proof in Appendix A). We have therefore proved the following:

Theorem 2 (The honest experim entalist). Let A be a quantum system with dimension n 3. Let and be density operators for A, and let figig be an orthonormal basis for A that depends only on . De ning $p_i = hg_i j$ $jg_i i$ and $s_i = hg_i j$ $jg_i i$, suppose that

$$X \qquad \qquad X \qquad \qquad p_{\underline{i}}R_{\underline{i}}(s_{\underline{i}}) \qquad p_{\underline{i}}R_{\underline{i}}(p_{\underline{i}}) \qquad \qquad (8)$$

is satis ed for all and , with equality if and only if = . Then fig_iig diagonalizes , and there exist constants C and D such that R $_i$ (s_i) = C + D ln s_i for all i.

The scheme outlined (HERS) is uniquely specified by Bob's need to guarantee Alice's honesty through self-interest. Relative entropy, S (k), appears naturally as the amount of money that Alice can expect to lose by lying. Any \boss" who wishes never to be lied to must use the HERS payment scheme.

Of course, rewards in the real world are often structured less wisely. We propose, however, that a maxim ally ethical scientist should act as if she were being motivated by HERS, and use S (k) as the universal measure of honesty. Hereafter, we will assume that the experimentalist is, in fact, honest.

The Uncertain Experimentalist: What does honesty" mean for an experimentalist who is not certain of? We assert that she should behave as if she were guided by a tangible, strictly proper, reward scheme. HERS is an excellent candidate, but our proofs hold for any strictly proper scheme.

Suppose that A lice does not know , but knows that it will be selected from an ensemble ()d (or simply () hereafter, for clarity). Equivalently, she thinks the true state is with probability (). Her expected reward (from HERS) for reporting is:

$$\overline{R} = R(:) ()d$$

$$Z \qquad Z$$

$$= C D H() ()d + S(k) ()d$$

$$Z$$

$$= const + D Tr(ln) ()d$$

$$= const + D [Tr(ln)] = const^0 D[S(k)];$$

where = R ()d. The \const " term s are independent of and therefore out of A lice's control. Therefore, A lice maxim izes her honesty by reporting the mean of her probability distribution.

The uniqueness of HERS depends on the Likelihood Principle. However, the mean of the probability distribution is maximally honest for any strictly proper scoring rule:

Theorem 3. Let A lice believe that is selected from a distribution (). Let her expected reward for reporting be R (:) = $_{i}p_{i}R_{i}$ (), where p_{i} = TrE $_{i}$, and R (:) > R (:) for all $_{f}$. Aplice maxim izes her expected reward by reporting = $_{i}$ ()d.

Proof: Since A lice expects to appear with probability (), her expected reward is:

$$\overline{R} = R(:) ()d (9)$$

$$Z X$$

$$= Tr(E_{i})R_{i}() ()d (10)$$

$$X Z$$

$$= TrE_{i} ()d R_{i}() (11)$$

$$= R (:) :$$
 (12)

R is strictly proper, so the unique maximum of R \subset :) is at = =.

Consider, instead if A lice had tried to maxim ize delity [15], which is not derived from an operational procedure, but would guarantee A lice's honesty when she knows exactly [32]. Suppose that A lice knows that is either $0 \text{ in} 0 \text{ jor } j + j + j \text{ with equal probability. The delity between any and a pure state is F (; j in j) = h j j i, so the average delity is just:$

$$\overline{F} = Tr(\overline{}); \tag{13}$$

where $\overline{} = \frac{1}{2}$ (0 in 0 j + j + in + j). To maxim ize $\overline{}$, A lice would choose the largest eigenstate of $\overline{}$ { not $\overline{}$ itself. Thus, while delity appears at rst like a good measure of honesty, it does not generally motivate A lice to report the mean of her distribution. Moreoever, it can motivate her to report a pure state that she knows is not the true state.

This is not simply a dierent de nition of honesty. An experim entalist who reports a pure state ji is predicting that some event will never occur. If A lice reports j0i, she is asserting that no measurement will ever yield j1i. In the presence of any uncertainty whatsoever, this is at best misleading, and at worst an outright lie.

This illustrates that HERS strongly penalizes overoptim ism. If B ob obtains an outcome jf_ihf_ij for which hf_ij $jf_ii=0$, A lice will lose in nitely much money! A truly zero-probability event is one against which a gambler would bet in nite money, at arbitrarily bad odds. Reporting p=0 for an event that could conceivably happen is in nitely misleading, and should be discouraged.

The Inform ed Experim entalist: How should Alice use the results of measurements (that she has performed) to reduce her uncertainty? Suppose that she has performed POVM measurements on N copies of , where the ith result corresponds to a positive operator E $_{\rm i}$. She knows two things:

- 1. is selected at random from an ensemble described by $_{0}$ ().
- 2. Through experiments on copies of , she has obtained a measurement record M = fE₁;E₂:::E_N g.

Suppose that she reports $_{j}$ when M $_{\underline{j}}$ occurs, and $_{\underline{j}}$ paid according to a SPSR where $_{\underline{R}}$ (:) = $_{j}$ Tr($_{\underline{j}}$ fihf_{$_{i}$} $_{\underline{j}}$)R_i().

Since appears with probability $_0$ (), the event \ appeared, M $_j$ was measured, and $_j$ was reported" occurs with probability $_0$ () p (M $_j$ j). A lice's expected reward over all possible events is:

We rewrite this using

$$Z$$
 p_{j} $p(M_{j}j)_{0}()d$, and (14)
 $-\frac{1}{p_{j}}$ $p(M_{j}j)_{0}()d$; (15)

to get

$$\overline{R} = X p_j Tr jf_i hf_i \overline{j}_j R_i (j) = X p_j R (\overline{j}_j : j):$$

 \overline{R} is strictly proper, so by setting $j = \overline{j}$ we uniquely maxim ize each term in the sum, and Equation 15 de nest the optim alestim ate of , given M j.

Equation 15 is nothing other than Bayes' Rule. Thus, the mean of a Bayesian—inferred distribution over states is the unique optimal estimate { for any strictly proper reward scheme. We formalize this in the following theorem:

Theorem 4. If is drawn from an ensemble $_0$ (), and a measurement M $_j$ with conditional probability p (M $_j$ j) is observed, then every strictly proper scoring rule \overline{R} (:) is maximized by:

1. U sing Bayes' Rule and Bom's Rule:

2. Reporting the mean of $_{\rm M}$ ().

This applies not only to relative entropy, our preferred measure of honesty, but to any honesty-guaranteeing reward scheme. We conclude that Bayesian inference is the unique solution to honest state estimation.

O ther procedures will not optimize any measure of honesty derived from a strictly proper scoring rule. A lternative measures ofhonesty will either (a) in some circumstances, motivate an experimentalist to at-out lie about the state, or (b) not be operationally implementable (e.g., delity). Our previous discussion of delity illustrates that a non-operational metric that guarantees the honesty of a knowledgeable experimentalist can fail dramatically in the face of uncertainty.

Inform ation theorists have previously interpreted relative entropy as a good measure of two states' distinguishability [16,17] { indeed, as the only meaningfulone in the limit of many copies. We have invoked the Likelihood Principle rather than the many-copy limit, but we can easily allow Bob to jointly measure N copies of . He must then apply a SPSR to the result, and A lice can expect a reward \overline{R} (N : N). As N ! 1 , relative entropy remains meaningful, unlike other measures (e.g., the B rier score [10]).

In the presence of uncertainty, pure (or rank-de cient) states are in nitely dishonest estimates. Estimating a pure state means predicting that some event will never happen. Anyone taking such a prediction seriously would be justified in betting in nitely much money, at arbitrarily bad odds, against that event { and should therefore expect to lose in nitely much, if the estimate is incorrect. This should be discouraged.

Bayesian state estimation has been discussed previously [18, 19, 20], especially for pure states [21, 22]. The predom inance of other methods such as maximum likelihoood, in the current literature (e.g., [23, 24, 25, 26, 27, 28, 29, 30], indicates that it has not received the attention it deserves. Our goal in this letter is to provide a concrete and compelling argument for Bayesian state estimation { and to call attention to the problematic implications of pure-state estimates.

A cknow ledgm ents

RBK thanks Daniel James (for pointing out the problem in the rst place), Jon Yard, John Preskill, and NSF contract PHY-0456720. PH thanks Howard Barnum, Simon Benjam in and especially Chris Fuchs for helpful

discussions, as well as the C anada R esearch C hairs program , the C anadian Institute for A dvanced R esearch and N SERC for support.

APPENDIX A: FURTHER RESTRICTING THE REWARD FUNCTION

We show that the quantum reward function is more tightly constrained than the classical one: the constants C_i of Theorem 1 must all be equal. We can assume that D=1. Assuming a reward function of the form specied by Theorem 1, the inequality \overline{R} (:) \overline{R} (:) is then equivalent to

$$S(k) + X$$
 $C_{i}(r_{i} p_{i}) 0:$ (A1)

Let U (t) be a sm ooth curve in the unitary group de ned on a neighborhood of t=0 and such that U(0)=I. A lso let (t)=U(t) $U^{Y}(t)$ and

$$g(t) = S(k(t)) + C_{i} r_{i} Tr U(t) j_{i} he_{i} j_{i} Y(t)$$

be the function de ned by substituting (t) into the expression of Eq. (A1). (Here $= i_1 r_1 \dot{p}_1 \dot{r}_2 \dot{r}_1 \dot{p}_2 \dot{r}_2 \dot{r}_2$

$$\frac{d}{dt}$$
S (k (t)) = Tr U-(ln)U^y + U (ln)U^y; (A2)

where U = dU = dt and the dependence of U on thas been suppressed. Because $U + U^y = 0$ when t = 0, $\frac{d}{dt}S$ (k (t)) $j_{=0} = 0$. Likewise, p_i equals 0 when t = 0. We must therefore determine the second derivative of g at t = 0 and show that for a suitable choice of curve, this derivative is negative. In that case, $g(t) = g(0) + g(0)t^2 = 2 + 0$ (t^3) with g(0) < 0, implying that g(t) is negative for su ciently small t.

So, di erentiating again, we nd

$$\frac{d^2}{dt^2}$$
S (k (t)) = 2Tr [X;](ln)X; (A3)

where X is a Herm itian matrix such that U=iX. We have made use of the identity $U+U^y=2U-U^y$ that can be proved by dierentiating $UU^y=I$. Because S (k (t)) 0, the expression in Eq. (A 3) must also be nonnegative.

Dierentiating the second term of g(t), we nd that

$$\frac{d^2g}{dt^2} = 2 \text{Tr} [X;] (\ln + B) X; \qquad (A4)$$

where $B = P_{i} C_{i} \dot{p}_{i} \dot{n}_{i} \dot{p}_{i}$

We will now show that all the C_i must be equal. Assume without loss of generality that $C_1 \in C_2$. Let $r_j = \exp((C_j + 2 \ln 2) = 2)$ for j = 1; 2 and $r_3 = 1$ r_1 r_2 .

W ith these choices, r_1 ; r_2 1=2, making a density operator. Now write ~ and B for the restriction of and B to the span of the rst two eigenvectors of . O beeve that there exists a choice of X also with support only on this subspace such that Tr [X; ~](ln ~)X > 0.

W ith these choices, and noting that Tr [X; ~]X = 0,

$$h$$
 i $g(0) = 2Tr [X; \sim](\ln \sim)X < 0$: (A 5)

Because g(0) < 0 contradicts the requirement that D (k (t)) 0, we conclude that $C_i = C_1$ for all i. \square

- [1] K. Vogeland H. Risken, Phys. Rev. A 40, 2847 (1989).
- [2] G.M.D'Ariano, M.G.A.Paris, and M.F.Sacchi, Advances in Imaging and Electron Physics 128, 205 (2003).
- [3] Z.H radil, Phys. Rev. A 55, R1561 (1997).
- [4] Z. Hradil, J. Rehcek, J. Fiursek, and M. Jezcaronek, in Quantum state estimation, edited by M. G. A. Paris and J. Rehacek (Berlin, Germany: Springer-Verlag, 2004, 2004), vol. 649 of Lecture Notes In Physics, pp. 59{112.
- [5] E.Bagan, A.Monras, and R.Muoz-Tapia, Phys.Rev.A 71,062318 (2005).
- [6] F. Tanaka and F. Komaki, Phys. Rev. A 71, 052323 (2005).
- [7] J. Rehacek, B.G. Englert, and D. Kaszlikowski, arxiv.org/quant-ph 05,0405084 (2004).
- [8] J. Bernardo, Annals of Statistics 7, 686 (1979).
- [9] M. Ohya and D. Petz, Quantum entropy and its use, Texts and monographs in physics (Springer-Verlag, Berlin, 1993).
- [10] T. G neiting and A. E. Raffery, Tech. Rep. 463R, Dept. of Statistics, Univ. of W ashington (2005).
- [11] L.J.Savage, Journal of the Am erican Statistical Association 66, 783 (1971).
- [12] R.Bhatia, Matrix Analysis (Springer, 1996).
- [13] J.O.Berger and R.L.W olpert, The Likelihood Principle: A Review, Generalizations, and Statistical Implications (with discussion)., IM S Lecture Notes-Monograph Series (Institute of Mathematical Statistics, 1988).
- [14] J.A czel, Journal of Mathematical Analysis and Applications 75, 447 (1980).
- [15] R. Jozsa, Journal of Modern Optics 41, 2315 (1994).
- [16] V. Vedral, M. B. Plenio, K. Jacobs, and P. L. Knight, Phys. Rev. A 56, 4452 (1997).
- [17] C. A. Fuchs, PhD. thesis, University of New Mexico (1995).
- [18] R.Derka, V.Buzek, G.Adam, and P.L.Knight, Journal of Fine Mechanics and Optics 11-12, 341 (1996).

- [19] R. Schack, T.A. Brun, and C.M. Caves, Physical Review A 64, 014305 (2001).
- [20] F.Neri, arxiv.org/quant-ph p. 0508012 (2005).
- [21] K.R.W. Jones, Annals of Physics 207, 140 (1991).
- [22] K.R.W. Jones, Physical Review A 50, 3682 (1994).
- [23] D.F.V. James, P.G.Kwiat, W.J.Munro, and A.G. White, Physical Review A 64,052312 (2001).
- [24] K. J. Resch, P. W alther, and A. Zeilinger, Phys. Rev. Lett 94,070402 (2005).
- [25] Y. S. Weinstein, T. F. Havel, J. Emerson, N. Boulant, M. Saraceno, S. Lloyd, and D. G. Cory, Journal of Chemical Physics 121, 6117 (2004).
- [26] J.L.O'B rien, G.J.Pryde, A.G ilchrist, D.F.V. James, N.K. Langford, T.C. Ralph, and A.G. W hite, Physical Review Letters 93, 080502 (2004).
- [27] S. H. Myrskog, J. K. Fox, M. W. Mitchel, and A. M. Steinberg, Phys. Rev. A 72, 013615 (2005).
- [28] C. F. Roos, G. P. T. Lancaster, M. Riebe, H. H. ner, W. Hnsel, S. Gulde, C. Becher, J. Eschner, F. Schmidt-Kaler, and R. Blatt, Phys. Rev. Lett 92, 220402 (2004).
- [29] J.B.A Itepeter, D.Branning, E.Je rey, T.C.Wei, P.G. Kwiat, R.T.Thew, J.L.O'Brien, M.A.Nielsen, and A.G.White, Physical Review Letters 90, 193601 (2003).
- [30] H. Haener, W. Haensel, C. F. Roos, J. Benhelm, D. C. alkar, M. Chwalla, T. Koerber, U. D. Rapol, M. Riebe, P.O. Schmidt, et al., Nature 438, 643 (2005).
- [31] Characterizing Bob's non-projective measurements is more complicated. For now, we note that: (1) in general, a xed informationally complete POVM su ces; (2) our results remain largely unchanged, but Alice's expected reward may involve C[] and C[] instead of; , where C is a quantum channel.
- $\ensuremath{\mbox{[32]}}$ Jozsa showed [15] that F (;) is maximal if and only if