

Quantum particles trapped in a position-dependent mass barrier; a d-dimensional recipe

Omar Mustafa¹ and S. Habib Mazhar Mousavi²
 Department of Physics, Eastern Mediterranean University,
 G. Magusa, North Cyprus, Mersin 10, Turkey
¹e-mail: omarmustafa@emueu.edu.tr
²e-mail: habibmazhari@emueu.edu.tr

April 1, 2022

Abstract

We consider a free particle, $V(x) = 0$; with a position-dependent mass $m(x) = 1/(1 + \alpha^2 x^2)^2$ in the d-dimensional Schrödinger equation. The effective potential turns out to be a generalized Pöschl-Teller potential that admits its exact solution.

PACS numbers: 03.65.Ge, 03.65.Fd, 03.65.Ca

1 Introduction

Quantum mechanical particles endowed with position-dependent mass (PDM), $M(x)$, have attracted attention and inspired intense research activities over the years [1-16]. They form interesting and useful models for many physical problems. They are used, for example, in the energy density many-body problem [2], in the determination of the electronic properties of the semiconductors [3] and quantum dots [4], in quantum liquids [5], in the ^3He clusters [6] and metal clusters [7], in the Bohmian approach to quantum theory [8], full/partial Gaussian wave-packet revival inside an infinite potential [9], etc.

In addition to its practical applicability side, conceptual problems of delicate nature erupt in the study of quantum mechanical systems with position-dependent mass (e.g., momentum operator does not commute with $m(x)$, uniqueness of kinetic energy operator, etc.). Comprehensive discussion on such issues could be found in, e.g., [8, 10, and related references therein].

On the other hand, the exact solvability of the d-dimensional Schrödinger equation may very well form the major ingredient for methods that are based on a Liouvillean-type change of variables (cf., e.g., [17-19]) such as the point canonical transformation (PCT) method (cf., e.g., [13, 14]). Where, in such

methodologies, the exact solution of the so-called reference Schrodinger equation (eigenvalues and eigenfunctions) is mapped into an exact solution (eigenvalues and eigenfunctions) of the so-called target Schrodinger equation (cf, e.g., [1,14]).

In a recent study, we have introduced a d-dimensional regularization of the PCT-method for some PDM-quantum mechanical particles [1]. Therein, inter-dimensional degeneracies associated with the isomorphism between the angular momentum quantum number ℓ and dimensionality d are incorporated through the central repulsive/attractive core $\ell(\ell+1)=r^2 \rightarrow \ell_d(\ell_d+1)=r^2$ (where $\ell_d = \ell + (d-3)/2$ for $d \geq 2$) of the spherically symmetric Schrodinger equation (cf, e.g., [20], Gang in [10] and Quesne in [10] for more details).

In this work, we study "free" particles, $V(r) = 0$, trapped in their own position-dependent mass barriers (hence, labeled as quasi-free particles), where inter-dimensional degeneracies remain intact in the d-dimensional radial Schrodinger equation (in atomic units $\hbar = m = 1$, the position-dependent mass $M(r) = m(r)$; and with $\ell = 0$ and $\ell = 1$ in Eq.(1.1) of Tanaka in [10].)

$$\frac{d^2}{dr^2} \frac{\ell_d(\ell_d+1)}{r^2} + \frac{m^0(r)}{m(r)} \frac{d}{2r} \frac{d}{dr} + 2m(r)E_{n_r, \ell_d}(r) = 0: \quad (1)$$

Where $n_r = 0; 1; 2; \dots$ is the radial quantum number, and $\ell_d(r) = \ell(r) = dr$. Moreover, the $d = 1$ can be obtained through $\ell_d = -1$ and $\ell_d = 0$ for even and odd parity, $P = (-1)^{\ell_d+1}$, respectively (cf, e.g., Mustafa and Znojil in [20]). Nevertheless, the inter-dimensional degeneracies associated with the isomorphism between angular momentum ℓ and dimensionality d builds up the ladder of excited states for any given n_r and nonzero ℓ from the $\ell = 0$ result, with that n_r , by the transcription $d \rightarrow d+2$: That is, if $E_{n_r, \ell_d}(d)$ is the eigenvalue in d-dim ensions then

$$E_{n_r, \ell_d}(2) = E_{n_r, \ell_d+1}(4) = \dots = E_{n_r, \ell_d}(2\ell_d+2) \quad (2)$$

for even d , and

$$E_{n_r, \ell_d}(3) = E_{n_r, \ell_d+1}(5) = \dots = E_{n_r, \ell_d}(2\ell_d+3) \quad (3)$$

for odd d . For more details on inter-dimensional degeneracies the reader may refer to a sample of references in [20].

With the PCT-method in point (cf, e.g., [1]), a substitution of the form $R(r) = g(r) \sqrt{q(r)}$ in (1) would result in $g(r)^2 q^0(r) = m(r)$, manifested by the requirement of a vanishing coefficient of the first-order derivative of $q(r)$ (hence a one-dimensional form of Schrodinger equation is achieved), and $q^0(r)^2 = m(r)$ to avoid position-dependent energies-multiplicity (i.e., $2E_{n_r, \ell_d}(r) = q^0(r)^2 = 2E$). Hence,

$$q(r) = \int_r^{\infty} \frac{1}{m(t)} dt \Rightarrow g(r) = m(r)^{1/4}: \quad (4)$$

This in effect implies

$$\frac{1}{2} \frac{d^2}{dq^2} + V_{\text{eff}}(q(r)) = E_{n_r, \ell_d}(q(r)) = E_{d, n_r, \ell_d}(q(r)); \quad (5)$$

with an effective potentials

$$V_{\text{eff}}(q(r)) = \frac{l_d(l_d + 1)}{2r^2 m(r)} U_d(r) \quad (6)$$

where

$$U_d(r) = \frac{m^0(r)}{8m(r)^2} - \frac{7m^0(r)^2}{32m(r)^3} + \frac{m^0(r)(d-1)}{4rm(r)^2}; \quad (7)$$

2 Consequences of an asymptotically vanishing mass settings as $r \rightarrow 1$

A "free" particle with an asymptotically vanishing position-dependent mass $m(r) = 1/(1 + \xi^2 r^2)^2$ would experience an effective potential

$$V_{\text{eff}}(q(r)) = \frac{\xi^2}{2} \left\{ \frac{(l_d - 1)}{\sin^2(\xi q)} + \frac{(l_d - 1)}{\cos^2(\xi q)} - \frac{\xi^2}{2} \right\}; \quad (8)$$

where

$$q(r) = \frac{1}{\xi} \arctan(\xi r) \quad \&r = \tan(\xi q); \quad (9)$$

and

$$U_d(r) = \xi^2 d \tan^2(\xi q) + \frac{\xi^2}{2} (1 - 2d) \quad (10)$$

In such settings, Eq. (5) reads

$$\frac{1}{2} \frac{d^2}{dq^2} + \frac{\xi^2}{2} \left\{ \frac{(l_d - 1)}{\sin^2(\xi q)} + \frac{(l_d - 1)}{\cos^2(\xi q)} \right\} E_{n_r, l_d}(q) = E_{n_r, l_d}(q); \quad (11)$$

where

$$\{ (l_d - 1) = l_d(l_d + 1); \quad (l_d - 1) = l_d(l_d + 1) + 2d \quad \text{and} \quad E = E + \frac{1}{2}\xi^2; \quad (12)$$

Equation (11) is obviously a standard one-dimensional form of Schrodinger equation with a generalized Pöschl-Teller effective potential which admits exact solution of the form

$$E_{n_r} = \frac{\xi^2}{2} \left(\{ + \} + 2n_r \right)^2 \quad (13)$$

$$E_{n_r, l_d}(q) = C \sin^{\{ }(\xi q) \cos^{\{ }(\xi q) {}_2F_1\left(n_r; \{ + \} + n_r; \{ + \frac{1}{2}\}; \sin^2(\xi q)\right) \quad (14)$$

with $\{; > 1$; $E_{n_r, l_d}(0) = 0$ and $E_{n_r, l_d} \frac{1}{2\xi} = 0$; as reported by Salem and Montemayor (see Eq.(4.7) in [21]). This in turn would lead to

$$E_{n_r, l_d} = \frac{\xi^2}{2} \left(\left(c + \frac{1}{2} + 2n_r \right)^2 - 1 \right); \quad = \frac{P}{(2l_d + 1)^2 + 8d} \quad (15)$$

$$R_{n_r; l_d}(r) = C^{-l_d+1} (1 + \epsilon^2)^{-\frac{1}{4}(2l_d+5+\epsilon)} {}_2F_1\left(-n_r; c + \frac{\epsilon}{2} + n_r; c; \frac{\epsilon^2}{1 + \epsilon^2}\right) \quad (16)$$

where $\epsilon = \epsilon r$; and $c = l_d + \frac{3}{2}$.

However, for $\epsilon = 0; 1$ (a requirement suggested by relation (12) when $l_d = 0; 1$) the effective potential in (11) collapses into

$$V_{eff}(q(r)) = \frac{\epsilon^2}{2} \frac{(1 - \cos^2(q))}{\cos^2(q)}; \quad (17)$$

Which admits an exact solution

$$E_{n_r} = 2\epsilon^2 \left(n_r + \frac{1}{2}\right)^2 - \frac{\epsilon^2}{2} \quad (18)$$

$$R_{n_r; l_d}(q) = A \cos(q) {}_2F_1\left(-n_r; n_r + \frac{1}{2}; \frac{1}{2}; \sin^2(q)\right); \quad (19)$$

and consequently

$$E_{n_r; 0} = 2\epsilon^2 \left(n_r + \frac{1}{2}\right)^2 - \frac{\epsilon^2}{2} \quad (20)$$

$$R_{n_r}(r) = A^{-l_d+1} (1 + \epsilon^2)^{-\frac{1}{4}(2l_d+5+\epsilon)} {}_2F_1\left(-n_r; c + \frac{\epsilon}{2} + n_r; c; \frac{\epsilon^2}{1 + \epsilon^2}\right); \quad (21)$$

where $\epsilon = (1 + \epsilon^2)^{1/2}$:

3 Concluding Remarks

In this letter, we considered a quasi-free particle with an asymptotically vanishing position-dependent mass $m(r) = 1/(1 + \epsilon^2 r^2)^2$ and radial potential $V(r) = 0$ (i.e., "free" particle in this sense). We have shown that under these settings the particle experiences an effective potential of the form of a Pöschl-Teller, Eq.(8). The exact solution of which is mapped to match the attendant settings of our quasi-free particle with the above mentioned position-dependent mass.

References

- [1] O Mustafa and S H M azharim ousavi 2006 "Point canonical transformation d-dimensional regularization" (arXiv: math-ph/0602044)
- [2] A Puente and M Casas Comput. Mater Sci. 2 (1994) 441
- [3] Bastard G 1988 "Wave Mechanics Applied to Semiconductor Heterostructures" , Les Editions de Physique, Les Ulis
- [4] L I Serra and E Lipparini Europhys. Lett. 40 (1997) 667
- [5] F Arias de Saaverda, J Boronat, A Polls , and A Fabrocini Phys. Rev. B 50 (1994) 4248
- [6] M Barranco , M Pi, S M Gatica ., E S Hernandez ., and J. Navarro Phys. Rev. B 56 (1997) 8997
- [7] A Puente , L I Serra, and M Casas Z. Phys. D 31 (1994) 283
- [8] A R Plastino, M Casas and A . Plastino Phys. Lett. A 281 (2001) 297 (and related references therein)
- [9] A Schmidt Phys. Lett. A (2006) (in press)
- [10] T Tanaka J. Phys. A 39 (2006) 219
C Quesne, Ann. Phys. 321 (2006) 1221
C Gang, Phys. Lett. A 329 (2004) 22
A R Plastino, A Rigo, M Casas, F Garcias, and A . Plastino Phys. Rev. A 60 (1999) 4318
- [11] S H Dong and M . Lozada-Cassou Phys. Lett. A 337 (2005) 313
I O Vakarchuk J. Phys. A ; Math and Gen 38 (2005) 4727
C Y Cai, Z Z Ren and G X Ju Commun. Theor. Phys. 43 (2005) 1019
- [12] B Bagchi, A Banerjee, C Quesne and V M Tkachuk J. Phys. A ; Math and Gen 38 (2005) 2929
J Yu and S H Dong Phys. Lett. A 325 (2004) 194
L Dekar, L Chetouani and T F Hammann J. Math. Phys. 39 (1998) 2551
- [13] C Quesne and V M Tkachuk J. Phys. A ; Math and Gen 37 (2004) 4267
L Jiang, L Z Yi, and C S Jia Phys. Lett. A 345 (2005) 279
A D Alhaidari Int. J. Theor. Phys. 42 (2003) 2999
- [14] A D Alhaidari Phys. Rev. A 66 (2002) 042116
- [15] R De, R Dutt and U Sukhatme J. Phys. A ; Math and Gen 25 (1992) L843

- [16] G Junker J. Phys.A ;M ath and Gen 23 (1990) L881
- [17] J Liouville J.M ath.Pure Appl.1 (1837)16
- [18] M Znojil and G Levai J.M ath.Phys.42 (2001)1996
M Znojil "P T -sym metric form of the Hulthen potential" (2000) (arX iv: m ath-ph/0002017)
- [19] M Znojil and G Levai Phys.Lett.A 271 (2000) 327
- [20] D R Herschbach et al "D im ensionalScaling in Chem icalPhysics" (K luwer Academ ic Publishers 1993, D ordrecht, N etherlands.)
D R Herschbach J.Chem .Phys.84 (1986) 838
H Taseli J.M ath.Chem .20 (1996) 235
O M ustafa and M Znojil J.PhysA ;M ath and Gen.35 (2002) 8929
O M ustafa and M O deh J.Phys.A ;M ath and Gen.32 (1999) 6653
O M ustafa and M O deh J.Phys.A ;M ath and Gen.33 (2000) 5207
M M Nieto Am .J.Phys.47 (1979) 1067
- [21] L D Salem and R M ontanayor Phys.Rev.A 47 (1993) 105
S F lügge 1974, P racticalQ uantum M echanics, Springer, Berlin.