

The certainty principle II

Proof of the uncertainty principle

D. A. Arbaty

March, 2006

Abstract

A more detailed derivation of the Heisenberg uncertainty principle from the certainty principle is given.

Introductory remarks. After publication of the paper [1] I received many letters, including those with references. In this connection, I think it is necessary to specify the following:

1. The metric introduced by me ("quantum angle") is known to mathematicians from 1904 as Fubini-Study metric.
2. The correct "uncertainty relation" (in fact, certainty relation) for the quantities energy - time was first suggested by Mandelshtam and Tamm [2].
3. Mandelshtam and Tamm studied a quantum system in the Schrodinger representation and did not use group theory methods. For this reason they could not understand that their result has more general character.
4. In contrast, I used group theory formulations. And implied that the system can be studied, in particular, in the representation of relativistic canonical quantization [4]. This allowed me to formulate the certainty principle and to suggest more general inequalities.

Some of my critics refused to believe that the Heisenberg uncertainty principle is really a consequence of the certainty principle. They claimed that "this can not be so, because this can never be so".

Nevertheless, this is so. And here I give a more detailed explanation.

Derivation of the uncertainty principle. Suppose that for a given quantum system we succeeded to find some observable X , that can be considered in some sense a "coordinate operator".

Suppose that X is a self-adjoint operator with continuous spectrum, $X = X^\dagger$. Let us denote $E_{(a;b)}$ its spectral projector¹ for an arbitrary real interval $(a;b)$.

Suppose also that we have a self-adjoint operator P , $P = P^\dagger$, such that for any a, b and x we have equality:

$$e^{+i x P} E_{(a+x; b+x)} e^{-i x P} = E_{(a;b)};$$

i. e. P is a "generator of spectral shifts" for X . As we know, such an operator is usually an operator of momentum.

Suppose now that the system is in state i , $\langle i | i \rangle = 1$.

The quantity $\langle i | E_{(a;b)} | i \rangle$, obviously, denotes the probability to find the system inside the interval $(a;b)$. Let us denote such l and r , that

$$\langle i | E_{(l;r)} | i \rangle = \langle i | E_{(r;l+1)} | i \rangle = \frac{1}{2} - \frac{1}{2} \sqrt{1 - \cos^2 l} \quad 0.07926:::$$

<http://daarb.narod.ru/>, <http://wave.front.ru/>

¹Roughly speaking, spectral projector is an operator nullifying wave function in X -representation outside of the given interval.

- [2] L. I. Mandelshtam, I. E. Tamm, "The uncertainty relation between energy and time in nonrelativistic quantum mechanics", J. Phys. (USSR) 9, 249-254 (1945).
 Rus.: L. I. Mandelshtam, I. E. Tamm "Sootnoshenie neopredelyonnosti energiiya-vremya v nerelyativistskoy kvantovoy mekhanike", Izv. Akad. Nauk SSSR (ser. z.) 9, 122-128 (1945), in [3], p. 306-315.]
- [3] L. I. Mandelshtam, Polnoe sobranie trudov, v. 2, M.-L. (1947).
- [4] D. A. Arbatsky "What is 'relativistic canonical quantization'?" (2005),
<http://daarb.narod.ru/wiroq-eng.html>, <http://wave.front.ru/wiroq-eng.html>, arXiv:math-ph/0503029.
 Rus.: D. A. Arbatsky "Chto takoe 'relyativistskoe kanonicheskoe kvantovanie'?" (2005),
<http://daarb.narod.ru/wiroq-rus.html>, <http://wave.front.ru/wiroq-rus.html>, arXiv:math-ph/0503029.]
- [5] J. Hilgevoord, J. Uink "The uncertainty principle", in The Stanford encyclopedia of philosophy, ed. E. N. Zalta (2001), <http://plato.stanford.edu/archives/win2001/entries/qt-uncertainty/>.