

Generation of NPT Entanglement from Nonclassical Photon Statistics

J. Solomon Ivan,^{*} N. Mukunda,[†] and R. Simon[‡]

The Institute of Mathematical Sciences, CIT Campus, Taramani, Chennai 600 113, India

(Dated: April 03, 2006)

With a product state of the form $\rho_{\text{in}} = \rho_a \otimes |0\rangle_{bb}\langle 0|$ as input, the output two-mode state ρ_{out} of the beam splitter is shown to be NPT whenever the photon number distribution (PND) statistics $\{p(n_a)\}$ associated with the possibly mixed state ρ_a of the a-mode is antibunched or otherwise nonclassical, i.e., if $\{p(n_a)\}$ fails to respect any one of an infinite sequence of classicality conditions.

PACS numbers: 03.67.Mn, 42.50.Dv, 03.67.-a, 42.50.Ar

Considerable progress has been achieved in recent years towards an understanding of entanglement in the context of continuous variable bipartite systems[1, 2, 3, 4]. Phase space descriptions and Gaussian states have played a significant role in this progress.

While the quantum uncertainty principle places restrictions on the moments of the phase space variables, separability of a state places additional demands on these moments. Gaussian states are fully determined by their first and second moments, or variances, and so also is the issue of their separability. But non-Gaussian states which satisfy the separability demand on their variances could still be entangled by virtue of violation of one of the separability requirements on their higher order moments. It is only relatively recently that interest in this direction has started emerging[5].

There is a useful connection between nonclassicality and inseparability, and the beam splitter plays an important role as a bridge between these two attributes. Asboth *et al.*[6] have shown that the output of a beam splitter whose input is a product state $\rho_{\text{in}} = \rho_a \otimes |0\rangle_{bb}\langle 0|$ is entangled if and only if the single-mode state ρ_a at the input is nonclassical. [Thus any measure of entanglement of the output state in this configuration is a computable measure of nonclassicality of ρ_a , the entanglement potential (EP)[6]]. For use in quantum information tasks, however, it is important to know if the entanglement generated in this manner is NPT or PPT.

Two canonical manifestations of nonclassicality have been extensively studied in the quantum optics context: (1) squeezing[7], and (2) antibunching or sub-Poissonian fluctuation[8], which is a particular manifestation of nonclassical photon statistics; these are respectively nonclassicalities of the phase-sensitive and phase-insensitive types. Whereas the former has been well explored as a source of entanglement in the context of Gaussian states, the same cannot be said in respect of the latter.

The principal aim of this Letter is to show that with $\rho_{\text{in}} = \rho_a \otimes |0\rangle_{bb}\langle 0|$ as input, the two-mode state after the beam splitter is definitely NPT if ρ_a is nonclassical of the phase-insensitive type, that is if the photon number distribution (PND) statistics $\{p(n_a)\}$ associated with ρ_a of the a-mode is antibunched, or possesses any other higher order nonclassicality. We prove this result in

two stages: first for the restricted case of antibunched input ρ_a wherein we exhibit a surprisingly simple witness; then in the case of arbitrary PND we take advantage of a complete characterization of nonclassicality based on the theory of moments[9, 10]. Our result and mode of proof extends the domain of effectiveness of the partial transpose criterion for separability considerably beyond the traditional Gaussian regime.

Consider therefore a bipartite system, each part being a single mode radiation field, with respective creation and annihilation operator pairs \hat{a}^\dagger, \hat{a} and \hat{b}^\dagger, \hat{b} acting on Hilbert spaces $\mathcal{H}^a, \mathcal{H}^b$. Their only nonvanishing commutators are $[\hat{a}, \hat{a}^\dagger] = [\hat{b}, \hat{b}^\dagger] = 1$. The Fock or photon number states for the two modes provide convenient ONB's for $\mathcal{H}^a, \mathcal{H}^b$ respectively:

$$\begin{aligned} |n_a\rangle &= (n_a!)^{-1/2} (\hat{a}^\dagger)^{n_a} |0\rangle_a, \\ |n_b\rangle &= (n_b!)^{-1/2} (\hat{b}^\dagger)^{n_b} |0\rangle_b, \quad n_a, n_b = 0, 1, 2, \dots \end{aligned} \quad (1)$$

The products $|n_a, n_b\rangle \equiv |n_a\rangle \otimes |n_b\rangle$ form an ONB for $\mathcal{H}^a \otimes \mathcal{H}^b$. The partial transpose of a bipartite state ρ is the operator $\tilde{\rho}$ on $\mathcal{H}^a \otimes \mathcal{H}^b$ defined in this ONB thus:

$$\langle n_a, n_b | \tilde{\rho} | n'_a, n'_b \rangle \equiv \langle n_a, n'_b | \rho | n'_a, n_b \rangle. \quad (2)$$

We have at our disposal this key fact[11]: if the partial transpose $\tilde{\rho}$ is not a valid density operator (NPT), i.e., if $\tilde{\rho} \not\geq 0$, then ρ is definitely an entangled state.

Clearly, a way to show that ρ is NPT is to exhibit an operator \hat{A} of the bipartite system such that the 'expectation value' of the *positive* operator $\hat{A}^\dagger \hat{A}$ in $\tilde{\rho}$ is negative. We can then conclude:

$$\text{Tr}(\tilde{\rho}_{AB} \hat{A}^\dagger \hat{A}) < 0, \text{ for some } \hat{A} \Rightarrow \rho \text{ is NPT.} \quad (3)$$

Alternatively, and in a sense more directly, we may be able to find some principal submatrix of the matrix $\langle n_a, n_b | \tilde{\rho} | n'_a, n'_b \rangle$ representing $\tilde{\rho}$, i.e., a submatrix formed by intersections of any subset of *rows of this matrix and the corresponding columns*, such that this submatrix is not positive definite. Then again we can conclude:

$$\text{A principal submatrix of } \tilde{\rho} \not\geq 0 \Rightarrow \rho \text{ is NPT.} \quad (4)$$

We use both strategies (3) and (4) in what follows.

Nonclassicality and Photon Statistics: For a moment let us restrict attention to states ρ_a of the \hat{a} -mode alone. The separation of its (quantum) states into so called ‘classical’ and ‘nonclassical’ types is based on the diagonal coherent state representation[12] of ρ_a :

$$\rho_a = \int_C \pi^{-1} d^2 z_a \phi(z_a) |z_a\rangle \langle z_a|,$$

$$|z_a\rangle = e^{-\frac{1}{2}|z_a|^2} \sum_{n_a=1}^{\infty} \frac{z_a^{n_a}}{\sqrt{n_a!}} |n_a\rangle, \quad \hat{a}|z_a\rangle = z_a|z_a\rangle.$$

The weight function $\phi(z_a)$ is in general a singular distribution of a certain class. While it is real on account of hermiticity of ρ_a and normalized to unity, it is in general not pointwise nonnegative over the complex plane \mathcal{C} . Classical states are identified as follows.

$$\rho_a \text{ ‘classical’} \Leftrightarrow \phi(z_a) \geq 0 \text{ for all } z_a \in \mathcal{C}. \quad (5)$$

All other states are declared ‘nonclassical’.

All information about any state ρ_a is captured by the expectation values of *all* possible hermitian observables \hat{A} of the system, namely $\text{Tr}(\rho_a \hat{A})$. While any operator \hat{A} can be written in the *normal ordered form* $\hat{A} = \sum_{j,k=0}^{\infty} c_{jk} \hat{a}^{\dagger j} \hat{a}^k$, we limit ourselves to observables that are ‘phase invariant’ i.e., remain unchanged under the map $\hat{a} \rightarrow e^{i\alpha} \hat{a}$, $\hat{a}^{\dagger} \rightarrow e^{-i\alpha} \hat{a}^{\dagger}$ for all α . These are of the simple form $\hat{A} = \sum_{j=0}^{\infty} c_j \hat{a}^{\dagger j} \hat{a}^j$, involving a single sum, and are just functions of the number operator $N_a = \hat{a}^{\dagger} \hat{a}$.

We can now ask for the way in which nonclassicality can manifest itself if only measurements of such operators are carried out. Clearly these expectation values involve only the phase-averaged information contained in $\phi(z_a)$:

$$\text{Tr}(\rho_a \hat{A}) = \int_0^{\infty} dI_a P(I_a) \left(\sum_{j=0}^{\infty} c_j I_a^j \right),$$

$$P(I_a) = \int_0^{2\pi} \frac{d\theta}{2\pi} \phi(I_a^{1/2} e^{i\theta}), \quad \int_0^{\infty} dI_a P(I_a) = 1.$$

While $P(I_a)$ is of course real, it may not be pointwise nonnegative. For a given $P(I_a)$, the probabilities constituting the photon number distribution (PND) are

$$p(n_a) \equiv \langle n_a | \rho_a | n_a \rangle = \int_0^{\infty} dI_a P(I_a) \frac{e^{-I_a} (I_a)^{n_a}}{n_a!}. \quad (6)$$

Whatever the nature of $P(I_a)$ may be, these probabilities are real nonnegative and add up to unity.

We now present the following definition which will suffice for our purposes[9]:

$$\rho_a \text{ is classical in phase invariant sense} \Leftrightarrow P(I_a) \geq 0. \quad (7)$$

Otherwise ρ_a is ‘nonclassical’ in the phase invariant sense. This definition is a coarse-grained version of the earlier

definition (5) given in terms of $\phi(z_a)$. Namely for the three mutually exclusive situations

$$\begin{aligned} (i) \quad & \phi(z_a) \geq 0 \quad (\text{hence } P(I_a) \geq 0), \\ (ii) \quad & \phi(z_a) \not\geq 0 \quad \text{but } P(I_a) \geq 0, \\ (iii) \quad & P(I_a) \not\geq 0 \quad (\text{hence } \phi(z_a) \not\geq 0), \end{aligned} \quad (8)$$

while definition (5) would describe (i) alone as ‘classical’ and (ii), (iii) as two levels of ‘nonclassicality’, the phase-invariant definition (7) clubs (i) and (ii) together as ‘classical’ and only (iii) as ‘nonclassical’.

One very familiar and well known signature of phase invariant nonclassicality is the antibunching condition

$$\begin{aligned} \langle \Delta n_a \rangle^2 - \langle n_a \rangle &\equiv \langle n_a^2 \rangle - \langle n_a \rangle^2 - \langle n_a \rangle \\ &\equiv \int_0^{\infty} dI_a P(I_a) (I_a - \langle I_a \rangle)^2 \equiv (\Delta I_a)^2 < 0, \end{aligned} \quad (9)$$

which definitely implies $P(I_a) \not\geq 0$. But this is by no means the only such signature. A complete set of necessary and sufficient conditions expressing the content of classicality, i.e., $P(I_a) \geq 0$, in terms of PND $\{p(n_a)\}$ will be described later in this Letter..

Transpose Operation and Expectation Values for Operators: While transpose operation on the density operator transcribes into ‘momentum reversal’ in the Wigner phase space description[2], it is more convenient for our present purposes to transfer this operation directly onto operators and their expectation values. The transpose ρ_a^T of the density operator ρ_a (of the a-mode) is defined in the Fock basis through $\langle n | \rho_a^T | n' \rangle \equiv \langle n' | \rho_a | n \rangle$. Now the key point here is that in the Fock basis the operators \hat{a} , \hat{a}^{\dagger} are represented by *real matrices*, so the transposition operation for the matrix of $\hat{a}^{\dagger l} \hat{a}^m$ coincides with hermitian conjugation, and we have $\text{Tr}(\rho_a^T \hat{a}^{\dagger j} \hat{a}^k) = \text{Tr}(\rho_a \hat{a}^{\dagger k} \hat{a}^j)$. A linear combination $\sum_{j,k} c_{j,k} \hat{a}^{\dagger j} \hat{a}^k$, with complex coefficients $\{c_{j,k}\}$, would thus be taken under the transpose map to $\sum_{j,k} c_{j,k} \hat{a}^{\dagger k} \hat{a}^j$, in contradistinction to (the antilinear) hermitian conjugation which would have taken it to $\sum_{j,k} c_{j,k}^* \hat{a}^{\dagger k} \hat{a}^j$. It is the *linearity* of the transpose map that permits its implementation on a subsystem, the b-mode, of a two-mode system through

$$\text{Tr}(\tilde{\rho}_{\text{out}} \hat{a}^{\dagger j} \hat{a}^k \hat{b}^{\dagger l} \hat{b}^m) = \text{Tr}(\rho_{\text{out}} \hat{a}^{\dagger j} \hat{a}^k \hat{b}^{\dagger m} \hat{b}^l). \quad (10)$$

This partial transpose relation proves very useful.

Beam Splitter and Conversion of Nonclassicality into Entanglement: The beam splitter is a passive, energy conserving system. Its unitary action U on the two-mode Hilbert space $\mathcal{H}^a \otimes \mathcal{H}^b$ can be viewed in two equivalent ways, and we will make use of both in succession. In the Heisenberg-type view, the effect of the beam splitter is to produce a $\pi/4$ rotation in the (\hat{a}, \hat{b}) plane:

$$\begin{aligned} U \hat{a} U^{-1} &= \frac{1}{\sqrt{2}} (\hat{a} + \hat{b}), & U \hat{b} U^{-1} &= \frac{1}{\sqrt{2}} (\hat{b} - \hat{a}), \\ U^{-1} \hat{a} U &= \frac{1}{\sqrt{2}} (\hat{a} - \hat{b}), & U^{-1} \hat{b} U &= \frac{1}{\sqrt{2}} (\hat{b} + \hat{a}). \end{aligned} \quad (11)$$

Alternatively, in the Schrödinger-type view, wherein U acts on the state rather than on dynamical variables, the fact that U commutes with the operator $\hat{a}^\dagger \hat{a} + \hat{b}^\dagger \hat{b}$ and hence will not connect subspaces of the two-mode Hilbert space $\mathcal{H}^a \otimes \mathcal{H}^b$ differing in the total number of photons will prove very useful[13].

We will now employ the beam splitter in a simple situation, and later extend its use to a less restricted situation.

1. *The case of antibunched input:* Let us assume that our beam splitter is fed with a product state

$$\rho_{\text{in}} = \rho_a \otimes |0\rangle_{bb}\langle 0|, \quad (12)$$

with the b-mode in the vacuum state. After passing through the beam splitter we have the output state $\rho_{\text{out}} = U\rho_{\text{in}}U^{-1}$. To test the operator $\tilde{\rho}_{\text{out}}$, the partial transpose of ρ_{out} , for positivity we use the first strategy (3), making the choice $A = c_0 + c_1 \hat{a}\hat{b}$.

Arranging c_1, c_2 into a column C so that C^\dagger is the row vector (c_1^*, c_2^*) , we obtain for the left hand side of (3)

$$\text{Tr}(\tilde{\rho}_{\text{out}} A^\dagger A) = C^\dagger \begin{pmatrix} 1 & \text{Tr}(\tilde{\rho}_{\text{out}} \hat{a}\hat{b}) \\ \text{Tr}(\tilde{\rho}_{\text{out}} \hat{a}^\dagger \hat{b}^\dagger) & \text{Tr}(\tilde{\rho}_{\text{out}} \hat{a}^\dagger \hat{a} \hat{b}^\dagger \hat{b}) \end{pmatrix} C. \quad (13)$$

We now use the result (10) of the PT operation and eqs. (11), to relate these traces to corresponding expectation values in the original input state ρ_a of eq. (12):

$$\begin{aligned} \text{Tr}(\tilde{\rho}_{\text{out}} \hat{a}\hat{b}) &= \text{Tr}(\rho_{\text{out}} \hat{a}\hat{b}^\dagger) \\ &= \text{Tr}(\rho_{\text{in}} U^{-1} \hat{b}^\dagger \hat{a} U) \\ &= \frac{1}{2} \text{Tr}(\rho_{\text{in}} (\hat{b}^\dagger + \hat{a}^\dagger)(\hat{a} - \hat{b})) \\ &= \frac{1}{2} \text{Tr}(\rho_a \hat{a}^\dagger \hat{a}) = \frac{1}{2} \langle n_a \rangle. \end{aligned}$$

The other independent matrix element in (13) is, by (10) followed by (11),

$$\begin{aligned} \text{Tr}(\tilde{\rho}_{\text{out}} \hat{a}^\dagger \hat{a} \hat{b}^\dagger \hat{b}) &= \text{Tr}(\rho_{\text{out}} \hat{a}^\dagger \hat{a} \hat{b}^\dagger \hat{b}) \\ &= \text{Tr}(\rho_{\text{in}} U^{-1} \hat{a}^\dagger \hat{b}^\dagger \hat{a} \hat{b} U) \\ &= \frac{1}{4} \text{Tr}(\rho_{\text{in}} (\hat{a}^\dagger - \hat{b}^\dagger)(\hat{a}^\dagger + \hat{b}^\dagger)(\hat{a} - \hat{b})(\hat{a} + \hat{b})) \\ &= \frac{1}{4} \text{Tr}(\rho_a \hat{a}^\dagger \hat{a} \hat{a}^\dagger \hat{a}) = \frac{1}{4} (\langle n_a^2 \rangle - \langle n_a \rangle). \end{aligned}$$

As a result, (13) reads:

$$\text{Tr}(\tilde{\rho}_{\text{out}} A^\dagger A) = C^\dagger \begin{pmatrix} 1 & \frac{1}{2} \langle n_a \rangle \\ \frac{1}{2} \langle n_a \rangle & \frac{1}{4} (\langle n_a^2 \rangle - \langle n_a \rangle) \end{pmatrix} C.$$

The determinant of this matrix is exactly the expression $\frac{1}{4}(\langle \Delta n_a^2 \rangle - \langle n_a \rangle)$ which appears in the antibunching condition (9). That is, if the input ρ_a is antibunched, then ρ_{out} is necessarily NPT. We have thus proved

Theorem 1: If a product state of the form $\rho_{\text{in}} = \rho_a \otimes |0\rangle_{bb}\langle 0|$, with antibunched ρ_a , is fed into a beam splitter, the output is definitely NPT entangled. That is,

$$\begin{aligned} (\langle \Delta n_a^2 \rangle - \langle n_a \rangle) < 0 \text{ for } \rho_a &\Rightarrow \tilde{\rho}_{\text{out}} \not\geq 0 \\ &\Rightarrow \rho_{\text{out}} \text{ is NPT.} \end{aligned} \quad (14)$$

2. *The case of general nonclassical PND input:* In the previous case we allowed the input a-mode state ρ_a to be any antibunched state. Now we retain the product form (12) for the input two mode state, but choose for ρ_a a more general phase invariant state determined completely by some given PND probabilities $\{p(n_a)\}$:

$$\rho_a(\{p(n_a)\}) \equiv \sum_{n_a=0}^{\infty} p(n_a) |n_a\rangle \langle n_a|. \quad (15)$$

The probabilities $p(n_a)$ are of course real nonnegative and normalized to unity, but are otherwise free. Given them, the problem of inverting eq (6) to obtain $P(I_a)$ is one of the classical moment problems, in fact the Stieltjes moment problem[10] as we are concerned with the variable $I_a \in [0, \infty)$. In this case $P(I_a)$ is uniquely determined by $\{p(n_a)\}$; and the condition $P(I_a) \geq 0$ translates, in a *necessary and sufficient* manner, into positivity conditions for a double hierarchy of real symmetric matrices of increasing dimensions, formed out of the probabilities $\{p(n_a)\}$. These are best described in the following form[9]:

$$\begin{aligned} q_{n_a} &= n_a! p(n_a) \\ \rho_a(\{p(n_a)\}) \text{ is } &\text{a 'classical' state} \\ &\Leftrightarrow \text{the PND } \{p(n_a)\} \text{ is 'classical'}. \\ &\Leftrightarrow P(I_a) \geq 0 \\ &\Leftrightarrow L^{(N)}, \tilde{L}^{(N)} \geq 0 \text{ for } N = 0, 1, 2, \dots \end{aligned}$$

$$\begin{aligned} L^{(N)} &= \begin{pmatrix} q_0 & q_1 & q_2 & \dots & q_N \\ q_1 & q_2 & q_3 & \dots & q_{N+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_N & q_{N+1} & q_{N+2} & \dots & q_{2N} \end{pmatrix}, \\ \tilde{L}^{(N)} &= \begin{pmatrix} q_1 & q_2 & q_3 & \dots & q_{N+1} \\ q_2 & q_3 & q_4 & \dots & q_{N+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{N+1} & q_{N+2} & q_{N+3} & \dots & q_{2N+1} \end{pmatrix}. \end{aligned} \quad (16)$$

Now pass the two-mode product state

$$\rho_{\text{in}} = \rho_a(\{p(n_a)\}) \otimes |0\rangle_{bb}\langle 0|, \quad (17)$$

through the beam splitter. The output state is

$$\begin{aligned} \rho_{\text{out}} &= U \rho_{\text{in}} U^{-1} \\ &= U \sum_{n_a=0}^{\infty} \frac{p(n_a)}{n_a!} (\hat{a}^\dagger)^{n_a} |0, 0\rangle \langle 0, 0| (\hat{a})^{n_a} U^{-1} \\ &= \sum_{n_a=0}^{\infty} \frac{p(n_a)}{2^{n_a} n_a!} (\hat{a}^\dagger + \hat{b}^\dagger)^{n_a} |0, 0\rangle \langle 0, 0| (\hat{a} + \hat{b})^{n_a} \\ &= \sum_{n_a=0}^{\infty} \frac{p(n_a) n_a!}{2^{n_a}} \sum_{r,s=0}^{n_a} \frac{|r, n_a - r\rangle \langle s, n_a - s|}{\sqrt{r!(n_a - r)!s!(n_a - s)!}}. \end{aligned}$$

The general matrix element of this density matrix is

$$\begin{aligned} \langle n_a', n_b' | \rho_{\text{out}} | n_a, n_b \rangle \\ = \delta_{n_a' + n_b', n_a + n_b} \frac{(n_a + n_b)! p(n_a + n_b)}{2^{n_a + n_b} \sqrt{n_a'! n_b'! n_a! n_b!}}. \end{aligned}$$

The matrix elements of the partial transpose $\tilde{\rho}_{\text{out}}$ are obtained simply by interchanging n_b and n_b' :

$$\begin{aligned} \langle n_a', n_b' | \tilde{\rho}_{\text{out}} | n_a, n_b \rangle \\ = \delta_{n_a' + n_b, n_a + n_b'} \frac{q_{n_a + n_b'}}{2^{n_a + n_b'} \sqrt{n_a'! n_b'! n_a! n_b!}}. \end{aligned} \quad (18)$$

Now we test for possible lack of positivity of $\tilde{\rho}_{\text{out}}$ by using the second strategy (4). Consider first the principal submatrix of (18) obtained by setting $n_b' = n_a'$, $n_b = n_a$. This submatrix $H^{(N)}$ has elements

$$H_{n_a', n_a}^{(N)} \equiv \langle n_a', n_a' | \tilde{\rho}_{\text{out}} | n_a, n_a \rangle = \frac{q_{n_a + n_a'}}{2^{n_a + n_a'} n_a'! n_a!}.$$

Clearly, this (N+1) dimensional submatrix coincides with $L^{(N)}$ modulo congruence by a diagonal matrix $D^{(N)}$:

$$\begin{aligned} H^{(N)} &= D^{(N)} L^{(N)} D^{(N)}, \\ D_{n_a, n_a'}^{(N)} &= (2^{n_a} n_a!)^{-1} \delta_{n_a, n_a'}. \end{aligned} \quad (19)$$

Consider similarly the principal submatrix of (18) corresponding to the row (and column) choices $n_b' = n_a' + 1$ and $n_b = n_a + 1$. This submatrix $\tilde{H}^{(N)}$ has elements

$$\begin{aligned} \tilde{H}_{n_a', n_a}^{(N)} &\equiv \langle n_a', n_a' + 1 | \tilde{\rho}_{\text{out}} | n_a, n_a + 1 \rangle \\ &= \frac{q_{n_a + n_a' + 1}}{2^{n_a + n_a' + 1} \sqrt{(n_a' + 1)! n_a'! (n_a + 1)! n_a!}}. \end{aligned}$$

This (N+1) dimensional submatrix is seen to coincide with $\tilde{L}^{(N)}$ modulo congruence by a diagonal matrix:

$$\begin{aligned} \tilde{H}^{(N)} &= \tilde{D}^{(N)} \tilde{L}^{(N)} \tilde{D}^{(N)} \\ \tilde{D}_{n_a, n_a'}^{(N)} &= \left(2^{n_a + \frac{1}{2}} \sqrt{(n_a + 1)! n_a!} \right)^{-1} \delta_{n_a, n_a'}. \end{aligned} \quad (20)$$

We see that in both (19) and (20), the exhibited principal submatrices of $\tilde{\rho}_{\text{out}}$ are essentially the matrices $L^{(N)}$ and $\tilde{L}^{(N)}$ (apart from trivial congruences) appearing in the result (16) of the moment problem to ensure positivity of $P(I_a)$ [9]. Thus if $\rho_a(\{p(n_a)\})$ is nonclassical, i.e., if $P(I_a) \not\geq 0$, then one of the matrices $L^{(N)}$, $\tilde{L}^{(N)}$ will necessarily fail to be positive semidefinite, for some N , as seen from (16), rendering one of the above submatrices of $\tilde{\rho}_{\text{out}}$ nonpositive. Thus, we arrive at
Theorem 2: If the input PND statistics $\{p(n_a)\}$ has any (phase-invariant) nonclassicality, then the output two-mode state emerging after the beam splitter is NPT:

$$L^{(N)} \quad \text{or} \quad \tilde{L}^{(N)} \not\geq 0 \quad \text{for some } N$$

$$\begin{aligned} &\Leftrightarrow \rho_a(\{p(n_a)\}) \text{ correspondingly 'nonclassical'} \\ &\Rightarrow \text{corresponding submatrices of } \tilde{\rho}_{\text{out}} \not\geq 0 \\ &\Rightarrow \rho_{\text{out}} \text{ is NPT entangled.} \end{aligned} \quad (21)$$

To conclude, the output nonclassicality, the transform by the beam splitter of the nonclassicality residing locally in the input a-mode photon statistics, manifests itself nonlocally as NPT entanglement. It is rather remarkable that the infinite set of matrices in (19), (20) whose positivity is equivalent to the positivity of $\tilde{\rho}_{\text{out}}$ should essentially coincide with the infinite set of matrices in (16) designed, with the help of the classical Stieltjes moment problem[10], to capture the classicality condition $P(I_a) \geq 0$.

* Electronic address: solomon@imsc.res.in

† Electronic address: nmukunda@cts.iisc.ernet.in; Permanent Address: Centre for High Energy Physics, Indian Institute of Science, Bangalore 560 012

‡ Electronic address: simon@imsc.res.in

- [1] A. Furusawa *et al.* Science **282**, 706 (1998); W. P. Bowen *et al.* Phys. Rev. Lett. **89**, 253601 (2002); F. Grosshans *et al.* Nature **421**, 238 (2003).
- [2] R. Simon, Phys. Rev. Lett. **84**, 2726 (2000); L.-M. Duan, G. Giedke, J.I. Cirac, and P. Zoller, **84**, 2722 (2000).
- [3] R. F. Werner and M. M. Wolf, Phys. Rev. Lett. **86**, 3658 (2001); G. Giedke *et al.* Phys. Rev. Lett. **91**, 107901 (2003); G. Giedke and J. I. Cirac, Phys. Rev. A **66**, 032316 (2002).
- [4] G. Giedke, B. Kraus, M. Lewenstein, and J. I. Cirac, Phys. Rev. Lett. **87**, 167904 (2001); M. M. Wolf, J. Eisert, and M. B. Plenio, Phys. Rev. Lett. **90**, 047904 (2003).
- [5] A. Biswas and G. S. Agarwal, New Journal of Physics **7**, 211 (2005); E. Shchukin and W. Vogel, Phys. Rev. Lett. **95**, 230502 (2005).
- [6] J. K. Asboth, J. Calsamiglia, and H. Ritsch, Phys. Rev. Lett. **94**, 173602 (2005).
- [7] R. E. Slusher *et al.*, Phys. Rev. Lett. **55**, 2409 (1985); Ling-An Wu *et al.*, Phys. Rev. Lett. **57**, 2520 (1986).
- [8] H. J. Carmichael and D. F. Walls, J.Phys. B **9**, 1199 (1976); H. J. Kimble, M. Dagenais, and L. Mandel, Phys. Rev. Lett. **39**, 691 (1977).
- [9] R. Simon, Mary Salvadoray, Arvind, and N. Mukunda, quant-ph/9709030; Arvind, N. Mukunda, and R. Simon, J.Phys. A **31**, 565 (1998).
- [10] See, for example, J. A. Shohat and J. D. Tamarkin, *The Problem of Moments* (American Mathematical Society, Providence, R. I., 1943).
- [11] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996); P. Horodecki, Phys.Lett.A **232**, 333 (1997).
- [12] E. C. G. Sudarshan, Phys. Rev. Lett. **10**, 277 (1963).
- [13] R. A. Campos, B. E. A. Saleh, and M. C. Teich, Phys. Rev. A **40**, 1371 (1989).