

Can one build a quantum hard drive? A no-go theorem for storing quantum information in equilibrium systems

Robert Alicki and Michał Horodecki

*Institute of Theoretical Physics and Astrophysics,
University of Gdańsk, Wita Stwosza 57, PL 80-952 Gdańsk, Poland*

(Dated: April 1, 2022)

We prove a no-go theorem for storing quantum information in equilibrium systems. Namely, quantum information cannot be stored in a system with time-independent Hamiltonian interacting with heat bath of temperature $T > 0$ during time that grows with the number of used qubits. We prove it by showing, that storing quantum information for macroscopic time would imply existence of perpetual mobile of the second kind. The general results are illustrated by the Kitaev model of quantum memory. In contrast, classical information can be stored in equilibrium states for arbitrary long times. We show how it is possible via phase-transition type phenomena.

Our result shows that there is a fundamental difference between quantum and classical information in *physical* terms.

I. INTRODUCTION

Quantum computing needs at least quantum memory. Quantum memory is based on encoding of (say 1-qubit) states into metastable states of a larger system. The states are expected to survive the interaction with a heat bath of the temperature $T > 0$. One requires that the life-time of the memory grows with the number N of subsystems. One can consider two methods: time-dependent [1, 2, 3, 4], time independent "self-correcting" systems (cf. [5]). The first method is non-equilibrium one, where energy is constantly dissipated and moreover the life-time τ satisfies $N \sim \text{poly}(\log \tau)$ which means that the memory is claimed to be exponentially stable. The results involve a kind of phase transition (cf. [3, 4]), and are mainly formulated within phenomenological approach (see however [6, 7, 8, 9]). The second method uses equilibrium states.

In this paper we consider time-independent method (time dependent case will be considered in future). We show that this method is not useful, i.e. quantum memory cannot be stored in equilibrium states. Put it into different way: one cannot build a "quantum hard drive" - device that would store quantum information without dissipation of energy constant in time. Conversely, we show how classical information can be stored in equilibrium states (this happens for all sorts of memory devices) by means of phase transition type phenomena.

To prove our no-go result use theorems from mathematical physics concerning the notion of passivity (inability of drawing work by external cyclic forces) and its stronger version complete passivity [10]. We invoke the highly nontrivial result, stating that set of completely passive states constitute simplex, so that no quantum superpositions can be encoded into such states. Thus our argument does not rely on a particular model. We refer to The Second Law, so that the result works in all the situations in which the law remains valid. This differs our results from the issues concerning decoherence of Schroedinger cat states (see e.g. [11, 12, 13]).

Our results imply that in equilibrium one can store

quantum information only over microscopic periods of time. Note however, that these microscopic time scale may vary from system to system, depending on the type of environment to which the system couples and the coupling strength, and for some systems may be pretty long.

II. CLASSICAL MEMORY AND PHASE TRANSITIONS

We will show on a simple model, how classical information can be efficiently stored. We will use mean-field Curie-Weiss model of ferromagnetic. Though unphysical, it reflects very well all the essential features that lead to efficient storage of classical information. It is well known that this model exhibits phase transition. We will show now that the phase transition gives rise to exponentially stable states. We consider N spins interacting by means of Hamiltonian

$$H = -JNX^2 \quad (1)$$

where

$$X = \frac{1}{N} \sum_j \sigma_i \quad (2)$$

where σ_j is Pauli matrix σ_z acting on j -th spin, and summation is taken over all pairs of sites. Thus, due to interaction, the energetically favourable configurations are when spins point the same direction. For a fixed value of the mean magnetization X , denoted by x , the energy amounts to

$$E(x) = -JNx^2. \quad (3)$$

We see that there is energy barrier between the two configuration minimizing the energy.

However, not only the height of the barrier is relevant, but also the number of microstates for a fixed macrostate ρ_x . Crossing of the barrier may be likely, if the number of states on the top of it is large enough. The quantity that




FIG. 1: Brownian particle in double potential well.

takes into account both energy difference and number of states is a free energy. Let us consider it in more detail. Basic processes induced by noise relevant for our problem are flips of single spin. Single spin-flip causes flip between neighbouring x 's changing $x \rightarrow x \pm \frac{1}{N}$. Thus the whole process is a random walk on x line. The transition probabilities between macrostates ρ_x are determined by transition probabilities between microstates. To avoid combinatorial issues, we adopt the heuristic rule

$$\frac{p(x \rightarrow x')}{p(x' \rightarrow x)} \simeq e^{-(E(x')-E(x))/k_B T} \frac{\mathcal{N}(x')}{\mathcal{N}(x)} \quad (4)$$

where $\mathcal{N}(x) = \exp[Nh(\frac{x+1}{2})]$ denotes number of microstates, $h(x) = -x \ln x - (1-x) \ln(1-x)$. Since the free energy is given just by $F(x, T) = E(x) - TS(x)$ with $S(x) = Nk_B h(x)$ we obtain

$$\frac{p(x \rightarrow x')}{p(x' \rightarrow x)} \simeq e^{-(F(x', T) - F(x, T))/k_B T} \quad (5)$$

The terms E and TS in free energy have here clear interpretation: the first one gives rise to the Boltzmann factor, while the second one reports the contribution of number of states to transition probability. Thus it is now clear that it is the shape of free energy rather than energy itself, that will determine stability. To see it more in detail, we consider x as continuous variable. In such limit, the problem is equivalent to Brownian motion in the potential V given by free energy (see [14]).

We can thus apply Kramers formula for mean exit time from a potential well (see fig. 1)

$$t(a \rightarrow b) \simeq \frac{2\pi k_B T}{D} |V''(a)V''(b)|^{-\frac{1}{2}} e^{\frac{1}{2}(V(b)-V(a))/k_B T} \quad (6)$$

where D is diffusion constant [14]. Thus, in our case we obtain, that probability of flip is exponential in the difference of free energy between x_{eq} and $x = 0$ (the top of energy barrier). Thus we have to examine the shape of free energy, explicitly given by

$$F(x, T) = N(-kT h(\frac{x+1}{2}) - Jx^2) \quad (7)$$

The function has two minima for $T < T_c$ and one minimum for $T > T_c$ (where T_c is critical temperature). When the entropic terms dominates, the density of final states is so large, that the Boltzmann factor cannot prevent from flip between the two states, and we have single minimum. There is only one stable state, with $x = 0$ (note that this is the state of the largest energy).

If the energy part dominates, then the probability of flipping is exponentially suppressed, and we have two minima divided by barrier proportional to N . Thus we have two stable phases, and according to (6) probability of flipping between them is exponentially decreasing with N .

The presented above mechanism is quite universal. Classical information can be efficiently stored in metastable states corresponding to local minima of the free energy separated by barriers which grow with the size of the system (e.g. large molecules, mesoscopic systems). For example the structure of protein molecules is determined by analysing free-energy landscape [15, 16].

III. EQUILIBRIUM STATES AND PASSIVITY

A spatially confined quantum system weakly interacting with a heat bath at the temperature T tends to the unique equilibrium state represented by the Gibbs density matrix

$$\rho_\beta = \frac{e^{-\beta H}}{\text{Tr} e^{-\beta H}} \quad (8)$$

where H is a Hamiltonian of the system and $\beta = 1/kT$ is the inverse temperature. This behavior illustrates the Zeroth Law of Thermodynamics and can be rigorously proved for simple but generic models of reservoirs both in the weak coupling limit [17] and for finite, but small enough coupling constant [18]. The only essential assumption is a kind of irreducibility of the coupling between the system and the bath which eliminates the appearance of conserved observables impairing equilibration process. This condition can be seen as the absence of decoherence free subspaces and subsystems in the finite system.

We are interested in the relaxation processes in the case of a system composed of many subsystems (say N qubits). For noninteracting subsystems and local coupling to the bath we observe individual and independent relaxation processes which implies the thermalization time independent of the number N of subsystem. On the other hand, for strongly interacting subsystems collective phenomena can produce metastable states with life-times growing with N . We observed such phenomenon for classical systems in the Section II and discussed their applications to classical information storage.

In principle, similar metastable states which become stationary, and resistant to local perturbations for $N \rightarrow \infty$ might be used to store quantum information as well.

Therefore, it is crucial to analyse mathematical structure of such quantum states. They should satisfy the restrictions imposed by The Second Law of Thermodynamics. Let us recall Kelvin formulation of the law:

It is impossible to construct an engine which, operating in a cycle will produce no other effect than the extraction of heat from a reservoir and the performance of an equivalent amount of work.

This can be rephrased in terms of *passivity*. We say that a state ρ is passive (with respect to the dynamics \mathcal{U}_t), when it is impossible to extract the energy from the system at a given state by means of a cyclic process (Second Law of Thermodynamics). A state is n -passive, if the state $\rho^{\otimes n}$ with respect to product evolution $\mathcal{U}_t^{\otimes n}$ is still passive. Finally, a state is called completely passive (CP), if it is n -passive for all n . Thus if a state is not CP, then one can extract energy from a finite amount of copies.

Let us now recall how passivity and complete passivity is described within quantum mechanics [10]. The cyclic process is represented by time dependent perturbation $h(t)$ satisfying $h(0) = h(\tau) = 0$. Consider first a finite system described by the Hamiltonian H with the eigenvectors $|j, \mu\rangle$ and the possibly degenerated eigenvalues ϵ_j

$$H|j, \mu\rangle = \epsilon_j|j, \mu\rangle, \quad \mu \in I_j. \quad (9)$$

The energy change due to a time dependent perturbation $h(t)$ is given by $\Delta E = \text{Tr}(\rho(U_\tau H U_\tau^{-1} - H))$ where $U_\tau = \mathbf{T} \exp \int_0^\tau dt (H + h(t))$. A state ρ is *passive* if $\Delta E \geq 0$ for all h . One easily shows that ρ is passive if and only if

- (i) $[H, \rho] = 0$ and hence for every passive ρ one can choose $|j, \mu\rangle$ in such a way that $\rho = \sum_{j, \mu} \lambda(j, \mu) |j, \mu\rangle \langle j, \mu|$,
- (ii) $(\epsilon_j - \epsilon_k)(\lambda(j, \mu) - \lambda(k, \nu)) \leq 0$.

Thus the state is passive if and only if it commutes with the Hamiltonian, and there is no inversion of population, in the sense, that for any two energy levels, the upper level is not more populated than the lower one. This still leaves freedom on degeneracies of Hamiltonian. Following [10] it is not hard to find that if a state is not Gibbs one, then for some n the state $\rho^{\otimes n}$ will get inversion of population with respect to sum of single system Hamiltonians $\sum_i H_i$. Therefore, for finite systems completely passive states are Gibbs states, hence at the given temperature we have only a single completely passive state.

For the infinite systems the situation is less trivial, we can have many CP states (e.g. different thermodynamical phases) which are the limits of metastable states (with increasing life-times) of the corresponding sequence of finite systems. Nevertheless, the structure of CP states is determined by the following theorem [28] valid for any system described by a C^* algebra of observables \mathcal{A} and a group of automorphisms (Hamiltonian dynamics) \mathcal{U}_t .

Theorem 1 *For arbitrary system $(\mathcal{A}, \mathcal{U}_t)$ the set of CP states constitutes a simplex.*

This means that any CP state can be uniquely decomposed into extremal states of the set of all CP states.

IV. NO-GO THEOREM FOR STABLE QUANTUM MEMORY

Suppose, that we want to construct stable quantum memory by means of a quantum system consisting of N subsystems with a specially designed Hamiltonian. We assume that the system interacts with a heat bath of the temperature $T > 0$.

Suppose, now that by increasing N we can arbitrarily increase the life-time of quantum memory. In particular, we can make the time much longer than any microscopic time scale, so that the violations of the Second Law due to fluctuations will be suppressed. Consider the set of states that can survive for macroscopic time. We call them metastable states. They have to be completely passive, as otherwise we could build *perpetuum mobile of the second kind*. Indeed, as explained in the previous section, if a system is in a non-CP state, then n systems constitute a system in a nonpassive state, for some finite n . Putting now k of such n -tuples into heat bath, we obtain a system in equilibrium, from which one can draw work proportional to k [29].

Thus employing the Second Law, we obtain that the only states that can be used to store information for arbitrarily long time are CP states [30]. All other states must decay within microscopic time scales.

However, according to Theorem 1, CP states form a simplex, hence the set of metastable states possesses a fully classical representation. They can be treated as probability measures on a certain configuration space. Therefore CP states cannot be used for the faithful representation of the quantum states. It follows for example from the fact, that quantum bits cannot be faithfully conveyed by classical channel (see e.g. [19, 20]). As a consequence only classical information can be preserved over macroscopic time scales.

V. EXAMPLE: KITAEV'S MODEL

In the following we would like to discuss the model introduced by Kitaev [5] which is supposed to be a good candidate for a self-correcting quantum memory.

Kitaev considered $k \times k$ square lattice torus. On each edge there is a qubit (so that there are $n = 2k^2$ qubits). For each vertex s and face p one defines operators

$$A_s = \sum_{j \in \text{star}(s)} \sigma_j^x, \quad B_p = \sum_{j \in \text{boundary}(p)} \sigma_j^z, \quad (10)$$

where "star" denotes edges that touch vertex s and "boundary" denotes edges that surround face p . The

Hamiltonian is given by

$$H = - \sum_s A_s - \sum_p B_p \quad (11)$$

The operators A_s and B_p are dichotomic, and they all mutually commute. We can form basis consisting of eigenvectors of those operators. One finds, that for any basis vector, it has eigenvalue 1 for even number of operators A_s and also for even number of operators B_p . Thus the energy levels are given by the number of pairs of operators A_s and B_p to which there is eigenvalue 1. This can be interpreted as number of pairs of particles. For each level, there are many possible configurations of pairs. However for every configuration there is 4-dimensional degeneracy due to topology of the torus. Thus we can imagine the total system as tensor product of 4-dimensional space and the space determined by configurations of pairs of particles. The particles are (non-interacting) anyons: they exhibit non-standard statistics, which manifests by inducing phase, when one anyon winds around the second one. It is the 2-qubit subsystem is the one that is expected to be noiseless, when the size of torus becomes large [21].

In this section we will consider the system from two points of view. First, we will treat it as infinite system, and discuss in the context of our general result. Subsequently, we shall consider finite system, and analyse interaction with a heat bath. This will independently prove that in finite temperature, quantum information cannot be stored.

A. Infinite system picture

Let us see how Kitaev's model fits into our theorem. To this end we will consider the limiting case of infinite system. The infinite system of spins is defined in terms of *quasilocal algebra*, spanned by observables that are tensor products of only finite amount of single site observables such as $\dots \otimes I \otimes I \otimes A_1 \otimes \dots \otimes A_k \otimes I \otimes I \otimes \dots$ and closed in operator norm topology [22]. Now, one can show that the limits of all ground (vacuum) states in the Kitaev model produce the same expectation values for all local and hence also quasilocal observables. This could be expected, as *good codewords* should be indistinguishable by local measurements. Therefore the infinite Kitaev system possesses a single ground state and therefore there is no phase transition in this model and even a bit of classical information cannot be stored [31].

The physical meaning of this result is the following. The highly nonlocal observables which describe the encoded two qubits become, with increasing size of the system, more and more vulnerable to external perturbations and hence less and less accessible to meaningful measurements. We will see in the next section, that indeed the noise will wash out all the quantum information in time that for sure does not increase with the size of the system.

B. Finite system interacting with heat bath

Let us then consider the Kitaev system coupled to a heat bath via local operators, for example σ_x^i and σ_z^j . We will work in Markovian approximation (weak coupling limit).

The basic processes that can occur is i) creation of pair of neighboring anyons, ii) annihilation of such pair, and ii) move of an anyon. For example, when we hit a particular qubit with σ_x , then, if the qubit belong to pair of anyons of type B , then the pair will be annihilated. If there were no anyons at this place, a pair will be created. If there was one anyon, it will be moved. The amplitude of the basic processes is determined by spectral density of the noise, always however we have $p_c = e^{-\Delta E/k_B T} p_a$, where ΔE is the energy needed to create a pair, p_c and p_a are probabilities of creation, and annihilation of a pair of anyons respectively. In this way in equilibrium state there is a constant density of anyons.

One of the decoherence mechanisms is a random walk of the created anyon pairs which will be performed until the anyons meet. It is clearly seen, that such processes will equilibrate the system of anyonic configurations. What about the topological qubits where the quantum information is to be stored? Consider such a process: an anyon pair is created, one of them winds around torus, making a noncontractible loop, and they finally annihilate. Such a process performs a gate on one of the qubits (there are two kinds of noncontractible loops, and this gives gates on different qubits) [5]. Thus such process corresponds to error, and if it is not suppressed, we cannot store quantum information on topological qubits.

We will now argue that quantum memory will be spoiled at least with the probability independent of the size of the system (number of qubits), so that enlarging a system will not bring any improvement.

Let us concentrate for a while on a single pair that has been created. Without loss of generality, we can imagine, that one anyon is resting, while the other one is performing random walk.

It is known that in a random walk on plane, the particle comes back to the origin infinitely many times. In our case, we do not have usual random walk, because coming back to the origin means annihilating the pair, and the walk is ended. If the path was short, then the process does not affect quantum memory. Thus we want to estimate the probability that a long path occur. Since the process is Markovian (i.e. memoryless) and the average number of anyons is constant, we can imagine, that the walk of the just annihilated pair is *continued* by a new pair that has been just created (see Fig. 2) In this way we have reduced the problem to the usual random walk.

It is well known, that probability of getting away from a disk of the radius L is proportional to $1/L^2$ and probability of a single step p . This follows e.g. from the fact that random walk is a discrete version of diffusion of Brownian particle. On the other hand, the number




FIG. 2: Random walk of anyons: the walk of one pair of anyons (white ones) after annihilation is continued by another pair of newly created anyons (black ones).

of pairs is again proportional to L^2 (the proportionality constant depending on temperature). Thus the overall probability p_{long} of making a path with a length of the order of a size of the system is proportional to p , and does not depend on the size itself [32].

$$p_{long} \propto \frac{p}{L^2} \times L^2 \quad (12)$$

This reasoning can be performed both for σ_x and σ_z noise operators. And existence of long paths will imply that both kind of nontrivial paths will be performed with constant probability. Thus both topological qubits will

be subjected to random X and Z gates, which means that their state will tend exponentially to the maximally mixed state. Therefore not only the quantum information, but also the classical one will be washed out. Hence also we will not have phase transition as predicted for infinite system with a quasi-local algebra. Thus we have shown, that Kitaev's system cannot provide us the means of storing quantum information, in accordance with our general theorem.

In conclusion, our result reveals a new, fundamental feature of quantum information: the latter cannot be stored in equilibrium systems. For practical matter, it implies, that it is impossible to construct quantum memories that do not require active protection of information and are stable for longer than microscopic time scales. On more fundamental level, our result shows for the first time that there is basic difference between quantum and classical information not only on logical level (as implied e.g. by the no-cloning theorem [23, 24, 25]) but also on physical level.

It is a pleasure to thank Paweł Horodecki and Ryszard Horodecki for several years of numerous discussions on decoherence problems in quantum computers as well as for useful discussion concerning the present work. R.A. is also grateful to Daniel Lidar, Dorit Aharonov, Michael Ben-Or and Gil Kalai for discussions on fault tolerance. M.H. acknowledges discussions on the same topic with Charles Bennett, Masato Koashi, Debbie Leung, Noah Linden, Raymond Laflamme, John Preskill and Alexiey Kitaev.

This work is supported by Polish Ministry of Science and Education, grant PBZ-MIN-008/P03/2003 and EC IP SCALA.

-
- [1] D. Aharonov and M. Ben-Or, quant-ph/9910081.
 - [2] E. Knill, R. Laflamme, and W. H. Żurek, Science **279**, 342 (1998).
 - [3] E. Dennis, A. Kitaev, A. Landahl, and J. Preskill, J. Math. Phys. **43**, 4452 (2002), quant-ph/0110143.
 - [4] D. Aharonov, quant-ph/9910081.
 - [5] A. Y. Kitaev, Annals Phys. **303**, 2 (2003), quant-ph/9707021.
 - [6] R. Alicki, M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. A **65** (2001), quant-ph/0105115.
 - [7] B. M. Terhal and G. Burkard, Phys. Rev. A **71**, 012336 (2005), quant-ph/0402104.
 - [8] R. Alicki (2004), quant-ph/0402139.
 - [9] D. Aharonov, A. Kitaev, and J. Preskill, Phys. Rev. Lett **96**, 050504 (2006), quant-ph/0510231.
 - [10] W. Pusz and S. L. Woronowicz, Commun. Math. Phys. **58**, 273 (1978).
 - [11] E. B. Davies, Ann. Inst. H. Poincaré A **28**, 91 (1978).
 - [12] E. Joos and H. Zeh, Z. Phys. B **59**, 223 (1985).
 - [13] W. H. Żurek, Phys. Rev. D **24**, 1516 (1981).
 - [14] C. W. Gardiner, *Handbook of Stochastic Methods for Physics, Chemistry and the Natural Sciences* (Springer-Verlag, 1983).
 - [15] A. Liwo, M. Khalili, and H. A. Scheraga, PNAS **102**, 2362 (2005).
 - [16] S. Oldziej *et al.*, PNAS **102**, 7547 (2005).
 - [17] E. B. Davies, Commun. Math. Phys. **39**, 91 (1974).
 - [18] J. Dereziński and V. Jaksic, Ann. H. Poincaré **4**, 739 (2003).
 - [19] S. Popescu, Phys. Rev. Lett **72**, 797 (1994).
 - [20] G. Alber, T. Beth, M. Horodecki, P. Horodecki, R. Horodecki, M. Rötteler, H. Weinfurter, R. Werner, and A. Zeilinger, *Quantum Information: An Introduction to Basic Theoretical Concepts and Experiments* (Springer, 2001).
 - [21] E. Joos and H. Zeh, Phys. Rev. Lett **90**, 067902 (2003), quant-ph/0208132.
 - [22] D. W. Robinson and O. Bratteli, *Operator algebras and quantum statistical mechanics 2* (Springer Verlag, Berlin, 1997).
 - [23] E. P. Wigner, *Symmetries and Reflections* (Indiana University Press, Bloomington, 1967).

- [24] W. K. Wootters and W. H. Zurek, Nature **299**, 802 (1982).
- [25] D. Dieks, Phys. Lett. A **92**, 27 (1982).
- [26] R. Alicki, Physica A **150**, 455 (1988).
- [27] P. Zanardi and M. Rasetti, Phys. Rev. Lett **79**, 3306 (1997).
- [28] The theorem implied by the following two seminal results: (i) for arbitrary system $(\mathcal{A}, \mathcal{U}_t)$ at temperature $T > 0$ the CP states are exactly so called KMS states (generalizations of Gibbs state to include infinite systems) [10]; (ii) for arbitrary system $(\mathcal{A}, \mathcal{U}_t)$ at temperature $T > 0$ the KMS states constitute simplex [22].
- [29] Note, that since we requiring complete passivity we have to work in the limit of infinite N , as for finite N only Gibbs state is CP.
- [30] In particular, this rules out systems with so called decoherence free subspaces [26, 27]. This can be explained by the fact, that conditions to obtain decoherence free subspaces are pretty unstable.
- [31] The rigorous analysis of the infinite Kitaev model will be published elsewhere
- [32] One can expect that another process of *fusion* of two anyonic strings into a single one will be even more efficient and yield the decoherence rate of encoded qubits increasing with the size of the system