

Untouched Aspects of the Wave Mechanics of a Particle in One Dimensional Box

Yatendra S. Jain

Department of Physics
North-Eastern Hill University Shillong - 793 022, India

Abstract

Wave mechanics of a particle in 1-D box (size = d) is critically analyzed to reveal its untouched aspects. When the particle rests in its ground state, its zero-point force (F_o) produces non-zero strain by modifying the box size from d to $d' = d + \Delta d$ in all practical situations where the force (F_a) restoring d is not infinitely strong. Assuming that F_a originates from a potential $\propto x^2$ (x being a small change in d), we find that: (i) the particle and strained box assume a mutually bound state (under the equilibrium between F_o and F_a) with binding energy $\Delta E = -\epsilon'_o \Delta d/d'$ (with $\epsilon'_o = h^2/8md'^2$ being the ground state energy of the particle in the strained box), (ii) the box size oscillates around d' when the said equilibrium is disturbed, (iii) an exchange of energy between the particle and the strained box occurs during such oscillations, and (iv) the particle, having collisional motion in its excited states, assumes collisionless motion in its ground state. These aspects have desired experimental support and proven relevance for understanding the physics of widely different systems such as quantum dots, quantum wires, trapped single particle/ion, clusters of particles, superconductors, superfluids, *etc.* It is emphasized that the physics of such a system in its low energy states can be truly revealed if the theory incorporates F_o and related aspects.

Keywords : wave mechanics, single particle, 1-D box.

PACS : 03.65-w, 03.65-Ca, 03.65-Ge

Email ysjain@email.com

1. Introduction

Wave mechanics emerged as a general theory of all natural phenomena about eight decades back. While it is believed to have been successful in explaining the observed behavior of widely different systems under widely different physical conditions, its several basic aspects and their interpretations still form a subject of debate [1-4]. This reveals that answer to several philosophical questions related to the basic principles of wave mechanics are unclear and the intricacies of wave particle duality need detailed discussion for their better understanding. In this context one may consider even the wave mechanics of a particle of mass m in a 1-D box of size d (*the simplest possible quantum system used to introduce the subject at its elementary level* [5]) which reveals the following.

1.1 Energy eigenfunction Ψ_n of n-th quantum state

$$\Psi_n = \sqrt{\frac{2}{d}} \sin(q_n x) \quad (1)$$

with $q_n = n\pi/d$, $n = 1, 2, 3, \dots$, energy eigenvalue,

$$E_n = \langle \Psi_n | -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} | \Psi_n \rangle = \frac{n^2 \hbar^2}{8md^2}, \quad (2)$$

with \hbar = Plank constant, and position expectation of the particle

$$\langle x \rangle = \langle \Psi_n | x | \Psi_n \rangle = \frac{d}{2} \quad (3)$$

1.2 Ψ_n , representing a kind of *stationary matter wave*, modulates the configuration (x) and the phase ($\phi_n = q_n x$) positions of the particle. It has n anti-nodal regions of size $\lambda_n/2 = d/n$ and $n-1$ nodes (*excluding two nodes at the two walls of the box*) and spreads over the entire size (d) of the box. It is an eigenfunction of q^2 operator ($-\partial_x^2$) not of q ($-i\partial_x$) which means that an experiment on the particle can determine only the magnitude (not the direction) of q . The fact, that Ψ_n is a result of the superposition of two plane waves, $u' = \exp(iq_n x)$ and $u'' = \exp(-iq_n x)$ of momenta q_n and $-q_n$, implies that there can be no means that may fix \pm direction for q_n in such a state of the particle.

1.3 The particle in its n -th quantum state exerts a *real* force [6]

$$f_n = -\partial_d \varepsilon_n = \frac{n^2 \hbar^2}{4md^3} \quad (4)$$

on the walls of the box that tends to increase d .

While these results seem to provide full understanding of the wave mechanics of the system, however, motivated by our recent analysis of the wave mechanics of two hard core (HC) particles in 1-D box [7] we analyzed the wave mechanics of the titled system further and discovered its untouched aspects, particularly, related to the role of quantum size, zero-point energy, zero-point force, *etc.* Interestingly, these aspects are found to have desired experimental foundations and proven relevance for understanding of the true behavior of widely different physical systems such as quantum dots, quantum wires, trapped ions [8, 9], metallic clusters [10, 11] and even complex systems such as superconductors and superfluids [12, 13]. In this context we may, particularly, refer to our recent papers related to the microscopic theory of superconductivity [14] and the unification of the physics of widely different systems interacting bosons and fermions including low dimensional systems such as N HC quantum particles in 1-D box [15]. As such this study would benefit the learners of wave mechanics and many body physics.

2. Untouched Aspects

1.1 Strain in the box:

We note that f_n (Eqn. 4) can also be obtained from

$$f_n = 2\hbar q_n \frac{\hbar q_n / m}{2d} = \frac{n^2 \hbar^2}{4md^3} \quad (5)$$

with $2\hbar q_n$ = change in momentum of the particle during its collision with a wall of the box and $(\hbar q_n/m)/(2d)$ = frequency of collision. This indicates that: (i) the particle between the walls, moving with group velocity $v_g = \partial E_n / \partial p_n = (\hbar q_n/m)$, periodically collides with each wall at a frequency $(\hbar q_n/m)/(2d)$, and (ii) f_n has an apparent identity with the force that a gas particle (as a classical entity) exerts on a wall of its container and contributes to the gas pressure which tends to inflate the size of the container. Evidently, this is an interesting aspect of the particle dynamics in a 1-D box. In variance with what one learns from classical mechanics, wave mechanics concludes E_n to depend on d and E_1 (lowest possible allowed energy) to have non-zero value which have an interesting impact on the behavior of our system which can be demonstrated by a thought experiment where the system is kept in contact with a thermal bath whose temperature (T) is slowly reduced to zero. Since the probability for the particle to occupy n -th quantum state goes proportionally with $\exp[-(E_n - E_1)/k_B T] = \exp[-(n^2 - 1)\varepsilon_o/k_B T]$ (with $E_1 = \hbar^2/8md^2 = \varepsilon_o$), it is evident that such probability even for the first excited state ($n=2$) would be an order of magnitude smaller than that for the ground state ($n=1$) when bath T is $\approx T_o$ (the temperature equivalent of ε_o). Evidently, to a good approximation, the particle at all $T \leq T_o$, stays in its ground state. Here we note that: (i) the particle in $n > 1$ state can have lower energy, not only by inflating d , but also by following an alternative path (where increase in d is not necessary) of jumping to a lower energy state and (ii) $f_{n>1}$ decreases in its strength when the particle moves to lower energy state with decreasing T . Evidently, the inflation in d (if any) produced by f_n would decrease with decreasing T and the box would exhibit $+ve$ thermal expansion coefficient $(1/d)\partial_T d$. However, the situation changes when the particle rests in its ground state ($n = 1$) because it can have lower energy only if d gets increased. Naturally $f_1 (= \hbar^2/4md^3)$, which can be called as *zero-point force* (hereafter denoted by F_o), can produce non-zero strain ($+\Delta d$ change in d) with T decreasing below $T \approx T_o$ when it reaches in equilibrium with forces (F_a) that restore d . It is evident that F_o and $+\Delta d$ would remain unchanged at all $T \leq T_o$ and the system would have $-ve$ $(1/d)\partial_T d$ around T_o . The observation of $-ve$ $(1/d)\partial_T d$ of the system at T around T_o should obviously prove the presence of non-zero strain in the box and the fall of the particle in its ground state.

2.2 Bound state of Particle and strained box

We note that Eqns. 1-3 assume that the force F_a , which controls the box size, is infinitely strong for which d remains unchanged in spite of the inflating action of f_n . However, since F_a is not infinitely strong in all practical situations, d is expected to change (say, by Δd) in the state of equilibrium between F_a and f_n . Assuming that F_a can be derived from $V(x) = (k/2)x^2$ potential, the total energy of the system (particle in its ground state + strained box) can be expressed as

$$E = \frac{\hbar^2}{8m(d+x)^2} + \frac{1}{2}kx^2 \quad (6)$$

with x being an increase in d . Solving Eqn. 6 for equilibrium where $x = \Delta d$, we have

$$\left. \frac{\partial E}{\partial x} \right|_{x=\Delta d} = -\frac{\hbar^2}{4m d^3} + k\Delta d = 0 \quad \text{with} \quad d' = d + \Delta d \quad (7)$$

which renders

$$k\Delta d = \frac{h^2}{4md'^3} = \frac{2\varepsilon'_o}{d'} \quad \text{with} \quad \varepsilon'_o = \frac{h^2}{8md'^2} \quad (8)$$

Evidently, a change in E with a change in x from $x = 0$ to $x = \Delta d$ (*i.e.*, $\Delta E = E_{|x=\Delta d} - E_{|x=0}$) can be obtained from

$$\Delta E = \frac{h^2}{8md'^2} + \frac{1}{2}k\Delta d^2 - \frac{h^2}{8md'^2} \approx -\frac{h^2}{8md'^3}\Delta d = -\frac{\varepsilon'_o\Delta d}{d'} \quad (9)$$

where we use Eqns. 6 and 8. Since ΔE represents the net change in energy of the particle and the strained box as a single unit, its $-ve$ value signifies that the two assume a kind of single bound state and they remain in this state unless ΔE energy is supplied from outside. We also note that: (i) strain (Δd) as well as strain energy ($k\Delta d^2/2$) of the box are the functions of the ground state energy ($\varepsilon'_o = h^2/8md'^2$) or momentum ($h/2d'$) of the particle. In principle, one may observe similar binding ($\Delta E_n = -n^2\varepsilon_n\Delta d_n/d'$) for any Ψ_n state, however, a real bound state can be observed only for the particle in Ψ_1 state, because the particle in a $\Psi_{n>1}$ state is free to jump to any lower state by releasing out the difference in their energies which is always $+ve$, while the particle in the ground state (Ψ_1) does not have this option. It is for reason that we wrote Eqn. 6 for the particle in its ground state only.

2.3 Energy exchange between particle and strained box

We note that the total energy of the particle in Ψ_1 state in its equilibrium with strained box is

$$E_1 = \frac{h^2}{8md'^2} + \frac{1}{2}k\Delta d^2 \quad (10)$$

However, the equilibrium is disturbed if d' is changed to $d' \pm \eta$ (with $|\eta| < \Delta d$) and the energy of the disturbed system would be

$$E'_1 = \frac{h^2}{8m(d' \pm \eta)^2} + \frac{1}{2}k(\Delta d \pm \eta)^2 \quad (11)$$

which can be rearranged as

$$E'_1 \approx \varepsilon'_o + \epsilon_s \mp \left(\frac{2\varepsilon'_o}{d'} - k\Delta d\right)\eta + \frac{1}{2}k'\eta^2 \quad (12)$$

with $\epsilon_s = \frac{1}{2}k\Delta d^2$ being the strain energy of the box and modified force constant

$$k' = k + \frac{6\varepsilon_o}{d'^2} \quad (13)$$

Since the first two terms on the right hand side of Eqn. 12 are constant and $(2\varepsilon'_o/d' - k\Delta d)\eta$ vanishes for the equilibrium condition (Eqn. 8), it can be argued that, with the particle occupying its ground state energy, the box size oscillations are controlled by an increased force constant k' (Eqn. 13). In addition $(2\varepsilon'_o/d' - k\Delta d)\eta = 0$ implies that zero point energy of the particle increases (or *decreases*) by $(2\varepsilon'_o\eta/d')$ when strain energy decreases (

or *increases*) by equal amount, $(k\Delta d)\eta$) indicating that the particle and the strained box keeps exchanging energy with each other during η oscillations.

2.4 Self superposition, effective size and nature of motion:

Since Ψ_n is the superposition of two plain waves (*viz.*, $\exp(iqx)$ and its reflection $\exp(-iqx)$ representing one and the same particle), the particle in the box assumes a kind of self superposition state. Identifying these waves to represent two separate particles, one may find that Ψ_n can identically define the state of mutual superposition of two particles moving with q_n and $-q_n$ momenta. In fact as shown in [7, 14, 15], there is no means to ascertain whether the wave superposition of two particles represented by a function like Ψ_n defines their mutual superposition or the self superposition assumed separately by each of them.

We note that the spread (d) of Ψ_n is an integer multiple of $\lambda_n/2$ and $\lambda_n/2$ equals d only for the ground state $\Psi_{n=1}$. One may call $\lambda_n/2$ as the unit spread of Ψ_n or identify it as the quantum size (or the uncertainty size) of a particle of momentum q_n . It is well known that no particle can be accommodated in a box if its $\lambda/2 > d$ which means that no object can share the $\lambda/2$ space occupied by a particle, particularly, if the particle and chosen object interact through HC repulsion. Evidently, $\lambda/2$ represents the effective size of a quantum particle in a sense that any effort to reduce this size needs energy which implies that the particle *repels* the object(s) trying to share the $\lambda/2$ space occupied it. Naturally, $\lambda/2$ could also be identified as the range of such repulsion.

In the light of what follows from Section 2.1 and the fact that quantum size of a particle in its ground state fits exactly with the size of the box, it can be concluded that the particle has collisionless motion in its ground state ($\Psi_{n=1}$) and collisional motion in its excited states, $\Psi_{n>1}$. This agrees with similar inferences of [7, 14, 15] which imply that collisionless motion, effective size ($=\lambda/2$) and self superposition state are the common features of a particle in the low energy states of widely different many body systems where each atom could be identified with a particle trapped in a box.

3. Relevance and Experimental Support

We note that the present study can help in understanding some interesting aspects of quantum particles subjected to certain physical situations. For example an electron free to move within the space of an atomic vacancy in a 1-D crystal. We presume that: (i) the trapped electron is unable to cross the atoms on its two sides and (ii) the potential responsible for the forces restoring the positions of these atoms varies as x^2 (with x = displacement of an atom from its position decided by the inter-atomic interactions in the said crystal). While the former presumption implies that the electron is trapped between two infinite potential walls, the latter means that the forces restoring the size of the trapping vacancy are identical to F_a . Since this represents a case identical to the system of present study, we use its inferences to find that the said electron uses its F_o to displace the neighboring atoms to produce local strain in the crystal which leads to a binding of the electron with crystalline lattice (with binding energy ΔE , Eqn. 9) and facilitates an

exchange of energy with lattice oscillations (*i.e.*, phonons), particularly, when T of the crystal is low enough to keep the electron in its ground state.

Similarly, for two electrons occupying their ground state in two separate atomic vacancies, one may find that their F_o should produce local strain in the crystal leading to their binding with the lattice under the thermal conditions which keep them in their ground state. Both these electrons can, obviously, be visualized to have an *indirect mutual binding* (detailed mathematical analysis given in [16]) which helps them to have a correlated zero-point motion with an energy exchange through phonons. Naturally, if there are many such electrons in the crystal, this binding can be envisaged for each of their pair.

As shown in [6], we note that forces identical to f_n (Eqn. 4) operate also in case of a particle trapped in a 3-D box (a cavity in a crystal). Evidently, the inferences of this study (Section 2.0) are relevant to the physics of real systems like quantum dot, quantum wire, trapped single particle/ion [8,9], *etc.* and in this context, the experimentally observed *-ve* value of volume expansion coefficient, $(1/V)\partial_T V$ (an analogue of *-ve* $(1/d)\partial_T d$), for liquids ${}^4\text{He}$ and ${}^3\text{He}$ [17], respectively, around $\approx 2.2\text{K}$ and $\approx 0.6\text{K}$ (not significantly different from $T_o \approx 1.4\text{K}$ estimated for these liquids), provides *strong experimental support*. This experimental fact also implies that the ground state of each atom in systems like these liquids should be identical to the ground state of a particle trapped in a 3-D box (*i.e.* a cavity of neighboring particles) and F_o of each particle should, undoubtedly, have an important role in deciding the low T behavior of such systems. Interestingly, our recent papers, related to microscopic theory of superconductivity [14] and unification of the quantum behavior of widely different systems of interacting fermions and bosons [15], clearly prove that: (i) theories giving due importance to F_o are mathematically simple, (ii) they explain several unexplained experimental facts such as, high T_c of ceramic superconductors, superfluid T_c of liquid ${}^3\text{He}$, logarithmic singularity of specific heat of liquid ${}^4\text{He}$ at T_λ , *etc.*, and (iii) their results agree closely with experiments. Evidently, our inferences have enough experimental support and proven relevance to the microscopic understanding of widely different many body systems.

4. Concluding Remarks

We find that : (i) a particle confined to a box assumes a state (Ψ_n) which resembles with that of two particles in a state of equal and opposite momenta (*i.e.*, q and $-q$), (ii) a zero-point force (F_o) operates when the particle occupies ground state and produces non-zero strain (*viz.*, an expansion in d by Δd) since the forces restoring d , in all real situations, are not infinitely strong, (iii) the particle in its ground state has a kind of binding with strained box, (iv) the experimental observation of *-ve* $(1/d)\partial_T d$ around $T = T_o$ should be an evidence for the occurrence of this strain and the fall of particle into its ground state, and (v) the particle motion changes from collisional to collisionless when it moves from its excited state ($\Psi_{n>1}$) to ground state (Ψ_1). Since these results are as natural consequences of wave particle duality as Eqns. 1-4, the experimentally observed *-ve* $(1/V)\partial_T V$ for liquids ${}^4\text{He}$ and ${}^3\text{He}$ (respectively, around 2.2 and 0.6 K) unquestionably support the above mentioned inferences and this implies that: (i) mechanical strain resulting from F_o , (ii) *-ve* expansion coefficient around T_o and (iii) loss of relative motion leading to collisionless motion of quantum particles (*as concluded in [7] with more detailed*

discussion) should be common aspects of the low energy states (*close to ground state*) of widely different many body systems. It may, therefore, be emphasized that a microscopic theory of a many body system, like electron fluid, liquids ^4He and ^3He , *etc.*, can not truly explain their low T behavior unless it incorporates the role of F_o (*a basic consequence of the wave particle duality*) and related aspects. As a proof of this point one may find recent studies related to the basic foundations of microscopic theory of superconductivity [14] and unification of the physics of widely different systems of interacting bosons and fermions such as liquids ^4He and ^3He [15].

This study should be useful for the learners of wave- mechanics because it reveals some of the basic aspects of a particle in a box (*e.g.*, the consequences of F_o) for the first time and to this effect it deserves to be a part of the elementary texts on wave mechanics. It should also be interesting to those who desire to contribute towards a true and complete understanding of the low energy states of widely different many body systems and the exotic phenomena of superfluidity and superconductivity.

Acknowledgement : The author is thankful to Dr. R. Moro (School of Physics, Georgia Institute of Technology, Atlanta, USA) for his very useful suggestions.

References

- [1] D. Giulini, E. Joos, C. Kiefer, J. Kupsch, I. -O. Stamatescu and H.D. Zeh, *Decoherence and the appearance of classical world in quantum theory* Springer, Berlin (1996).
- [2] C. Monroe, D.M. Meekhof, B.E. King and D.J. Wineland, *A “Schrödinger cat” superposition state of an atom*, Science **272**, 1131-36(1996).
- [3] C. Kiefer, *On the interpretation of quantum theory - from Copenhagen to present day*, e-print : www.arxiv.org/quant-ph/0210152 (22 Oct, 2002), pp 1-11.
- [4] C.S. Unnikrishnan, *Is the quantum mechanical description of physical reality complete ? proposed resolution of EPR puzzle*, Foundations of Physics Letters **15** 1-25 (2002).
- [5] L.I. Schiff, *Quantum Mechanics* (3rd edition), McGraw Hill, Tokyo (1968), pp 38-39.
- [6] Yung-Kuo Lim, *Problems and solutions on quantum mechanics*, Sarat Book House Kolkata (2001) pp 20-21 (Originally published by World Scientific (1998)).
- [7] Y. S. Jain, *Wave mechanics of two hard core quantum particles in 1-D box*, Central European J. Phys. **2**, 709-719 (2004); arxiv.org/quant-ph/0603233.
- [8] L. Zhuang, L. Guo and S.Y. Chou, *Silicon single-electron quantum-dot transistor switch operating at room temperature* Appl. Phys. Lett. **72**, 1205-1209 (1998).
- [9] D.M. Meekhof, C. Monroe, B.E. King, W.M. Itano, and D.J. Wineland, *Generation of nonclassical motional states of a trapped atom* Phys. Rev. Lett. **76** 1796-99 (1996).

- [10] R. Moro, X. Xu, S. Yin and W.A. De Heer, Science **300**, 1265 (2003); R. Moro, S. Yin, X. Xu and W.A. De Heer, Phys. Rev. Lett. **93**, 08680 (2004)
- [11] W.A. De Heer, Rev. Mod. Phys. **65**, 611 (1993).
- [12] D.R. Tilley and J. Tilley, *Superfluidity and superconductivity*, Adam Hilger Ltd. Bristol (1986).
- [13] F. Dalfovo, S. Giorgini, L.P. Pitaevskii and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999), and arxiv.org/Cond-mat/9806038
- [14] Y.S. Jain, *Basic foundations of the microscopic theory of superconductivity*, arxiv.org/cond-mat/0603784.
- [15] Y.S. Jain, *Unification of the physics of bosonic and fermionic systems*, Ind. J. Phys. **79**, 1009 (2005) (A brief report of the invited talk given at CMDAYS-04, August 25-27, 2004, Department of Physics, NEHU, Shillong-793 022, India.);
Note: While the factor $|\sin(\mathbf{k}\cdot\mathbf{r}/2)|$ in Eqn. 5 for Ψ^+ in this reference should be read as $\sin(|\mathbf{k}\cdot\mathbf{r}|/2)$, $E_g(T)/Nk_B T$ in Eqn. 25 be read as $E_g(T)/Nk_B$. Our detailed research papers on the microscopic theories of liquids ^4He and ^3He which not only form the basis of the unification discussed in this paper but also establish F_o and related aspects as their basic foundations would soon be available on this arxiv.
- [16] Y.S. Jain, *Wave mechanics of three different systems of a particle in 1-D box and a 1-D oscillator* (Technical Report SSP/YSJ-02 (2005)).
- [17] J. Wilks, *The properties of liquid and solid helium*, Clarendon Press, Oxford (1967).