

Value Assignments to Observables Depending on a History of Context Change

Satoshi Uchiyama^y

Department of Life and Creative Sciences
Hokusei Gakuin University Junior College
Atsubetsu-ku, Sapporo 004-8631, Japan.

April 17, 2024

Abstract

The functional composition principle is generalized by taking into account of history of context change. Analysis of Peres' example shows hysteresis of value assignments. It is shown that value assignments which depend on the history of context change are possible in the case that the Hilbert space of state vectors is finite dimensional.

1 A generalized functional composition principle depending on history of context change

The Kochen-Specker theorem states that there is no value assignment for quantum mechanical observables that satisfies the functional composition principle (FUNC) when the dimension of the Hilbert space of state vectors is greater than two [1, 2]. By analogy with Riemannian surfaces which settle contradictory concept of multivalued functions by analytic continuation, we investigate a value assignment depending on a history of context change.

The eigenvalues of an observable represented by a self-adjoint operator are values which are obtainable when one makes an observation. If there was hidden variables that specify the observed value, we can assign the value to the pair of the observable and the hidden variables. If we denote the hidden variables, the observable, and the eigenvalue by ω , \hat{O} , and ω_i , respectively, then a value assigning map v is defined as a mapping of the set of all observables into a set of random variables such that for the possible ω ,

$$v(\hat{O})(\omega) = \omega_i : \quad (1)$$

email: uchiyama@hokuseiac.jp

^yAddress until the end of Feb. 2007: Centro Vito Volterra, Università degli Studi di Roma "Tor Vergata", Via Columbia, Rome 00133, Italy.

The functional composition principle (FUNC) states that for observables \hat{A} and \hat{B} , if there exists a function f such that $\hat{A} = f(\hat{B})$, then a value assigning map v satisfies that

$$v(\hat{A})(!) = f(v(\hat{B})(!)) : \quad (2)$$

Physically, a context of a measurement of an observable \hat{O} is determined by a setting of the measurement apparatus. Mathematically, a maximal Boolean sublattice of a lattice of observational propositions on a quantum system determines a context of a measurement. An observational proposition of an yes-no experiment is represented by a projection operator. There is a one-to-one mapping of the set of all unit vectors into the set of projection operators of rank 1. By using this mapping, we can specify a maximal Boolean sublattice, i.e., a context, by a complete orthonormal system (CONS) of vectors. We define an equivalence relation on the set of all CONSs. We shall say that two CONSs $(j_1 i; \dots; j_n i)$ and $(j_1 i; \dots; j_n i)$ are equivalent if there exist a permutation p of the index set $1; 2; \dots; n$ and phase factors $\exp(i\theta_i)$ ($i = 1; 2; \dots; n$) such that $j_{ji} = \exp(i\theta_{p(j)}) j_{p(j)i}$ ($j = 1; 2; \dots; n$). Thus a context is an equivalent class with respect to this equivalence relation and some CONS is a representative of a context. We denote a set of contexts by C . C is a subset of the quotient space with respect to the equivalence relation of the complex Stiefel manifold of orthonormal n -frames.

If a measurement apparatus that is not set in a context of a measurement of an observable \hat{O} gave outcomes, then results would be invalid, i.e., the obtained values could be different from the eigenvalues. On the contrary, a measurement apparatus that is set in the context of a measurement of the observable \hat{O} gives eigenvalues of \hat{O} as outcomes. In this sense, the measurement apparatus gives stable results for \hat{O} in the context. We say that an observable \hat{O} is stable in a context $_ \in C$ and denoting it by $\hat{O} / _$, if \hat{O} is diagonalized in a CONS $(j_1 i; \dots; j_n i)$ that is a representative of $_$.

When an observable \hat{O} is not stable in a context $_$, there is no experimental restriction on values of \hat{O} . Hence it is natural to consider that a value assigning map depends on contexts, and therefore the domain of a value assigning map is the set of observables that are stable in a given context. Thus the contextual version of FUNC becomes as follows.

Contextual version of FUNC (cFUNC) If $\hat{A} = f(\hat{B})$ and $\hat{B} / _$ then,

$$v_(\hat{A}) = f(v_(\hat{B})); \text{ a.s. on } _ : \quad (3)$$

Here, $_$ is the set of hidden variables and a probability measure on $_$ characterizing the ensemble of the hidden variables is supposed to be.

It is more general to consider that a value assigning map may depend on the contexts that the system experienced in the past. A history of context change is a finite sequence of contexts $(_ ; \dots; _)$, where $_ ; \dots; _ \in C$. We use a more intuitive notation about this such as $_ ! _ ! \dots ! _$.

Generalized version of FUNC (gFUNC) If $\hat{A} = f(\hat{B})$ and $\hat{B} / _$ then,

$$v_ ! _ ! _ (\hat{A}) = f(v_ ! _ ! _ (\hat{B})); \text{ a.s. on } _ : \quad (4)$$

for $8_{-}; \quad 2_{-} C$, where $!$ represents the same history of context change in both sides of the equation.

Proposition 1 Provided $gFUNC$, if observables \hat{A} and \hat{B} are stable in a context $\underline{\quad}$, then

$$v_{\underline{\quad}}(\hat{A} + \hat{B}) = v_{\underline{\quad}}(\hat{A}) + v_{\underline{\quad}}(\hat{B}); \quad (5)$$

$$v_{\underline{\quad}}(\hat{A} \hat{B}) = v_{\underline{\quad}}(\hat{A}) \cdot v_{\underline{\quad}}(\hat{B}); \text{ a.s. on } \underline{\quad}; \quad (6)$$

Proof. Since $\hat{A}, \hat{B} / \underline{\quad}$, they can be diagonalised simultaneously. Then there exist an observable \hat{C} and functions g and h such that $\hat{A} = g(\hat{C})$ and $\hat{B} = h(\hat{C})$.

$$\begin{aligned} & v_{\underline{\quad}}(\hat{A} + \hat{B}) \\ &= v_{\underline{\quad}}(g(\hat{C}) + h(\hat{C})) \\ &= v_{\underline{\quad}}((g + h)(\hat{C})) \\ &\stackrel{gFUNC}{=} (g + h)(v_{\underline{\quad}}(\hat{C})) \\ &= g(v_{\underline{\quad}}(\hat{C})) + h(v_{\underline{\quad}}(\hat{C})) \\ &\stackrel{gFUNC}{=} v_{\underline{\quad}}(g(\hat{C})) + v_{\underline{\quad}}(h(\hat{C})) \\ &= v_{\underline{\quad}}(\hat{A}) + v_{\underline{\quad}}(\hat{B}); \end{aligned}$$

$$\begin{aligned} & v_{\underline{\quad}}(\hat{A} \hat{B}) \\ &= v_{\underline{\quad}}(g(\hat{C}) \cdot h(\hat{C})) \\ &= v_{\underline{\quad}}((g \cdot h)(\hat{C})) \\ &\stackrel{gFUNC}{=} (g \cdot h)(v_{\underline{\quad}}(\hat{C})) \\ &= g(v_{\underline{\quad}}(\hat{C})) \cdot h(v_{\underline{\quad}}(\hat{C})) \\ &\stackrel{gFUNC}{=} v_{\underline{\quad}}(g(\hat{C})) \cdot v_{\underline{\quad}}(h(\hat{C})) \\ &= v_{\underline{\quad}}(\hat{A}) \cdot v_{\underline{\quad}}(\hat{B}); \end{aligned}$$

□

Remark. Difficulty of defining a value assigning map with the $FUNC$ appears when the \hat{A} is degenerated. Consider the following matrices:

$$\begin{aligned} \hat{B} &= \begin{array}{c|c|c} 0 & 1 & 0 \\ \hline 0 & 0 & 2 \end{array} \begin{array}{c} 0 \\ 1 \\ 0 \end{array} \hat{A}; \\ \hat{C} &= \begin{array}{c|c|c} 0 & \frac{1}{2} & \frac{1}{2} \\ \hline \frac{1}{2} & \frac{1}{2} & 0 \end{array} \begin{array}{c} 0 \\ 0 \\ 1 \end{array} \hat{A} \oplus \begin{array}{c|c|c} 1 & 0 & 0 \\ \hline 0 & 2 & 0 \end{array} \begin{array}{c} 1 \\ 0 \\ 0 \end{array} \hat{A} \oplus \begin{array}{c|c|c} 0 & \frac{1}{2} & \frac{1}{2} \\ \hline \frac{1}{2} & \frac{1}{2} & 0 \end{array} \begin{array}{c} 0 \\ 0 \\ 1 \end{array} \hat{A} = \begin{array}{c|c|c} 0 & \frac{3}{2} & \frac{1}{2} \\ \hline \frac{1}{2} & \frac{3}{2} & 0 \end{array} \begin{array}{c} 0 \\ 0 \\ 3 \end{array} \hat{A}; \end{aligned}$$

Suppose functions f and g are give by

$$f(x) = \begin{cases} 2 & \text{if } x < 3 \\ x & \text{otherwise} \end{cases} ; \quad g(x) = \begin{cases} 2 & \text{if } x = 2 \\ x & \text{otherwise} \end{cases} ; \quad (7)$$

respectively. Then

$$f(\hat{B}) = \begin{array}{c|cc|c} 0 & 2 & 0 & 0 \\ \hline 0 & 0 & 2 & 0 \\ \hline 0 & 0 & 0 & 3 \end{array} \begin{array}{c} 1 \\ A \end{array} ; \quad (8)$$

$$g(\hat{C}) = \begin{array}{c|cc|c} 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ \hline \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 \\ \hline 0 & 0 & 0 & 1 \end{array} \begin{array}{c} 1 \\ C \\ A \end{array} \begin{array}{c|cc|c} 0 & 2 & 0 & 0 \\ \hline 0 & 0 & 2 & 0 \\ \hline 0 & 0 & 0 & 3 \end{array} \begin{array}{c} 1 \\ A \\ B \end{array} \begin{array}{c|cc|c} 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ \hline \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 \\ \hline 0 & 0 & 0 & 1 \end{array} \begin{array}{c} 1 \\ C \\ A \end{array}$$

$$= \begin{array}{c|cc|c} 0 & 2 & 0 & 0 \\ \hline 0 & 0 & 2 & 0 \\ \hline 0 & 0 & 0 & 3 \end{array} \begin{array}{c} 1 \\ A \end{array} = f(\hat{B}) = \hat{A} :$$

Although \hat{B} and \hat{C} are stable in different contexts respectively, \hat{A} is stable in both of the contexts. Thus one has to assign a value to \hat{A} so that the assigned value would be consistent with values assigned to \hat{B} and \hat{C} .

Since cFUNC and gFUNC give no constraint among value assignments in different contexts, the space of hidden variables is broken up into independent probability spaces of different contexts. To synthesize them, we require the following condition.

Non-transition condition (n-T R N S). For all contexts $\omega, \omega' \in C$, if \hat{B} / ω and \hat{B} / ω' , then

$$v_{\omega'}(\hat{B})(!) = v_{\omega}(\hat{B})(!); \text{ a.s. on } \omega ; \quad (9)$$

where $!$ represents the same history of context change in both sides of the equation.

Remark. In the EPR {Bohm Gedankenexperiment, since the spin observables of particles in a pair commute, n-T R N S means that an assigned value of a component of the spin of one particle does not change, when we change the setting of the measuring apparatus for the spin of the other particle. But locality in this sense is weaker than the genuine locality, since it allows hysteresis of value assignments.

In the rest of this section, we clarify what kinds of observables are stable in two different contexts.

Proposition 2 There exists a nonzero self-adjoint operator that is diagonalized in two different CONSS $(j_1 i; :::; j_n i)$ and $(j_1 i; :::; j_n i)$ if and only if there

exist partitions $fI_1; \dots; I_m g$ and $fJ_1; \dots; J_m g$ of the index set $X = f1; \dots; ng$ ($m \leq n$) such that

$$\text{span } f j_{i1} : i2 I_k g = \text{span } f j_{j1} : j2 J_k g; \quad (10)$$

for $k = 1; \dots; m$.

Proof. (i) Let \hat{O} be a self-adjoint operator that is diagonalized in both CONSSs $(j_{i1}; \dots; j_{in})$ and $(j_{j1}; \dots; j_{jn})$. Suppose \hat{O} has eigenvalues $o_1; \dots; o_m$ that are distinct from each other ($m \leq n$). We define a partition $fI_k g$ of the index set X by

$$I_k = \{ i2 X : \hat{O} j_{i1} = o_k j_{i1} \}; \quad (11)$$

Similarly, we define a partition $fJ_k g$ of X by

$$J_k = \{ j2 X : \hat{O} j_{j1} = o_k j_{j1} \}; \quad (12)$$

Since $\text{span } f j_{i1} : i2 I_k g$ and $\text{span } f j_{i1} : i2 J_k g$ is the eigenspace of \hat{O} with the eigenvalue o_k , they coincide with each other.

(ii) Let o_k s are increasing real numbers, i.e., $o_1 < o_2 < \dots < o_m$. We define a self-adjoint operator \hat{O} by

$$\hat{O} = \sum_{k=1}^m o_k \sum_{i2 I_k} j_{i1} h_{i1} j; \quad (13)$$

this is diagonalized in $(j_{i1}; \dots; j_{in})$. Since $\sum_{i2 I_k} j_{i1} h_{i1} j = \sum_{j2 J_k} j_{j1} h_{j1} j$, \hat{O} is diagonalized in $(j_{j1}; \dots; j_{jn})$. \square

Remark. When $m = 1$, the self-adjoint operator is I , i.e., the identity operator. $m = n$ means that the two CONSSs are representatives of the same context.

Lemma 1 For two different CONSSs $(j_{i1}; \dots; j_{in})$ and $(j_{j1}; \dots; j_{jn})$, there exists uniquely the finest partitions $fI_1; \dots; I_m g$ and $fJ_1; \dots; J_m g$ of the index set $X = f1; \dots; ng$ of the CONSSs respectively ($m \leq n$) such that

$$\text{span } f j_{i1} : i2 I_k g = \text{span } f j_{j1} : j2 J_k g; \quad (14)$$

for $k = 1; \dots; m$.

Proof. Consider partitions $fI_1^0; \dots; I_m^0 g$ and $fJ_1^0; \dots; J_m^0 g$ of the index set X ($m^0 \leq n$) such that

$$\text{span } f j_{i1} : i2 I_k^0 g = \text{span } f j_{j1} : j2 J_k^0 g;$$

for $k = 1; \dots; m^0$. If there exist another partitions $fI_1^{00}; \dots; I_m^{00} g$ and $fJ_1^{00}; \dots; J_m^{00} g$ of the index set X ($m^{00} \leq n$) such that

$$\text{span } f j_{i1} : i2 I_k^{00} g = \text{span } f j_{j1} : j2 J_k^{00} g;$$

for $k = 1; \dots; m^0$, then put

$$\begin{aligned}\tilde{I}_{(k;l)} &:= I_k^0 \setminus I_1^0; \\ \tilde{J}_{(k;l)} &:= J_k^0 \setminus J_1^0;\end{aligned}$$

for $k = 1; \dots; m^0, l = 1; \dots; m^0$. Since

$$\bigcap_{i: i \in I_{(k;l)}} \tilde{I}_{(k;l)} = \text{span} \{f_{j: i \in I_k^0} g\} \setminus \text{span} \{f_{j: i \in I_1^0} g\}$$

and

$$\bigcap_{j: j \in J_{(k;l)}} \tilde{J}_{(k;l)} = \text{span} \{f_{j: j \in J_k^0} g\} \setminus \text{span} \{f_{j: j \in J_1^0} g\};$$

we obtain

$$\bigcap_{i: i \in I_{(k;l)}} \tilde{I}_{(k;l)} = \text{span} \{f_{j: j \in J_{(k;l)}} g\} : \quad (15)$$

Thus $f_{\tilde{I}_{(k;l)}} g$ and $f_{\tilde{J}_{(k;l)}} g$ are finer partitions of X satisfying the condition (15) than $f_{I_k^0} g$ and $f_{J_1^0} g$ or coincide with $f_{I_k^0} g$ and $f_{J_1^0} g$. Since the index set X is finite, above procedure ends with finite times and we obtain the finest partitions $f_{I_i} g$ and $f_{J_i} g$ satisfying the condition (14). The uniqueness follows from the maximality of the $f_{I_i} g$ and the $f_{J_i} g$. \square

Corollary 1 For two different contexts $\underline{\omega} = (j_1 i; \dots; j_n i)$ and $\underline{\omega}' = (j_1 i; \dots; j_n i)$, an observable \hat{O} is stable in $\underline{\omega}$ and $\underline{\omega}'$ if and only if \hat{O} can be written as

$$\hat{O} = \sum_{k=1}^m o_k P_k; \quad (16)$$

where o_k is an eigenvalue of \hat{O} and P_k is given as $P_k := \prod_{i \in I_k} \prod_{j \in J_i} f_{j: i \in I_i} g$ by using the finest partitions $f_{I_1}; \dots; f_{I_m} g$ and the $f_{J_1}; \dots; f_{J_m} g$ of the index set given by Lemma 1.

Proof . It follows from the spectral decomposition of \hat{O} , Proposition 2 and Lemma 1. \square

2 Contextuality in the Peres' example of nonexistence of noncontextual hidden variables

Peres showed that it is impossible to assign a value to $(x \quad y) \quad (y \quad x)$ noncontextually in the singlet state [3]. We denote the eigenvectors with eigenvalue

1 of x by $X = x; y; z$. Here are the contexts that appear in the Peres' example:

$$\begin{aligned}
- & (x + i \quad x + i; x \quad i \quad x \quad i; x + i \quad x \quad i; x \quad i \quad x + i); (17) \\
- & (y + i \quad y + i; y \quad i \quad y \quad i; y + i \quad y \quad i; y \quad i \quad y + i); (18) \\
- & (z + i \quad z + i; z \quad i \quad z \quad i; z + i \quad z \quad i; z \quad i \quad z + i); (19) \\
- & (x + i \quad y + i; x \quad i \quad y \quad i; x + i \quad y \quad i; x \quad i \quad y + i); \\
- & (y + i \quad x + i; y \quad i \quad x \quad i; y + i \quad x \quad i; y \quad i \quad x + i); \\
- & \frac{1}{2} (x + i \quad z + i + \frac{p}{1} z \quad i \quad z \quad i); \\
- & \frac{1}{2} (y + i \quad z + i + \frac{p}{1} z \quad i \quad z \quad i); \\
- & \frac{1}{2} (x + i \quad z \quad i + \frac{p}{1} z \quad i \quad z + i); \\
- & \frac{1}{2} (y + i \quad z \quad i + \frac{p}{1} z \quad i \quad z + i) :
\end{aligned}$$

We can see the following:

$$\begin{aligned}
- & \cdot \quad x \quad I; I \quad x; \quad x \quad x; \\
- & \cdot \quad y \quad I; I \quad y; \quad y \quad y; \\
- & \cdot \quad z \quad I; I \quad z; \quad z \quad z; \\
- & \cdot \quad x \quad I; I \quad y; \quad x \quad y; \\
- & \cdot \quad y \quad I; I \quad x; \quad y \quad x; \\
- & \cdot \quad x \quad y; \quad y \quad x; \quad z \quad z;
\end{aligned}$$

We analyze the Peres' example by exploiting gFUNC and n-TRNS. In order to do this, we need one more condition. Perfect anti-correlation (a-CRL). For the contexts $_$ of (17), $_$ of (18), $_$ of (19),

$$\begin{aligned}
v_{!}(_ (x \quad I) (!)) &= v_{!}(_ (I \quad x) (!)); \\
v_{!}(_ (y \quad I) (!)) &= v_{!}(_ (I \quad y) (!)); \\
v_{!}(_ (z \quad I) (!)) &= v_{!}(_ (I \quad z) (!));
\end{aligned}$$

respectively, for all possible $! \in \{0, 1\}$. Here $!$ represents the same history of context change in both sides of the equations.

Remark. If we add a rule that we can replace an observable by its eigenvalue in the corresponding eigenstate, then a-CRL follows from gFUNC.

Proposition 3 If gFUNC, n-TRNS, a-CRL hold and eigenvalues are assigned to the corresponding observables, then

$$v_{!}((x \quad y) \quad (y \quad x)) = 1; \text{ a.s.} \quad : \quad (20)$$

Proof .

$$\begin{aligned}
& \underline{V}((x \quad y) \quad (y \quad x)) \\
&= \underline{V}(z \quad z) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ (z \quad z) \\
&= \underline{V}_! _ ((z \quad I) \quad (I \quad z)) \\
&\stackrel{gF \text{ UNC}}{=} \underline{V}_! _ (z \quad I) \underline{V}_! _ (I \quad z) \\
&\stackrel{a \text{ CRL}}{=} (\underline{V}_! _ (z \quad I))^2 = 1: \tag{21}
\end{aligned}$$

The last equality follows from that the eigenvalues of $z \quad I$ are ± 1 . \square

Proposition 4 If gFUNC, n-TRNS, a-CRL hold, then

$$\begin{aligned}
& \underline{V}((x \quad y) \quad (y \quad x)) \\
&= \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ \underline{V}_! _ (x \quad I) \tag{22}
\end{aligned}$$

a.s. on \mathcal{S} .

Proof .

$$\begin{aligned}
& \underline{V}((x \quad y) \quad (y \quad x)) \\
&\stackrel{gF \text{ UNC}}{=} \underline{V}(x \quad y) \underline{V}(y \quad x) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ (x \quad y) \underline{V}_! _ (y \quad x) \\
&= \underline{V}_! _ ((x \quad I) (I \quad y)) \underline{V}_! _ (y \quad x) \\
&\stackrel{gF \text{ UNC}}{=} \underline{V}_! _ (x \quad I) \underline{V}_! _ (I \quad y) \underline{V}_! _ (y \quad x) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ (I \quad y) \underline{V}_! _ (y \quad x) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (I \quad y) \underline{V}_! _ (y \quad x) \\
&\stackrel{a \text{ CRL}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ (y \quad x):
\end{aligned}$$

We can proceed along calculating.

$$\begin{aligned}
& \underline{V}((x \quad y) \quad (y \quad x)) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ (y \quad x) \\
&\stackrel{gF \text{ UNC}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ (y \quad I) \underline{V}_! _ (I \quad x) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ (y \quad I) \underline{V}_! _ \underline{V}_! _ (I \quad x) \\
&\stackrel{a \text{ CRL}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ (y \quad I) \underline{V}_! _ \underline{V}_! _ (x \quad I) \\
&\stackrel{n \text{ TRNS}}{=} \underline{V}_! _ \underline{V}_! _ (x \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ \underline{V}_! _ (y \quad I) \underline{V}_! _ \underline{V}_! _ (x \quad I):
\end{aligned}$$

□

Proposition 5 If $gFUNC$, $n-TRNS$, $a-CRL$ hold, then a value assignment has hysteresis.

Proof. By Proposition 3, the r.h.s. of (22) is equal to 1. Because every term s of the r.h.s. of (22) assume only the values ± 1 , exactly one of $v_{x-}(x, I)$, $v_{x-}(y, I)$, $v_{y-}(y, I)$, $v_{y-}(x, I)$ must have opposite sign from the ones of the rests. We note that $v_{x-}(x, I) = v_{x-}(x, I)$ and $v_{x-}(y, I) = v_{y-}(y, I)$ do not follow from $n-TRNS$, because x, I is not stable in the context $_$ and y, I is not stable in the context $_$. Therefore we obtain

$$v_{x-}(x, I) = v_{x-}(x, I); \quad v_{x-}(y, I) \neq v_{y-}(y, I); \quad (23)$$

or

$$v_{x-}(x, I) \neq v_{x-}(x, I); \quad v_{x-}(y, I) = v_{y-}(y, I); \quad (24)$$

This shows hysteresis of the value assignment to x, I or y, I . □

Remark. $v_{x-}(y, I)$ shows that the system experienced the context $_$ in which y, I was not stable. $v_{x-}(x, I)$ shows that the system experienced the context $_$ in which x, I was not stable. Only one of $v_{x-}(y, I)$ and $v_{y-}(x, I)$, however, can assume values different from the values through a different values through a different history of context change. This is similar to the case of the Kocken{Specker paradox on spin-1 observables [1].

Proposition 6 If $gFUNC$, $n-TRNS$, $a-CRL$ hold and eigenvalues are assigned to the corresponding observables, then there is no noncontextual value assigning map in the EPR {Bohm Gedankenexperiment.

Proof. Suppose that the value assigning map is noncontextual. Then (22) becomes

$$\begin{aligned} & v_{x-}(x, I) v_{x-}(y, I) v_{y-}(y, I) v_{y-}(x, I) \\ &= v(x, I) v(y, I) v(y, I) v(x, I) \\ &= v(x, I)^2 v(y, I)^2 = +1: \end{aligned}$$

Thus this contradicts (20); there is no such value assigning map. □

Remark. This is nothing else but what Peres shows in ref. [3].

3 Context changing maps

We associate a mapping $\underline{\cdot}^!$ of $\underline{\cdot}$ into $\underline{\cdot}$ with a context change from a context $\underline{\cdot} = (j_1 i; \dots; j_n i)$ to a context $\underline{\cdot} = (j_1 i; \dots; j_n i)$.

Let $fI_k g$ and $fJ_k g$ be the finest partitions of the index set $f1; \dots; ng$ of Lemma 1. Let q be a permutation of the index set such that $i \in I_k$ implies $q(i) \in J_k$ ($k = 1; \dots; m$). We define a unitary transformation $U_{\underline{\cdot}^! \underline{\cdot}}$ by

$$U_{\underline{\cdot}^! \underline{\cdot}} = \prod_{k=1}^m \prod_{i \in I_k} \prod_{j \in J_k} j_{q(i)} i h_{ij} \quad (25)$$

$U_{\underline{\cdot}^! \underline{\cdot}}$ changes the context $\underline{\cdot}$ to $\underline{\cdot}$. Symbolically, we write

$$\underline{\cdot} = U_{\underline{\cdot}^! \underline{\cdot}} \underline{\cdot} \quad (26)$$

Proposition 7 If $\hat{O} / \underline{\cdot}; \underline{\cdot}$, then $U_{\underline{\cdot}^! \underline{\cdot}} \hat{O} U_{\underline{\cdot}^! \underline{\cdot}}^y = \hat{O}$.

Proof. By Corollary 1 and the definition of $U_{\underline{\cdot}^! \underline{\cdot}}$, denoting eigenvalues of \hat{O} by o_k s, we have

$$\begin{aligned} U_{\underline{\cdot}^! \underline{\cdot}} \hat{O} U_{\underline{\cdot}^! \underline{\cdot}}^y &= \prod_{k=1}^m \prod_{i \in I_k} \prod_{j \in J_k} j_{q(i)} i h_{ij} o_1 \prod_{i \in I_1} j_{i^0} i h_{i^0 j} U_{\underline{\cdot}^! \underline{\cdot}}^y \\ &= \prod_{k=1}^m \prod_{i \in I_k} j_{q(i)} i o_k h_{ij} U_{\underline{\cdot}^! \underline{\cdot}}^y \\ &= \prod_{k=1}^m \prod_{i \in I_k} o_k j_{q(i)} i h_{ij} \prod_{i \in I_1} j_{i^0} i h_{i^0 j} \\ &= \prod_{k=1}^m \prod_{i \in I_k} o_k j_{q(i)} i h_{q(i) j} \\ &= \prod_{k=1}^m \prod_{j \in J_k} o_k j_j i h_{jj} \\ &= \hat{O} : \end{aligned}$$

□

We say that a value assigning map $v_{\underline{\cdot}^! \underline{\cdot}}$ admits a context changing map from $\underline{\cdot}$ to $\underline{\cdot}$, if for an arbitrary observable $\hat{O} / \underline{\cdot}; \underline{\cdot}$, there exists a one-to-one mapping $\underline{\cdot}^!$ of $\underline{\cdot}$ onto $\underline{\cdot}$ such that

$$v_{\underline{\cdot}^! \underline{\cdot}}(U_{\underline{\cdot}^! \underline{\cdot}} \hat{O} U_{\underline{\cdot}^! \underline{\cdot}}^y)(!) = v_{\underline{\cdot}^! \underline{\cdot}}(\hat{O})(\underline{\cdot}^!(!)); \quad \text{a.s. on } \underline{\cdot} : \quad (27)$$

We call such $\underline{\cdot}^!$ a context changing map from $\underline{\cdot}$ to $\underline{\cdot}$.

Proposition 8 If a value assigning map $v_{\underline{\cdot}^! \underline{\cdot}}$ admits a context changing map and an observable \hat{O} is stable in the contexts $\underline{\cdot}$ and $\underline{\cdot}$, then

$$v_{\underline{\cdot}^! \underline{\cdot}}(\hat{O})(\underline{\cdot}^!(!)) = v_{\underline{\cdot}^! \underline{\cdot}}(\hat{O})(!); \quad \text{a.s. on } \underline{\cdot} : \quad (28)$$

P roof . By P roposition 7, for all possible $! 2$,

$$\begin{aligned}
(27) \quad & \mathbf{v} \quad \text{!} \text{ } _ \text{ } (\hat{\mathcal{O}}) \text{ } (\text{!} \text{ } _ \text{ } \text{ } ^1 \text{ } (\text{!})) \\
& \mathbf{v} \quad \text{!} \text{ } _ \text{ } \text{ } \text{!} \text{ } _ \text{ } (\mathbf{U} \text{!} \text{ } _ \text{ } \text{ } \hat{\mathcal{O}} \text{ } \mathbf{U} \text{!} \text{ } _ \text{ } \text{ } ^y \text{ }) (\text{!}) \\
& = \mathbf{v} \quad \text{!} \text{ } _ \text{ } \text{ } \text{!} \text{ } _ \text{ } (\hat{\mathcal{O}}) (\text{!}) :
\end{aligned}$$

Proposition 9 Suppose that a value assigning map $v : _ \rightarrow _$ admits a context changing map $m : _ \rightarrow _$ from $_$ to $_$. Let f be a function such that $\hat{A} = f(\hat{B})$ for some observables $\hat{A}, \hat{B} / _$. If $\hat{A} / _$ and

$$v_{\perp}(\hat{A}) = f v_{\perp}(\hat{B}) \quad ; \text{ a.s. on } \quad (29)$$

holds, then

$$\hat{A} = f(\hat{U}_1, \hat{B}U_1^y); \quad (30)$$

$$v_{\alpha}(\hat{A}) = f v_{\alpha}(U_{\alpha}^{-1} \hat{B} U_{\alpha}) ; \text{ a.s. on } : \quad (31)$$

P r o o f . Since $\hat{A} / _ , _ ,$ we have by P r o p o s i t i o n 7

$$f(U_{-}, \hat{B}U_{-}^y) = U_{-}, \hat{A}U_{-}^y = \hat{A}:$$

By (27),

$$\begin{aligned}
& \mathbf{v} \quad \mathbf{!}_{-} \quad \mathbf{!}_{-} \quad (\hat{\mathbf{A}}) \quad (\mathbf{!}_{-}) \\
& = \mathbf{v} \quad \mathbf{!}_{-} \quad \mathbf{!}_{-} \quad (\mathbf{U}_{-} \mathbf{!}_{-} \quad \hat{\mathbf{A}} \quad \mathbf{U}_{-}^{\mathbf{y}} \quad \mathbf{!}_{-}) \quad (\mathbf{!}_{-}) \\
(27) \quad & \mathbf{v} \quad \mathbf{!}_{-} \quad (\hat{\mathbf{A}}) \quad (\mathbf{!}_{-} \quad \mathbf{!}_{-}^{-1} \quad (\mathbf{!}_{-})) \\
(29) \quad & \mathbf{f} \quad \mathbf{v} \quad \mathbf{!}_{-} \quad (\hat{\mathbf{B}}) \quad (\mathbf{!}_{-} \quad \mathbf{!}_{-}^{-1} \quad (\mathbf{!}_{-})) \\
(27) \quad & \mathbf{f} \quad \mathbf{v} \quad \mathbf{!}_{-} \quad \mathbf{!}_{-} \quad (\mathbf{U}_{-} \mathbf{!}_{-} \quad \hat{\mathbf{B}} \quad \mathbf{U}_{-}^{\mathbf{y}} \quad \mathbf{!}_{-}) \quad (\mathbf{!}_{-}) :
\end{aligned}$$

1

Proposition 10 For different contexts $_ (j_1 i; :::; j_n i), _ (j_1 i; :::; j_n i), _ (j_1 i; :::; j_n i),$ de ne

$$S_{\underline{a}}(\underline{a}! \underline{a}) \doteq U_{\underline{a}}^y \quad U_{\underline{a}}! \underline{a} \quad U_{\underline{a}}! \underline{a} : \quad (32)$$

Then $S_{(_! _)}^{_}$ is a permutation of the CONS $(j_1 i; ::; j_n i)$, i.e., there exists a permutation p of the index set $f1; ::; ng$ such that

$$j_{p(i)} = S_{-}(\quad)j_i; \quad i = 1, \dots, n: \quad (33)$$

Proof. By definition of U_{α}^{β} , for each i there is j such that $j_{\beta}i = U_{\alpha}^{\beta}j_{\alpha}i$. Similarly, there exist k and l such that $j_{\alpha}i = U_{\beta}^{\alpha}j_{\beta}i$ and $j_{\alpha}i = U_{\beta}^{\alpha}j_{\beta}i$. Hence $S_{\alpha}(\beta)j_{\beta}i = j_{\alpha}i$, and therefore $p(i) = l$. It is clear that this p is a permutation of the index set. \square

Proposition 11 For a history of context change $\alpha \rightarrow \beta \rightarrow \gamma \rightarrow \delta$, there exists a permutation p of the index set $\{1, \dots, n\}$ such that

$$U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}j_{\delta}i = U_{\alpha}^{\delta}j_{\alpha}i; i = 1, \dots, n; \quad (34)$$

Proof. In the history of context change $\alpha \rightarrow \beta \rightarrow \gamma \rightarrow \delta$, $j_{\delta}i$ is mapped to

$$U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}j_{\delta}i:$$

Using the notation $S_{\alpha}(\beta)$ defined as (32),

$$\begin{aligned} & U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}j_{\delta}i \\ &= U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}S_{\alpha}(\beta)j_{\alpha}i \\ &= U_{\alpha}^{\beta}U_{\beta}^{\gamma}S_{\alpha}(\beta)S_{\beta}(\gamma)j_{\beta}i \\ &= U_{\alpha}^{\beta}S_{\alpha}(\beta)S_{\beta}(\gamma)S_{\gamma}(\delta)j_{\delta}i \end{aligned}$$

Thus we obtain

$$U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}j_{\delta}i = S_{\alpha}(\beta)S_{\beta}(\gamma)S_{\gamma}(\delta)j_{\delta}i; \quad (35)$$

By successive applications of Proposition 10, there exist a permutation p of the index set such that

$$S_{\alpha}(\beta)S_{\beta}(\gamma)S_{\gamma}(\delta)j_{\delta}i = j_{\alpha}p(i)$$

and it is clear that this p satisfies (34). \square

Corollary 2 For a history of context change $\alpha \rightarrow \beta \rightarrow \gamma \rightarrow \delta$, there exists a permutation p of the index set $\{1, \dots, n\}$ such that

$$U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}j_{\delta}i = j_{\alpha}p(i); i = 1, \dots, n; \quad (36)$$

Proof. By Proposition 11, there is a permutation q of the index set such that

$$U_{\alpha}^{\beta}U_{\beta}^{\gamma}U_{\gamma}^{\delta}j_{\delta}i = U_{\alpha}^{\delta}j_{\alpha}q(i); i = 1, \dots, n; \quad (37)$$

By definition of U_{α}^{δ} , there is a permutation r of the index set such that $U_{\alpha}^{\delta}j_{\alpha}q(i) = j_{\alpha}r(q(i))$. It suffices to put $p = q^{-1} \circ r$. \square

4 A mathematical realization in the case of a finite-dimensional Hilbert space

It is necessary to show that there exists an example that realized the idea presented in the previous section at least mathematically. We shall show the following theorem.

Theorem 1 Let H be an n -dimensional complex Hilbert space ($2 \leq n < \infty$). Let $\psi \in H$ be an arbitrary quantum state vector. Let C be a set of contexts in H . Then for a sufficiently small positive real number $\epsilon > 0$, there are a symplectic manifold Ω and a subset $B(\epsilon)$ such that for an arbitrary history of context change $\omega = \{\omega_1, \omega_2, \dots\}$ that begins with a fixed context ω_1 , there exist

- i) a one-to-one mapping $T_{\omega_1} : B(\epsilon) \rightarrow \Omega$,
- ii) a probability distribution $P_{\omega_1}^{\psi}$ on Ω that is obtained under the mapping $T_{\omega_1} : B(\epsilon) \rightarrow \Omega$ from a probability distribution (f_i) that is uniform on $B(\epsilon)$ and vanishes outside of $B(\epsilon)$; hence

$$d(f_i) = dP_{\omega_1}^{\psi} \circ T_{\omega_1}^{-1}; \quad (38)$$

- iii) a value assigning map $v_{\omega_1} : \Omega \rightarrow \mathbb{R}^n$ such that the expectation value of $v_{\omega_1}(\hat{O})$ for any observable $\hat{O} \in \mathbb{R}^n$ calculated with $P_{\omega_1}^{\psi}$ coincides with the corresponding one provided by quantum probability theory. And the random variable $v_{\omega_1}(\hat{O})$ assumes only eigenvalues of \hat{O} almost surely on Ω .

Moreover, these value assigning maps satisfy gFUNC and n-TRNS.

The proof consists of six steps. In the first, we construct a symplectic manifold from H and define a volume element. In the second, we define a probability distribution (f_i) on Ω that represents an ensemble prepared before measurements. In the third, we introduce a mapping called a splitting which maps prepared ensemble to the one just before measurements, and define $P_{\omega_1}^{\psi}$ as the results of the splitting. In the fourth, we define value assignments and show that gFUNC is satisfied. In the fifth, we show that quantum mechanical expectation values are reproduced. In the final step, we show that n-TRNS is satisfied.

Step 1.

Let us fix a complete orthonormal system (j_1, \dots, j_n) of H that specifies a context $\omega \in C$. We define functions x^i, y^i ($i = 1, 2, \dots, n$) of H into \mathbb{R} as the real part and the imaginary part of the coefficient of j_i in expansion of a vector of H respectively, for $i = 1, 2, \dots, n$. Then for an arbitrary vector $\psi \in H$,

$$\psi = \sum_{i=1}^n x^i(j_i) + \sum_{i=1}^n iy^i(j_i) \quad (39)$$

Then $(x^1, \dots, x^n; y^1, \dots, y^n)$ becomes a coordinate system of H .

Let us define $\mathbb{R}^{2n} = \mathbb{R}^{2n}$. We denote this coordinate system by ω . Then we define ω as

$$\omega = \sum_{i=1}^n dx^i \wedge dy^i \quad (40)$$

Using this coordinate system, we define a symplectic structure on H by a symplectic form defined by

$$\omega = \sum_{i=1}^n dx^i \wedge dy^i; \quad (41)$$

where \wedge means the exterior product. For the sake of convenience, we use matrix notation: $x = [x^1, \dots, x^n]^T$, $y = [y^1, \dots, y^n]^T$, where T means transpose. Then the symplectic form can be written as

$$\omega = dx^T \wedge dy; \quad (42)$$

Lemma 2 Let U be a unitary transformation of an n -dimensional Hilbert space H . Then U is a canonical transformation of H with respect to the symplectic structure defined by (41).

Proof. Put $z^i = x^i + \sqrt{-1}y^i$ ($i = 1, \dots, n$). For convenience, let us use matrix notation; $z = [z^1, \dots, z^n]^T$, where T represents transpose. Since

$$\begin{aligned} \overline{z^i} dz^i &= x^i \sqrt{-1} y^i dx^i + \sqrt{-1} dy^i \\ &= \frac{1}{2} d(x^i)^2 + \frac{1}{2} d(y^i)^2 - \sqrt{-1} d(x^i y^i) + 2 \sqrt{-1} x^i dy^i; \end{aligned}$$

By taking the exterior derivatives, we obtain

$$d(\overline{z^i} dz^i) = 2 \sqrt{-1} dx^i \wedge dy^i; \quad (43)$$

where y represents complex conjugate transpose.

We denote by U the $n \times n$ matrix whose (i, j) -entry is h_{ij} . It is easy to see that for an arbitrary $j \in \{1, \dots, n\}$

$$z(U^j) = [z^i(U^j)] = [h_{ij}] = \sum_{j=1}^n h_{ij} z^j = U z^j; \quad (44)$$

The pullback of $\overline{z^i} dz^i$ under U becomes

$$\begin{aligned} U^*(\overline{z^i} dz^i) &= (\overline{U z^i}) d(U z^i) \\ &= \overline{z^i} U^* U dz^i \\ &= \overline{z^i} dz^i; \end{aligned}$$

Thus $z^y dz$ is invariant under U . Since the symplectic form is equal to the exterior derivative of $z^y dz$ except a scalar multiple as we saw in (43), U is a canonical transformation of H . \square

We define a volume element $d^n x d^n y$ of H by

$$d^n x d^n y = dx^1 \wedge \dots \wedge dx^n \wedge dy^1 \wedge \dots \wedge dy^n = (-1)^{n(n-1)/2} \frac{1}{n!} \wedge_{i=1}^n : \quad (45)$$

Corollary 3 Let U be a unitary transformation of an n -dimensional Hilbert space H . U preserves the volume element $d^n x d^n y$.

Proof. By Lemma 2, U preserves the symplectic form. Hence the volume element $d^n x d^n y$ which is an n th power of with respect to exterior product is preserved under U . \square

Step 2.

We define an n -ball with a center $f \in H$ by

$$B^{2n}(f; \rho) = \{f \in H : \|f - f\| < \rho\} \quad (46)$$

Let us consider an ensemble of vectors that are distributed uniformly on $B^{2n}(f; \rho)$. Define $B(\rho) = \frac{1}{\mathcal{B}^{2n}(f; \rho)}$. By using $\frac{1}{\mathcal{B}^{2n}(f; \rho)}$, this ensemble is characterized by a probability distribution $\frac{1}{\mathcal{B}^{2n}(f; \rho)}$ on $B^{2n}(f; \rho)$. Using matrix notation of the coordinates $(x; y)$, we can write it as

$$d(f; \rho)((x; y)) = \frac{1}{\mathcal{B}^{2n}(f; \rho)} B(\rho)((x; y)) d^n x d^n y; \quad (47)$$

where $B(\rho)$ is the characteristic function of $B(\rho)$ and $\mathcal{B}^{2n}(f; \rho)$ denotes the volume of $B^{2n}(f; \rho)$.

Step 3.

We call an injection $C_{f; \rho}^{f; \rho}$ of $B^{2n}(f; \rho)$ into H a splitting in a context \mathcal{C} , if there exists a split $\mathcal{D}_i : i = 1, 2, \dots, n$ of $B^{2n}(f; \rho)$, i.e., $B^{2n}(f; \rho) = \bigcup_{i=1}^n \mathcal{D}_i$ and $\mathcal{D}_i \cap \mathcal{D}_j = \emptyset$ for $i \neq j$, and the volume of \mathcal{D}_i is proportional to $\mathcal{B}^{2n}(f; \rho)$, and

$$C_{f; \rho}^{f; \rho}(\mathcal{D}_i) = \mathcal{B}^{2n}(f; \rho) \mathcal{B}^{2n}(f; \rho)^{-1} = 0 \quad (48)$$

($i = 1, 2, \dots, n$). Here, $X \setminus Y$ represents the symmetric difference of subsets X and Y defined by $X \setminus Y = (X \setminus Y) \cup (Y \setminus X)$.

Lemma 3 If $\rho > 0$ is sufficiently small, then there exists a splitting of $B^{2n}(f; \rho)$ in an arbitrary context \mathcal{C} :

Proof. A split \mathcal{D}_i of $B^{2n}(f; \rho)$ is obtained by the following procedure. Let $R_{f; \rho}^{f; \rho}(\theta)$ be a rotation around the direction f with an angle θ (0

$< 2\pi$) in \mathbb{R}^n such that $R_{\mathbf{u}(\mathbf{j}; i)}(2\pi) = \text{the identity mapping}$. Put $g = \frac{1}{2\pi}$.
 $R_{\mathbf{u}(\mathbf{j}; i)}(\cdot) = \cdot$.

Let B_0 be a subset of $B^{2n}(\mathbf{j}; i)$ such that

$$\begin{aligned} g(B_0) \setminus g \circ (B_0) &= \emptyset; \text{ if } \theta \in [0, 2\pi); \\ B^{2n}(\mathbf{j}; i) &= \bigcup_{\theta \in [0, 2\pi)} fg(B_0) : 0 \leq \theta < 2\pi. \end{aligned}$$

Let $\theta_0 (= 0) < \theta_1 < \dots < \theta_n (= 2\pi)$ be an increasing sequence of angles around the direction $\mathbf{u}(\mathbf{j}; i)$ such that the volume of $fg(B_0) : \theta_{i-1} < \theta_i$ is proportional to $\mathbf{h}_i \mathbf{j}_i^2$ ($i = 1; 2; \dots; n$). Put $D_i = fg(B_0) : \theta_{i-1} < \theta_i$ ($i = 1; 2; \dots; n$). For sufficient small ϵ , $B^{2n}(\mathbf{j}; i; \mathbf{h}_i \mathbf{j}_i^{i=n})$ s are pairwise disjoint. Then we can define a one-to-one mapping $C_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i}$ of $B^{2n}(\mathbf{j}; i)$ into H by defining an action of $C_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i}$ on D_i by the successive compositions of a stretching D_i along g -direction so as to make it a $2n$ -dimensional ball, a contraction along radial direction of the ball, and a parallel translation mapping the center of the ball from \mathbf{j}_i to \mathbf{j}_i ($i = 1; \dots; n$). This completes the proof. \square

First, we define $P_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i}$ as the probability distribution characterizing an ensemble that is obtained by an action of the splitting $C_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i}$ in \mathbf{u} (Lemma 3) on the ensemble characterized by $\mathbf{u}(\mathbf{j}; i)$. Since

$$d(\mathbf{j}; i)(\mathbf{x}; \mathbf{y}) = dP_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i} = C_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i}(\mathbf{x}; \mathbf{y}); \quad \int_{\mathbb{R}^{2n}} d(\mathbf{x}; \mathbf{y}) = 1 = \int_H d(\mathbf{h}); \quad (49)$$

$$dP_{\mathbf{u}(\mathbf{j}; i)}^{\mathbf{j}; i}(\mathbf{x}; \mathbf{y}) = \frac{1}{B^{2n}(\mathbf{j}; i)} \prod_{i=1}^n \frac{X^n}{B^{2n}(\mathbf{j}_i; \mathbf{h}_i \mathbf{j}_i^{i=n})} (\mathbf{x}; \mathbf{y}) d^n \mathbf{x} d^n \mathbf{y} \quad (50)$$

holds.

Suppose that up to k th context \mathbf{u} of the history of context change $\mathbf{u}!$, a splitting $C_{\mathbf{u}!}^{\mathbf{j}; i}$ in a context $\mathbf{u}!$ is defined; $P_{\mathbf{u}!}^{\mathbf{j}; i}$ is defined from $\mathbf{u}(\mathbf{j}; i)$ through $C_{\mathbf{u}!}^{\mathbf{j}; i}$. Now our aim is to define $C_{\mathbf{u}!}^{\mathbf{j}; i}$ in the $(k+1)$ th context \mathbf{u} of the history of context change $\mathbf{u}!$.

We denote the split of $B^{2n}(\mathbf{j}; i)$ with respect to the splitting $C_{\mathbf{u}!}^{\mathbf{j}; i}$ by $D_i(\mathbf{u}!) : i = 1; 2; \dots; n$. These $D_i(\mathbf{u}!)$ s may be different from the D_i s defined in the proof of Lemma 3, but by the assumption they satisfy the following:

$$D_i(\mathbf{u}!) \cap D_j(\mathbf{u}!) = \emptyset; \quad \text{if } i \neq j; \quad (51)$$

$$D_i(\mathbf{u}!) \cap D_j(\mathbf{u}!) = \emptyset; \quad \text{if } i \neq j; \quad (52)$$

$$\bigcup_{i=1}^n D_i(\mathbf{u}!) = B^{2n}(\mathbf{j}; i); \quad (53)$$

$$C_{\mathbf{u}!}^{\mathbf{j}; i}(D_i(\mathbf{u}!)) \cap C_{\mathbf{u}!}^{\mathbf{j}; i}(D_j(\mathbf{u}!)) = \emptyset; \quad \text{if } i \neq j; \quad (54)$$

By Lemma 1, there exist the nest partitions $f_{I_k}^j$ and $f_{J_k}^j$ of the index set $f_{1, \dots, n}$ such that $\text{span} f_{j_i} : i \in I_k = \text{span} f_{j_i} : i \in J_k$. By splitting $D_i(\underline{\alpha} \mid \underline{\beta})$ inner if necessary and collecting them, we can define a split $f_{D_i}(\underline{\alpha} \mid \underline{\beta})$ of $D_i(\underline{\alpha} \mid \underline{\beta})$ so that

$$\left[\sum_{j \in J_k} D_j(\underline{\alpha} \mid \underline{\beta}) \right]_{i \in I_k} = \left[\sum_{j \in J_k} D_i(\underline{\alpha} \mid \underline{\beta}) \right]_{i \in I_k} \quad (55)$$

$$D_i(\underline{\alpha} \mid \underline{\beta}) \neq 0 \iff \exists j \in J_k \text{ such that } f_{j_i}^i \neq 0; \quad (56)$$

$$D_i(\underline{\alpha} \mid \underline{\beta}) \setminus D_j(\underline{\alpha} \mid \underline{\beta}) = 0 \text{ if } i \neq j; \quad (57)$$

$$\prod_{i=1}^n D_i(\underline{\alpha} \mid \underline{\beta}) = B^{2n}(j_i); \quad (58)$$

$$C_{\underline{\alpha} \mid \underline{\beta}}^{j_i} D_i(\underline{\alpha} \mid \underline{\beta}) = B^{2n}(j_i); \quad (59)$$

hold. We define the splitting $C_{\underline{\alpha} \mid \underline{\beta}}^{j_i}$ as an injection of $B^{2n}(j_i)$ into H that satisfies

$$C_{\underline{\alpha} \mid \underline{\beta}}^{j_i} D_i(\underline{\alpha} \mid \underline{\beta}) = B^{2n}(j_i); \quad (60)$$

for $i = 1, \dots, n$.

We define $P_{\underline{\alpha} \mid \underline{\beta}}^{j_i}$ as the probability distribution characterizing the ensemble that is obtained as results of action of the splitting $C_{\underline{\alpha} \mid \underline{\beta}}^{j_i}$ on the ensemble characterized by (j_i) . Since

$$d(j_i)(x; y) = dP_{\underline{\alpha} \mid \underline{\beta}}^{j_i} C_{\underline{\alpha} \mid \underline{\beta}}^{j_i}(x; y); \quad \int_{\mathbb{R}^{2n}} d(x; y) = 1 = \int_{\mathbb{R}^{2n}} dP_{\underline{\alpha} \mid \underline{\beta}}^{j_i}(x; y); \quad (61)$$

$$dP_{\underline{\alpha} \mid \underline{\beta}}^{j_i}(x; y) = \frac{1}{B^{2n}(j_i)} \prod_{i=1}^n D_i(\underline{\alpha} \mid \underline{\beta}) (x; y) d^n x d^n y; \quad (62)$$

holds. It suffices to put $T_{\underline{\alpha} \mid \underline{\beta}} = \prod_{i=1}^n C_{\underline{\alpha} \mid \underline{\beta}}^{j_i}$. By repeating these constructions from $k = 1$ we obtain $C_{\underline{\alpha} \mid \underline{\beta}}^{j_i}$, $P_{\underline{\alpha} \mid \underline{\beta}}^{j_i}$, and $T_{\underline{\alpha} \mid \underline{\beta}}$.

Step 4.

For each pair of different contexts $\underline{\alpha}, \underline{\beta} \in C$, as the counterpart of the unitary transformation $U_{\underline{\alpha} \mid \underline{\beta}}$ of context change, we define $\underline{\alpha} \mid \underline{\beta} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ by

$$\underline{\alpha} \mid \underline{\beta} = \underline{\alpha} U_{\underline{\alpha} \mid \underline{\beta}}^{-1}; \quad (63)$$

First, we define a value assigning map $v_{\underline{\alpha}}(\cdot)$ in the context $\underline{\alpha}$ that corresponds to the orthonormal basis $f_{j_1}^1, \dots, f_{j_n}^n$ in the following way. Let \hat{O} be an observable that is stable in $\underline{\alpha}$. We denote the eigenvalue of \hat{O} associated with an eigenvector j_i by α_i . Let $f_{I_1}^0, \dots, f_{I_m}^0$ be a partition of the index set

find i_1, \dots, i_n such that $o_{i_1} = o_{j_1}, \dots, o_{i_n} = o_{j_n}$ for some k . For the sake of convenience, we write $o_{i_k}^0$ instead of o_{i_k} whose index i_k belongs to I_k^0 . We define $v_{\hat{O}}$ by

$$v_{\hat{O}}(\hat{O})(x; y) = \begin{cases} o_{i_k}^0; & \text{if } \hat{O}^{-1}(x; y) \in \text{span}\{j_i : i \in I_k^0\}; \\ \text{otherwise,} & \end{cases} \quad (64)$$

where \hat{O} represents a value depending on $(x; y)$.

For each context $\gamma \in C$, we define $v_{\hat{O}}^{\gamma}$ for \hat{O} / γ by

$$v_{\hat{O}}^{\gamma}(\hat{O})(x; y) = v_{\hat{O}}(U_{\gamma}^{\gamma} \hat{O} U_{\gamma}^{-1})(\hat{O}^{-1}(x; y)); \quad \forall (x; y) \in R^{2n} = \text{span}\{H\}; \quad (65)$$

In more general case, we define value assigning maps successively. Suppose that up to k th context γ_k of the history of context change $\gamma_1, \dots, \gamma_k, v_{\hat{O}}^{\gamma_k}$ is defined. For the $(k+1)$ th context γ_{k+1} , we define $v_{\hat{O}}^{\gamma_{k+1}}$ for \hat{O} / γ_{k+1} by

$$v_{\hat{O}}^{\gamma_{k+1}}(\hat{O})(x; y) = v_{\hat{O}}^{\gamma_k}(U_{\gamma_{k+1}}^{\gamma_{k+1}} \hat{O} U_{\gamma_{k+1}}^{-1})(\hat{O}^{-1}(x; y)); \quad \forall (x; y) \in R^{2n} = \text{span}\{H\}; \quad (66)$$

By definition, these value assigning maps admit context changing maps.

Proposition 12 Let U be a unitary transformation of H .

$$U B^{2n}(\hat{O}; r) = B^{2n}(U \hat{O} U^{-1}; r); \quad (67)$$

where $\hat{O} \in H, r > 0$.

Proof.

$$\begin{aligned} & U B^{2n}(\hat{O}; r) \\ &= U \{ \hat{O} \in H : k \hat{O} \hat{O}^{-1} \hat{O} \hat{O}^{-1} < r \} \\ &= U \{ \hat{O} \in H : k \hat{O} \hat{O}^{-1} \hat{O} \hat{O}^{-1} < r \} \\ &= \{ \hat{O} \in H : k U \hat{O} U^{-1} \hat{O} U^{-1} U \hat{O} U^{-1} < r \} \\ &= \{ \hat{O} \in H : k \hat{O} \hat{O}^{-1} \hat{O} \hat{O}^{-1} < r \} \\ &= B^{2n}(U \hat{O} U^{-1}; r); \end{aligned}$$

□

Proposition 13 The value assigning maps $v_{\hat{O}}^{\gamma_1, \dots, \gamma_k}(\hat{O}) \in T_{\hat{O}}^{\gamma_1, \dots, \gamma_k}$ defined in the above satisfy gFUNC.

Proof. Let \hat{A} and \hat{B} be observables such that there exists a function $f : R \rightarrow R$ which satisfies $\hat{A} = f(\hat{B})$. Suppose that \hat{B} is stable in a context $\gamma \in C$. For a history of context changes $\gamma_1, \dots, \gamma_k, \gamma_{k+1}$, there exist unitary

transformations $U_{i_1}^y; \dots; U_{i_n}^y$ of context changes corresponding to each step. It is easy to see that

$$\begin{aligned} U_{i_1}^y \dots U_{i_n}^y \hat{B} U_{i_1}^y \dots U_{i_n}^y &= \hat{O} / \dots; \\ U_{i_1}^y \dots U_{i_n}^y \hat{A} U_{i_1}^y \dots U_{i_n}^y &= \hat{Q} / \dots; \end{aligned}$$

and

$$\hat{Q} = f(\hat{O});$$

Let $(x; y)$ be an arbitrary point in $B(\cdot) = \bigcup_{i=1}^n B^{2n}(j_i; \dots)$. There exists $i \in \{1, \dots, n\}$ such that $T_{i_1} \dots T_{i_n}((x; y)) \in B^{2n}(j_{i_1}; \dots, j_{i_n})$ in the notation in the step 3. Put $j_i = \dots^1((x; y))$. Let a_i and b_i be eigenvalues of \hat{A} and \hat{B} associated with the eigenvector j_{i_1} , respectively. Then

$$v_{i_1} \dots v_{i_n}(\hat{A}) T_{i_1} \dots T_{i_n}((x; y)) = v_{i_1} \dots v_{i_n}(\hat{A}) \dots C_{i_1}^{j_{i_1}} \dots(j_i) = a_i;$$

In the same way,

$$v_{i_1} \dots v_{i_n}(\hat{B}) T_{i_1} \dots T_{i_n}((x; y)) = b_i;$$

Now our task is to show $a_i = f(b_i)$.

By the definition of $v_{i_1} \dots v_{i_n}$ and (32),

$$\begin{aligned} a_i &= v_{i_1} \dots v_{i_n}(\hat{A}) T_{i_1} \dots T_{i_n}((x; y)) \\ &= v_{i_1} \dots v_{i_n} (U_{i_1}^y \dots U_{i_n}^y \hat{A} U_{i_1}^y \dots U_{i_n}^y) \dots^1 \dots^1 T_{i_1} \dots T_{i_n}((x; y)) \\ &= v_{i_1} \dots v_{i_n}(\hat{Q}) \dots^1 \dots^1 \dots C_{i_1}^{j_{i_1}} \dots(j_i) \\ &= v_{i_1} \dots v_{i_n}(\hat{Q}) \dots (U_{i_1}^y \dots U_{i_n}^y C_{i_1}^{j_{i_1}} \dots(j_i)); \end{aligned}$$

By Corollary 2 and Proposition 12, there exists a permutation p^0 of the index set $\{1, \dots, n\}$ such that $U_{i_1}^y \dots U_{i_n}^y C_{i_1}^{j_{i_1}} \dots(j_i)$ belongs to $B^{2n}(j_{p^0(i)}; \dots, j_{p^0(n)})$.

Hence a_i is an eigenvalue of \hat{Q} associated with an eigenvector $j_{p^0(i)}$. Since $\hat{Q} = f(\hat{O})$, $j_{p^0(i)}$ is an eigenvector of \hat{O} with an eigenvalue o_i such that $a_i = f(o_i)$. By definition of $v_{i_1} \dots v_{i_n}$, the assigned value of \hat{O} is o_i . Therefore

$$\begin{aligned} a_i &= v_{i_1} \dots v_{i_n}(f(\hat{O})) \dots (U_{i_1}^y \dots U_{i_n}^y C_{i_1}^{j_{i_1}} \dots(j_i)) \\ &= f(v_{i_1} \dots v_{i_n}(\hat{O})) \dots (U_{i_1}^y \dots U_{i_n}^y C_{i_1}^{j_{i_1}} \dots(j_i)) \\ &= f(v_{i_1} \dots v_{i_n}(\hat{O}) \dots^1 \dots^1 T_{i_1} \dots T_{i_n}((x; y))) \\ &= f(v_{i_1} \dots v_{i_n} (U_{i_1}^y \dots U_{i_n}^y \hat{O} U_{i_1}^y \dots U_{i_n}^y) \dots T_{i_1} \dots T_{i_n}((x; y))) \\ &= f(v_{i_1} \dots v_{i_n}(\hat{B}) T_{i_1} \dots T_{i_n}((x; y))) \\ &= f(b_i); \end{aligned}$$

□

Step 5.

Proposition 14 Let \mathcal{B} be the Borel σ -algebra of \mathbb{R}^n . $(\Omega; \mathcal{B}; P_{\omega}^{j, i; \omega})$ is a probability space that reproduce quantum-mechanical statistical results, i.e., for an observable $\hat{B} \in \mathcal{A}$,

$$\int_{\Omega} dP_{\omega}^{j, i; \omega} v_{\omega}(\hat{B}) = \text{tr}(\hat{B} \rho_j) \quad (68)$$

Proof. For the history of context change $\omega \rightarrow \omega'$, there exists a sequence of unitary transformations of context changes $U_{\omega}^1, \dots, U_{\omega}^n$ such that $U_{\omega}^1 \dots U_{\omega}^n \hat{B} U_{\omega}^1 \dots U_{\omega}^n =: \hat{O}$ is stable in the context ω . Since $\hat{B} = \sum_{i=1}^n b_i |j_i\rangle \langle j_i|$ by using Corollary 2, we obtain

$$\begin{aligned} \hat{O} &= U_{\omega}^1 \dots U_{\omega}^n \hat{B} U_{\omega}^1 \dots U_{\omega}^n \\ &= \sum_{i=1}^n b_i U_{\omega}^1 \dots U_{\omega}^n |j_i\rangle \langle j_i| U_{\omega}^1 \dots U_{\omega}^n \\ &= \sum_{i=1}^n b_i |j_{p(i)}\rangle \langle j_{p(i)}| \end{aligned}$$

where p represents the permutation of the index set given by Corollary 2.

By the definition of $v_{\omega}(\hat{B})$,

$$v_{\omega}(\hat{B}) = v_{\omega}(\hat{O}) \quad (69)$$

Using Corollary 2 again and (69), we have

$$\begin{aligned} v_{\omega}(\hat{B})(|j_i\rangle) &= v_{\omega}(\hat{O})(U_{\omega}^1 \dots U_{\omega}^n |j_i\rangle) \\ &= v_{\omega}(\hat{O})(|j_{p(i)}\rangle) \\ &= b_i \end{aligned}$$

Therefore by (62)

$$\begin{aligned} &\int_{\Omega} dP_{\omega}^{j, i; \omega} v_{\omega}(\hat{B}) \\ &= \frac{1}{\mathcal{B}^{2n}(\mathcal{J}_i)} \int_{\Omega} d^n x d^n y \sum_{i=1}^n \mathcal{B}^{2n}(\mathcal{J}_i; \mathcal{H}_i \mathcal{J}_i^{j=n}) v_{\omega}(\hat{B}) \\ &= \sum_{i=1}^n b_i \frac{\mathcal{B}^{2n}(\mathcal{J}_i; \mathcal{H}_i \mathcal{J}_i^{j=n})}{\mathcal{B}^{2n}(\mathcal{J}_i)} \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^{X^n} b_i \mathfrak{h}_i \mathfrak{j}_i \mathfrak{j} \\
&= \mathfrak{h} \hat{\mathfrak{B}} \mathfrak{j}_i:
\end{aligned}$$

□

Step 6.

Finally, we show that n-TRNS is satisfied.

Proposition 15 For the history of context change $\underline{\omega} \rightarrow \underline{\omega}' \rightarrow \underline{\omega}''$, let $\hat{\mathcal{O}}$ be a degenerate observable that is stable in both $\underline{\omega}$ and $\underline{\omega}'$. Then

$$v_{\underline{\omega}''}(\hat{\mathcal{O}}) \cdot C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) = v_{\underline{\omega}'}(\hat{\mathcal{O}}) \cdot C_{\underline{\omega}}^{\mathfrak{j}_i}(\mathfrak{j}_i) \quad (70)$$

for $\mathfrak{j}_i \in B^{2n}(\mathfrak{j}_i)$.

Proof. We use the same notations in the step 3. If $\mathfrak{j}_i \in D_i(\underline{\omega} \rightarrow \underline{\omega}')$, $B^{2n}(\mathfrak{j}_i)$, then

$$C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) \in B^{2n}(\mathfrak{j}_i); \mathfrak{h}_i \mathfrak{j}_i \mathfrak{j}^{\mathfrak{i}=n} :$$

By the definition of the value assigning map $v_{\underline{\omega}}(\hat{\mathcal{O}})$, $v_{\underline{\omega}'}(\hat{\mathcal{O}})$ assumes an eigenvalue of $\hat{\mathcal{O}}$, say α_{k_0} , i.e.,

$$\alpha_{k_0} = v_{\underline{\omega}'}(\hat{\mathcal{O}}) \cdot C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) :$$

Let \mathfrak{I}_k and \mathfrak{J}_k be the finest partitions of the index set with respect to the pair of contexts $\underline{\omega}$ and $\underline{\omega}'$ (Lemma 1). Let $U_{\underline{\omega}'}^{\underline{\omega}}$ be the unitary transformation of the context change from $\underline{\omega}$ to $\underline{\omega}'$. There exists k such that $\mathfrak{i} \in \mathfrak{I}_k$. For this k , by (55), there exists $\mathfrak{j} \in \mathfrak{J}_k$ such that $\mathfrak{j}_i = U_{\underline{\omega}'}^{\underline{\omega}}(\mathfrak{j}_i)$ and $\mathfrak{j}_i \in D_j(\underline{\omega} \rightarrow \underline{\omega}') \cap B^{2n}(\mathfrak{j}_i)$. Hence by (60)

$$C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) \in B^{2n}(\mathfrak{j}_i); \mathfrak{h}_j \mathfrak{j}_i \mathfrak{j}^{\mathfrak{i}=n} :$$

By Proposition 12,

$$U_{\underline{\omega}'}^{\underline{\omega}} \cdot C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) \in B^{2n}(\mathfrak{j}_i); \mathfrak{h}_j \mathfrak{j}_i \mathfrak{j}^{\mathfrak{i}=n} :$$

By Proposition 8, $v_{\underline{\omega}'}(\hat{\mathcal{O}}) = v_{\underline{\omega}}(\hat{\mathcal{O}}) \cdot U_{\underline{\omega}'}^{\underline{\omega}}$, and therefore

$$\begin{aligned}
&v_{\underline{\omega}'}(\hat{\mathcal{O}}) \cdot C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) \\
&= v_{\underline{\omega}}(\hat{\mathcal{O}}) \cdot U_{\underline{\omega}'}^{\underline{\omega}} \cdot C_{\underline{\omega}'}^{\mathfrak{j}_i}(\mathfrak{j}_i) \\
&= \alpha_{k_0} :
\end{aligned}$$

□

Hence n-TRNS holds. Thus the proof of the theorem is completed.

5 Discussion

Theorem 1 seems to contradict to the no-go theorem for noncontextual hidden variable models [1] by the following consideration. By the no-go theorem, there exists a finite set of contexts $\underline{f}(\underline{j}) : j = 1; \dots; N$ such that it is impossible to define a value assigning map v for them, if the dimension of the Hilbert space of quantum state vectors is greater than two. For each context $\underline{f}(\underline{j})$, there exists nondegenerate observable $\hat{A}(\underline{j}) / \underline{f}(\underline{j})$. Let \hat{D} be an observable that cannot be assigned a value.

Suppose that \hat{D} is stable in every contexts $\underline{f}(\underline{1}); \dots; \underline{f}(\underline{N})$. Then there exists a function f^j such that $\hat{D} = f^j(\hat{A}(\underline{j}))$. By gFUNC,

$$f^k(v_{\underline{f}(\underline{1})!} \dots \underline{f}(\underline{k})\hat{A}(\underline{k})) = v_{\underline{f}(\underline{1})!} \dots \underline{f}(\underline{k})\hat{D}$$

for an arbitrary history of context change $\underline{f}(\underline{1})! \dots \underline{f}(\underline{k})$ ($k = 1; \dots; N$). By n-TRN,

$$v_{\underline{f}(\underline{1})}(\hat{D}) = \underline{f}(\underline{1})! \dots \underline{f}(\underline{k})\hat{D} = \underline{f}(\underline{1})! \dots \underline{f}(\underline{N})\hat{D};$$

and therefore

$$f^k(v_{\underline{f}(\underline{1})!} \dots \underline{f}(\underline{k})\hat{A}(\underline{k})) = v_{\underline{f}(\underline{1})!} \dots \underline{f}(\underline{k})\hat{D} = v_{\underline{f}(\underline{1})}(\hat{D});$$

If we define a value assigning map v by $v(\hat{O}) = v_{\underline{f}(\underline{1})!} \dots \underline{f}(\underline{k})\hat{O}$ if $\hat{O} / \underline{f}(\underline{k})$, then v satisfies FUNC partially, i.e., for \hat{f} 's ($j = 1; \dots; N$), but this contradicts the assumption for \hat{D} . The root of this contradiction comes from the assumption that $\hat{D} / \underline{f}(\underline{j})$; $\forall j$. Thus there must be a j such that \hat{D} is not stable in $\underline{f}(\underline{j})$.

Acknowledgments

I thank Professor L. Accardi for useful comments and people of the Volterra Center for their warm hospitality. I also thank my colleagues at Hokusei Gakuen University Junior College for their support, especially late Professor Reimei Kobayashi for his warm encouragement.

References

- [1] S. Kochen and E. P. Specker, "The Problem of Hidden Variables in Quantum Mechanics", J. Math. & Mech. 17, 59 (1967).
- [2] M. Redhead, "Incompleteness, Nonlocality, and Realism: a prolegomenon to the philosophy of quantum mechanics", (Clarendon Press, Oxford, 1987)
- [3] A. Peres, "Incompatible results of quantum measurements", Phys. Lett. A 151, 107 (1990).