

When Non-Gaussian States are Gaussian: Generalization of Non-Separability Criterion for Continuous Variables

Derek M. Hughes¹, Vladimír Bužek^{1,2}, and Mario Zimán^{1,2,3}

¹Research Center for Quantum Information, Slovak Academy of Sciences, Dubravská cesta 9, 84511 Bratislava, Slovakia

²Quniverse, L. Štefániho ul. 116, 841 04 Bratislava, Slovakia

³Faculty of Informatics, Masaryk University, Botanická 68a, 60200 Brno, Czech Republic

(Received February 14, 2021)

We present a class of non-Gaussian two-mode continuous variable states for which the separability criterion for Gaussian states can be employed to detect whether they are separable or not. These states reduce to the two-mode Gaussian states as a special case.

PACS numbers: 03.67.-a

Whether a quantum state is entangled or not represents a very important question in quantum information theory. Such knowledge reveals whether one can take advantage of the non-local properties of the state in quantum protocols such as quantum teleportation [1] and quantum cryptography [2]. This issue has been dealt with by many authors in recent years primarily in qubit systems where the Peres-Horodecki partial transpose separability condition was the first method to figure out if a two-qubit state was separable [3]. In general, for N qubits the solution is not known. Continuous variable systems have proven to be an extremely useful setting for quantum cryptography and communication [4]. In these protocols entangled states are required and the question of separability arises naturally. For two-mode systems separability criteria for Gaussian states were established in Refs. [5, 6] which proved to be both necessary and sufficient. More recently necessary and sufficient conditions for the partial transposition of bipartite harmonic quantum states to be nonnegative are formulated in Ref. [7, 8] and separability criteria based on uncertainty relations for two-mode representations of $SU(2)$ and $SU(1,1)$ algebras in Refs. [9, 10, 11] can be obtained from the former result as special cases. These criteria have particularly targeted uncovering whether non-Gaussian states are separable or not as previous criteria fail to detect relatively simple entangled states.

In this paper we approach the problem from a different perspective. The Wigner function of the reduced state of a two-mode quantum state is shown in Fig. 1. Clearly the Wigner function is non-Gaussian and hence the two-mode state from which it is derived is non-Gaussian as well. Say an experimentalist measures (that is reconstructs from a tomographically complete measurement) this state in the laboratory. What exactly can she say about the separability of the state? At first sight not a great deal. However we will show that in fact this state's separability is completely known and understood in terms of its interpretation as a Gaussian state. Furthermore we introduce a whole class of these states for which the usual two-mode Gaussian states are a special case.

Gaussian states of a two-mode continuous variable quantum system have been much studied in the liter-

ature and their entanglement properties are quite well established [12, 13]. Here we briefly review their characterization. Let \hat{a}^\dagger, \hat{a} be the bosonic creation and annihilation operators acting on a Fock Hilbert space as $\hat{a}^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle$ and $\hat{a} |n\rangle = \sqrt{n} |n-1\rangle$ and satisfying the Weyl-Heisenberg commutation relations $[\hat{a}, \hat{a}^\dagger] = \mathbb{I}$. We can introduce position and momentum operators $\hat{x} = \hat{x}^\dagger + \hat{x}$, $\hat{p} = i(\hat{a}^\dagger - \hat{a})$ (setting $\hbar = 1$) which define phase space through the continuous range of their eigenvalues. For two modes these position and momentum operators are defined in each Hilbert space. We gather these operators into a single vector $\hat{\mathbf{x}} = (\hat{x}_1; \hat{p}_1; \hat{x}_2; \hat{p}_2)$ for clarity. By definition the Wigner function of a Gaussian state, ρ , takes the form $W(\mathbf{x}) = \exp[-\mathbf{x}^0 \mathbf{x}^{0T} - 2\mathbf{x}^1] = \frac{1}{\sqrt{\det \hat{\mathcal{C}}}} \exp[-\mathbf{x}^0 \hat{\mathcal{C}}^{-1} \mathbf{x}^0 - 2\mathbf{x}^1]$, where $\mathbf{x}^0 = (x_1, x_2, p_1, p_2)$ and $\hat{\mathcal{C}}_{jk} = \langle \hat{x}_j \hat{x}_k + \hat{x}_k \hat{x}_j \rangle = 2 \langle \hat{x}_j \hat{x}_k \rangle$ is called the covariance matrix with $\hat{\mathcal{C}}_{ii} = \text{Tr}[\hat{\mathcal{O}}_i^2]$. The Wigner function thus depends only on the first and second moments of the position and momentum operators [14]. Furthermore, via local operations the covariance matrix can be brought to the simpler form

$$s.f. = \begin{pmatrix} 0 & b_1 & 0 & c_1 & 0 & 1 \\ 0 & b_2 & 0 & c_2 & 0 & 0 \\ c_1 & 0 & d_1 & 0 & 0 & 0 \\ 0 & c_2 & 0 & d_2 & 0 & 0 \end{pmatrix}; \quad (1)$$

where b_j, c_j and d_j satisfy $\frac{b_j - 1}{d_j - 1} = \frac{b_j + 1}{d_j + 1}$ and $|c_j| \leq \sqrt{(b_j - 1)(d_j - 1)}$ as shown in Ref. [6]. The necessary and sufficient condition for the state to be separable is then

$$h(\hat{u})^2 + h(\hat{v})^2 \leq \frac{1}{\mathcal{C}_0^2}; \quad (2)$$

with $\hat{u} = \frac{c_1}{\sqrt{d_1 - 1}} \hat{x}_1 + \frac{c_2}{\sqrt{d_2 - 1}} \hat{x}_2$, $\hat{v} = \frac{c_1}{\sqrt{d_1 + 1}} \hat{p}_1 + \frac{c_2}{\sqrt{d_2 + 1}} \hat{p}_2$; and $\mathcal{C}_0^2 = \frac{1}{\sqrt{(d_1 - 1)(d_2 - 1)}}$. Actually this expression simplifies to $\frac{1}{\sqrt{(d_1 - 1)(d_2 - 1)}}$ when both $b_1 = 0$ and $d_1 = 1$ or $\frac{1}{\sqrt{(d_1 + 1)(d_2 + 1)}}$ when both $b_1 = 1$ and $d_1 = 1$ otherwise. In the following section we propose an additional criterion: we will relate the degree of entanglement of Gaussian states to an energy of these states.

Gaussian states are an important class of states, but some quantum protocols require [15] the usage of non-

and

$$\langle \hat{X}_1^{(k)} \rangle^2 = \frac{1}{2} \left(\frac{1}{\hat{N}_1 + 1} \hat{a}_1^2 + \hat{a}_1^2 \frac{1}{\hat{N}_1 + 1} \right) \quad (11)$$

with $\hat{N}_1 = \hat{a}_1^\dagger \hat{a}_1$. While we readily concede that it is a non-trivial task to experimentally measure these expectation values we are motivated by what one is able to say about the separability of a state when full tomography has been carried out and the result is a state such as that in Fig. 1.

For a given squeezing parameter r the two-mode squeezed vacuum state $|j\rangle = \frac{1}{\sqrt{1 - e^{-2r}}} \sum_n e^{-nr} |jn\rangle$ and its multi-photon equivalent $|j\rangle = \frac{1}{\sqrt{1 - e^{-2r}}} \sum_n e^{-nr} |jn\rangle_k$ will possess the same degree of entanglement, a fact most easily seen using the von Neumann entropy of the reduced state of either mode, $S = -\text{Tr} \rho \ln \rho = -\text{Tr} \rho_1^{(k)} \ln \rho_1^{(k)} = S^{(k)}$. It should be noted however that uncovering the two-mode entanglement present in each of the above states requires measurement of different observables. In what follows we will compare two states of a two-mode system at a fixed energy value [13]. In terms of the average energy $\langle \hat{H} \rangle = \langle \hat{H}_1 \rangle + \langle \hat{H}_2 \rangle$ with $\hat{N}_i = \hat{a}_i^\dagger \hat{a}_i$ the multi-photon squeezed vacuum state with the same degree of entanglement has average energy $\langle \hat{H} \rangle = k \langle \hat{H}_1 \rangle$. If we fix the energy of both states at $\langle \hat{H} \rangle = 2n$ where $n = \langle \hat{N}_1 \rangle$ then the reduced state of j can be written as $\rho_j = \sum_n p_n |jn\rangle \langle jn|$ with $p_n = \frac{n}{1+n} (1+n)^{-(1+n)}$ and the reduced state of j can be similarly written as $\rho_j^{(k)} = \sum_n p_n^{(k)} |jn\rangle_k \langle jn|_k$ with $p_n^{(k)} = \frac{n}{k} (1 + \frac{n}{k})^{-(1 + \frac{n}{k})}$. Thus it is clear that for fixed energy the usual two-mode squeezed vacuum is more entangled than its multi-photon counterpart. Intuitively this makes sense given that the nature of the multi-photon state does not allow certain quantum correlations to exist; for instance, upon measurement of the state in the joint number basis there is zero probability to obtain the result $(j, k+n)$ for $m < k$ and $n \in \mathbb{Z}^+$. This result also ties in with the fact that among all continuous variable states with a given fixed energy, the maximally entangled states are Gaussian [18].

For mixed states we compare a k -photon two-mode mixed state with a usual two-mode mixed state having the same average energy $\langle \hat{H} \rangle$. To get an intuitive sense of the difference between two such states we look for the minimum purity (defined as $P(\rho) = \text{Tr}(\rho^2)$) allowed for the k -photon mixed state given this energy $\langle \hat{H} \rangle$. The dependence on k is

$$P_{\text{min}}^{(k)} = \frac{1}{\left(\frac{\langle \hat{H} \rangle}{k} + 1\right)^2}; \quad (12)$$

corresponding to a tensor product of two multi-photon thermal states. As k increases the minimum purity increases asymptotically toward 1 so that the k -photon mixed states tend toward the vacuum in the limit $k \rightarrow 1$. To re-enforce this point in Fig. 3 we plot the maximally

entangled multi-photon Gaussian mixed states, see Ref. [13], for various values of k , all at a fixed mean energy $\langle \hat{H} \rangle = 1$. Thus we can say that the maximally entangled k -photon mixed states are less entangled than those for $k = 1$. For general mixed states this statement is not always true.

We have presented a large class of non-Gaussian states for which the existing separability criterion for Gaussian states can be employed in order to detect their entanglement. In order to clarify our results we recall that an arbitrary unitary transformation $U : \mathcal{H} \rightarrow \mathcal{H}$ resulting in an "annihilation" operator $\hat{b}_j = U \hat{a}_j U^\dagger$ can be exploited to define another class of non-Gaussian states $C_U = \{ \rho : \rho = U^\dagger \rho_U U \}$, where G denotes the set of "standard" Gaussian states. Due to the fact that unitary transformations preserve an operator's spectra and the commutation relations, the operators $\hat{b}_j, \hat{b}_j^\dagger$ form a representation of the Weyl-Heisenberg group and Gaussian states with respect to these operators can be defined. In fact, as it was pointed out in Ref. [6] the inequality in Eq.(2) provides a sufficient separability criterion for all operators $\hat{\rho}, \hat{\sigma}$ that are locally unitary equivalent to \hat{a}, \hat{a}^\dagger , i.e. $\hat{\rho} = U \hat{a} U^\dagger, \hat{\sigma} = U \hat{a}^\dagger U^\dagger$, respectively. Moreover, these inequalities provide the necessary conditions for entanglement for all Gaussian states defined with respect to new phase-space coordinates, i.e. for all $\rho \in C_U$.

We have to note that even though the multi-photon non-Gaussian states analyzed in our paper seem to be of the similar form as discussed above, there is a significant difference. The operators \hat{a} and \hat{A}_k are not mutually related by a unitary transformation in the above sense (for more details see Ref. [19]). In fact, the number operators \hat{N}_k and \hat{n} have different spectra (\hat{N}_k is degenerated). The construction in our case is based on the fact that the (semi)infinite Hilbert space of the original harmonic oscillator can be expressed as a finite direct sum of (semi)infinite Hilbert spaces that are isomorphic to the original one, i.e. $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_k$. Here \mathcal{H}_j is a linear span of vectors $|H\rangle = \sum_{j,k} |jn\rangle_k$ ($n = 0, \dots, j-1$). Physically this means that we are restricted to states belonging to the subspace spanned on photon number states separated by a fixed energy k (representing the energy of k photons). The vacuum for \mathcal{H}_j is represented by the state $|j0\rangle = |jj\rangle$. The linear spaces \mathcal{H} and \mathcal{H}_j are related by a non-bijective transformation. However, since \mathcal{H}_j and \mathcal{H}_j^0 are in one-to-one correspondence, we can write $\mathcal{H} = \bigoplus_{j=0}^{k-1} \mathcal{H}_j = \mathcal{H}_k \oplus V_k$ ($\dim \mathcal{H}_k = 1$ and $\dim V_k = k$). Using this notation the multiphoton annihilation operators are unitarily related to the original annihilation operator (acting on \mathcal{H}_k) via the unitary transformation $U = \sum_{n,m} |jn\rangle_k \langle jm|$ ($n, m \in \mathbb{Z}^+$) [19] performing the transformation from \mathcal{H} to $\mathcal{H}_k \oplus V_k$. In particular, $U \hat{A}^{(k)} U^\dagger = \hat{a} \oplus I$.

The Gaussian states are naturally a special case of these non-Gaussian states as one would expect. For two




FIG. 1: (Color online) The Wigner function of the reduced state of a k -photon two-mode squeezed vacuum state as represented in the phase space defined by $\hat{x}_1 = \hat{a}_1^y + \hat{a}$; $\hat{p}_1 = i(\hat{a}_1^y - \hat{a}_1)$ with $k = 3$. This state is non-Gaussian.




FIG. 2: (Color online) The Wigner function of the reduced state of a k -photon two-mode squeezed vacuum state as represented in the phase space defined by $\hat{X}_1^{(k)} = \hat{A}_1^{(k)y} + \hat{A}^{(k)}$; $\hat{P}_1^{(k)} = i(\hat{A}_1^{(k)y} - \hat{A}_1^{(k)})$ with $k = 3$. The state is a thermal state and its Gaussian nature is clearly evident.

modes the operation moving from the basis in which the states have a Gaussian Wigner representation to that in which they don't is local unitary and as such preserves

the entanglement. This holds for \hat{b}_j , but also for multiphoton operators $\hat{A}^{(k)}$, hence the criterion derived for standard Gaussian states can be directly applied to multiphoton Gaussian states as it was demonstrated in the present work. A question remains is how to efficiently verify whether a given state belongs to a certain sector of the Hilbert space \mathcal{H}^j for a given k , or not. The answer can be given by analyzing the expression for the state under consideration in the Fock basis. If the populated (i.e. non-vanishing) levels are separated by the same energy (equivalently, by the same number of photons), then the state belongs to a multiphoton sector \mathcal{H}^k of the Hilbert space \mathcal{H} and its multiphoton Wigner function can be further analyzed.

Acknowledgement

This work was supported in part by the European Union projects INTAS-04-77-7289, CONQUEST and QAP, by




FIG. 3: The maximally entangled mixed states of the multiphoton Gaussian states plotted at the same average energy $\langle \hat{E} \rangle = 1$ (units are dimensionless) for different values of k . The log-negativity, E , is used to measure the entanglement while the purity, P , is used to indicate how mixed the state is.

the Slovak Academy of Sciences via the project CEP I/2/2005, by the project APVT-99-012304.

-
- [1] C. H. Bennett et al, Phys. Rev. Lett. 70, 1895 (1993); S. L. Braunstein et al, Phys. Rev. Lett. 80, 869 (1998).
 - [2] C. H. Bennett et al, Proc. of IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India (IEEE, New York, 1984), p. 175; M. Hillery, Phys. Rev. A 61, 022309 (2000).
 - [3] A. Peres, Phys. Rev. Lett. 77, 1413 (1996).
 - [4] P. van Loock, Fort. der Phys. 50, 1177 (2002).
 - [5] R. Simon, Phys. Rev. Lett. 84, 2726 (2001).
 - [6] L. M. Duan et al, Phys. Rev. Lett. 84, 2722 (2001).
 - [7] E. Schukin et al, Phys. Rev. Lett. 95, 230502 (2005); E. Schukin et al, Phys. Rev. A 74, 030302 (R) (2006).
 - [8] A. M. Iranowicz et al, quant-ph/0605001.
 - [9] G. S. Agarwal and A. Biswas, New J. Phys. 7, 211 (2005).
 - [10] M. Hillery et al, Phys. Rev. Lett. 96, 050503 (2006).
 - [11] H. Nha and J. Kim, Phys. Rev. A 74, 012317 (2006).
 - [12] G. Adesso et al, Phys. Rev. Lett. 92, 087901 (2004); G. Adesso et al, Phys. Rev. A 70, 022318 (2004).
 - [13] D. M. Clough et al, Phys. Rev. A 74, 042303 (2006).
 - [14] J. Marcinkiewicz, Math. Z. 44, 612 (1939).
 - [15] S. Olivares et al, Phys. Rev. A 70, 032112 (2004).
 - [16] R. A. Brandt et al, J. Math. Phys. 10, 1168 (1969).
 - [17] S. Lloyd et al, Phys. Rev. Lett. 82, 1784 (1999).
 - [18] M. M. Wolf et al, Phys. Rev. Lett. 96, 080802 (2006).
 - [19] A. Luis et al, Quantum Opt. 8, 33-41 (1993).