

Reply to Yang et al.'s comment

Nguyen Ba An

Institute of Physics and Electronics, 10 Dao Tan, Thu Le, Ba Dinh, Hanoi, Vietnam

This is to reply to a recent comment by Yang, Yuan and Zhang on "Teleportation of two-quN bit entanglement: Exploiting local resources".

The authors of the recent comment [1] have taken their patience to recheck the calculation in [2] on "Teleportation of two-quN bit entanglement: Exploiting local resources". However, Method 1 in [1] would not be necessary.

The simplest way to make all the formulae in [2] remain valid is just to do the exchange of indices 3 & 4 in the whole text of [2].

Or, as also noticed in [1] (Method 2), one needs just replace $X(Y)$ by $Y(X)$ in the definition (9) in [2], without making the exchange 3 & 4. In the latter option, the states $j_{mn}i_{13}$, with $m = 1;2$ and $n = 0;1;2$, above Eq. (2) in [2] should be changed to

$$j_{10}i_{13} = \frac{1}{\sqrt{3}}(e^{-\frac{2-i}{3}}j_{0i} + e^{\frac{2-i}{3}}j_{1i} + j_{2i})_{13};$$

$$j_{11}i_{13} = \frac{1}{\sqrt{3}}(e^{\frac{2-i}{3}}j_{0i} + j_{1i} + e^{-\frac{2-i}{3}}j_{2i})_{13};$$

$$j_{12}i_{13} = \frac{1}{\sqrt{3}}(j_{0i} + e^{-\frac{2-i}{3}}j_{1i} + e^{\frac{2-i}{3}}j_{2i})_{13};$$

$$j_{20}i_{13} = \frac{1}{\sqrt{3}}(e^{\frac{2-i}{3}}j_{0i} + e^{-\frac{2-i}{3}}j_{1i} + j_{2i})_{13};$$

$$j_{21}i_{13} = \frac{1}{\sqrt{3}}(e^{-\frac{2-i}{3}}j_{0i} + j_{1i} + e^{\frac{2-i}{3}}j_{2i})_{13};$$

$$j_{22}i_{13} = \frac{1}{\sqrt{3}}(j_{0i} + e^{\frac{2-i}{3}}j_{1i} + e^{-\frac{2-i}{3}}j_{2i})_{13};$$

This Reply, thanks to [1], could also serve as an Erratum to [2].

[1] J. Yang, H. Yuan and Z. J. Zhang, quant-ph/0611184.

[2] Nguyen Ba An, Phys. Lett. A 341 (2005) 9.