

Manipulating Multi-qudit Entanglement Witnesses by Using Linear Programming

M .A .Jafarizadeh^{a,b,c}, G .Najarbashi^{a,b,y}, H .Habibian^{a,z}

^aDepartment of Theoretical Physics and Astrophysics, Tabriz University, Tabriz 51664, Iran.

^bInstitute for Studies in Theoretical Physics and Mathematics, Tehran 19395-1795, Iran.

^cResearch Institute for Fundamental Sciences, Tabriz 51664, Iran.

April 17, 2024

E-mail: jafarizadeh@tabrizu.ac.ir

^yE-mail: najarbashi@tabrizu.ac.ir

^zE-mail: hesam-habibian@sicatechdec.com

Abstract

A new class of entanglement witnesses (EWs) called reduction type entanglement witnesses is introduced, which can detect some multi-qudit entangled states including PPT ones with Hilbert space of dimension $d_1 \times d_2 \times \dots \times d_n$. The novelty of this work comes from the fact that the feasible regions turn out to be convex polygons, hence the manipulation of these EWs reduces to linear programming which can be solved exactly by using simplex method. The decomposability and non-decomposability of these EWs are studied and it is shown that it has a close connection with eigenvalues and optimality of EWs. Also using the Jamiołkowski isomorphism, the corresponding possible positive maps, including the generalized reduction maps of Ref. [21], are obtained.

Keywords: Entanglement Witness, Multi-qudit, Optimality, Linear Programming, Feasible Region.

PACS Index: 03.65.Ud

1 Introduction

Quantum entangled states lie at the heart of the rapidly developing field of quantum information science, which encompasses important potential applications such as quantum communication, quantum computation [1, 2, 3]. However, the fundamental nature of entangled states has tantalized physicists since the earliest days of quantum mechanics, and even today is by no means fully understood. One of the most basic problems is that how can one tell if a quantum state is entangled?, and how entangled is it still after some noisy quantum process.

Here we will deal with the states of finite dimensional Hilbert space $H = H_{d_1} \otimes \dots \otimes H_{d_n}$. A density matrix ρ is called unentangled or separable if it can be written as a convex combination of pure product states as $\rho = \sum_j p_j \rho_j^{(1)} \otimes \dots \otimes \rho_j^{(n)}$. If no such convex linear combination exists for a given ρ , then the state is called entangled. Although, in the case of pure states of bipartite systems it is to check whether a given state is, or is not entangled, the question is yet an open problem in the case of mixed states (see the recent good reviews [4, 5, 6, 7]). There have been many efforts in recent years to analyze the separability and quantitative character of quantum entanglement. The Bell inequalities satisfied by a separable system give the first necessary condition for separability [8]. In 1996, Peres made an important step towards proving that, for a separable state, the partial transposition with respect to one subsystem of a bipartite density matrix is positive, $\rho^{T_A} \geq 0$ [9]. By establishing a close connection between positive map theory and separability, Horodecki et al. promptly showed that this is sufficient condition for separability for bipartite systems of $2 \otimes 2$ and $2 \otimes 3$ [10]. Regarding the quantitative character of entanglement, Wootters succeeded in giving an elegant formula to compute the entanglement of formation [11] of $2 \otimes 2$ mixtures, thus giving also a separability criterion [12]. An alternative method to detect entanglement is to construct so-called entanglement witnesses [10, 13, 14, 15] and positive maps [16]. Entanglement witnesses are physical observables that can detect the presence of entanglement. Recently many attempts have been made to use the

convex optimization methods as a robust tools in most quantum information areas including construction of EW s [17, 18, 19, 14, 15].

The main motivation for the present investigation is the remarkable fact that there is no evidence that when and where the manipulation of the EW s reduces to LP which can be solved exactly. For example, the first author and his collaborators have found certain examples of Bell diagonal EW s and generalized d d Choi entanglement witnesses [14, 15], by approximate LP method. Indeed in most cases determining feasible regions needs to use numerical calculation and consequently the problem is solved approximately. Therefore to figure out the problems which can be easily solved exactly in a optimal way and also can be generalized to an arbitrary number of particles with different Hilbert spaces is the main purpose of this paper. To this aim we introduce a new class of $(d_1 \times d_2 \times \dots \times d_n)$ -multi-qudit EW s called reduction type entanglement witnesses (REW s) and show that the computational difficulty in solving such problems reduces to LP which can be solved by simplex method [20]. In these cases the feasible regions are simplexes and the minimum value of optimization problem is achieved on their apexes. On the other hand we show that the EW s corresponding to hyperplanes surrounding feasible regions are optimal. Another advantage of this work is the fact that all REW s can be written in terms of some positive operators and optimal EW s. Also in most cases the decomposability of REW s is rather determined, where two of their eigenvalues plays an important role in this issue. Another important consequences of such EW s are the positive maps including the generalized reduction map [21], which can be obtained from these EW s or their tensor product, via Jamiołkowski isomorphism. Finally a class of entangled density matrix including PPTES are provided which can be detected by such REW s and consequently they determine the non-decomposability of REW s. At the end we carry out in detail some particular cases such as multi-qubit and $2 \times 3 \times 4$ and generalized reduction EW s.

The paper organized as follows: In section 2 we give a brief review of EW s. In section 3 we explain the general scheme of linear programming. In section 4, we introduce a new class of

EWs which can be put in the realm of LP, since their feasible regions are convex polygons (indeed simplexes) which can be exactly determined. In section 5, we show that all EWs corresponding to hyperplanes surrounding the feasible regions are optimal. Section 6 is devoted to some interesting examples such as: multi-qubit REWs, $(d \times d \times \dots \times d)$ multi-qudit REWs and $2 \times 3 \times 4$ REW. Section 7 deals with two examples of LP type, where the first one can be solved exactly by the prescription of this paper. In section 8 we introduce some entangled and PPT states which can be detected by REWs and the decomposability of REWs is discussed. In section 9 by using Jamiołkowski isomorphism, the relation between REWs and positive maps is explained. The paper is ended with a brief conclusion and three appendices.

2 Entanglement witness

As mentioned above in the introduction, one of the pragmatic approach to detect entanglement is to construct entanglement witnesses. Let us briefly recall what these operators are.

Definition 1. A Hermitian operator W is called an entanglement witness detecting the entangled state ρ_e if $\text{Tr}(W \rho_e) < 0$ and $\text{Tr}(W \rho_s) \geq 0$ for all separable state ρ_s .

So, if we have a state ρ and we measure $\text{Tr}(W \rho) < 0$, we can be sure that ρ is entangled. This definition has a clear geometrical meaning. The expectation value of an observable depends linearly on the state. Thus, the set of states where $\text{Tr}(W \rho) = 0$ holds is a hyperplane in the set of all states, cutting this set into two parts. In the part with $\text{Tr}(W \rho) > 0$ lies the set of all separable states, the other part (with $\text{Tr}(W \rho) < 0$) is the set of state detected by W . From this geometrical interpretation it follows that all entangled states can be detected by witness. Indeed for each entangled state ρ_e there exist an entanglement witness detecting it [10].

Definition 2. An EW is decomposable (d-EW) if there exists operators $P; Q_i$ with

$$W = P + Q_1^{T_A} + Q_2^{T_B} + \dots + Q_N^{T_Z} \quad P; Q_i \geq 0 \quad (2.1)$$

where partial transpose taken with respect to some subsystem s and it is non-decomposable if it can not be put in this form [19].

Only non-decomposable EWs can detect PPT entangled states that is those density matrices which have positive partial transposition with respect to each subsystem s [22].

3 Manipulating EWs by LP method

This section deals with basic definitions of linear programming (LP) and general scheme to construct EWs by an exact LP method.

To this aim first we consider a Hermitian operator W with some negative eigenvalues

$$W = \sum_i a_i \rho_i \quad (3.2)$$

where ρ_i are positive operators, with $0 \leq \text{Tr}(\rho_i \rho_s) \leq 1$ for every separable states ρ_s . Note that, the condition $0 \leq \text{Tr}(\rho_i \rho_s) \leq 1$ is not always required. It is used here only to simplify analyzing the problem and pave the way to generalize the prescription to multi-qudits with arbitrary higher dimensions as will be discussed in the following.

The minimum value of $F = \text{Tr}(W \rho_s)$ achieves for pure product state, since every mixed ρ_s can be written as a convex combination of pure product states (due to the convexity of separable region) as $\rho_s = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ with $p_i \geq 0$ and $\sum_i p_i = 1$, hence we have

$$\text{Tr}(\rho_s W) = \sum_i p_i \text{Tr}(W |\psi_i\rangle\langle\psi_i|) \geq C_{\min} \quad (3.3)$$

$$\text{with } C_{\min} = \min_{|\psi\rangle \in D_{\text{prod}}} \text{Tr}(W |\psi\rangle\langle\psi|);$$

where D_{prod} denotes the set of product states. Thus we need to find the pure product state $|\psi_{\min}\rangle$ which minimize $\text{Tr}(W |\psi\rangle\langle\psi|)$. Now, as the pure product state $|\psi\rangle$ varies, the map defined by $P_i = \text{Tr}(\rho_i |\psi\rangle\langle\psi|)$ maps the set D_{prod} into a region inside the hypercube defined by $0 \leq P_i \leq 1 \quad i = 1, 2, \dots, N$. This is not all circumstances. An important and difficult task

is to find the convex region (called feasible region) inside this hypercube which comes from $F_W := \text{Tr}(W_s)$ as s varies on D_{sep} , where D_{sep} denotes the set of separable states. Here in this work we are interested in the EWs with the feasible regions of simplex (or at most convex polygons) types, such that the manipulating these EWs amounts to

$$\begin{aligned} \text{minimize } F_W &:= \text{Tr}(W_s) = \sum_{i=1}^X a_i P_i \\ \text{subject to } &\sum_{i=1}^X (c_{ij} P_i - d_i) = 0 \quad j = 1, 2, \dots \end{aligned} \quad (3.4)$$

where $c_{ij}, d_i, i, j = 1, 2, \dots$ are parameters of hyper-planes surrounding the feasible regions. Therefore, the corresponding boundary points of feasible region will minimize exactly F_W , thus the required task reduces to LP problem which can be solved by the well-known simplex method [20].

4 (d_1, d_2, \dots, d_n) multi-qudit reduction type EWs

In this section we consider n Particles with arbitrary dimensions. Without loss of generality particles can be arranged so that d_1, d_2, \dots, d_n . The discussion of some special cases is postponed to Section 6. We introduce and parameterize the multi-qudit reduction-type entangled witnesses (REW)s labeled by subscript R as

$$W_R^{(n)} = \sum_{S \in N^0} b_S |s\rangle\langle s| + d_1 \sum_{j_2, \dots, j_n} |j_2 \dots j_n\rangle\langle j_2 \dots j_n| + \sum_{S \in N^0} a_S^0 |s\rangle\langle s|; \quad (4.5)$$

where $N^0 = \{2, \dots, n\}$, with $b_j = b_1, a_j^0 = a_1^0$ and $|s\rangle, |s^0\rangle$ defined as

$$|s\rangle = \bigotimes_{i=0}^{X-1} |i\rangle_{i_1} \otimes |i\rangle_{i_2}^{(2)} \otimes \dots \otimes |i\rangle_{i_n}^{(n)} \quad \text{with} \quad |i\rangle_{i_k}^{(k)} = \begin{cases} |i\rangle_{i_k} & \text{if } k \in S \\ |1\rangle_{i_k} & \text{if } k \notin S \end{cases} \quad (4.6)$$

and

$$|s^0\rangle = \bigotimes_{i_1=0}^{X-1} \bigotimes_{i_2=0}^{X-1} \dots \bigotimes_{i_n=0}^{X-1} |i_1 i_2 \dots i_n\rangle \otimes |i_2\rangle_{i_2}^{(2)} \otimes \dots \otimes |i_n\rangle_{i_n}^{(n)} \quad (4.7)$$

with

$$O_{i_k}^{0(k)} = \begin{cases} \sum_{j=1}^8 j_{i_1} i_{i_1} j_{i_2 i_k} & \text{if } k \in S \\ \sum_{j=1}^8 j_{i_k} i_{i_k} j_{\prod_{k \in m=1}^n (1 - i_k i_m)} & \text{if } k \notin S; \quad \exists j = n-3 \text{ and } i_2; \dots; i_n = d_1 - 1 \\ 0 & \text{if } k \notin S; \quad \exists j = n-2 \text{ and } i_2; \dots; i_n = d_1 - 1 \\ \sum_{j=1}^8 j_{i_k} i_{i_k} j_{(1 - i_k i_1)} & \text{otherwise} \end{cases}$$

, respectively and

$$j_{00\dots 0} i = \prod_{i=0}^{d_1-1} j_i^{(1)} i j_i^{(2)} i \dots j_i^{(n)} i \quad (4.8)$$

is maximally entangled state. In this notation we have $j_{00\dots 0} = j_{00\dots 0}^{(0)}$. Obviously for multi-qubit system none of $j_{00\dots 0}^{(0)}$ exists. The number of $P_S^{(0)}$ depends on the dimensions of particles d_i 's and it can take one of the following values

$$m = \begin{cases} \geq P_{\exists j=n-d}^{n-3} C_{\exists j}^{n-1} & \text{if } d_1 = d_2 = \dots = d_n = d \\ \geq P_{\exists j=0}^{n-2} C_{\exists j}^{n-1} (2^{m-1} - 1) & \text{if } d_1 = \dots = d_m < d_{m+1} \dots d_n \end{cases}$$

We introduce the new more convenient parameters $a_1 = b_1 = b$, $a_s = b_s + \prod_{s_0 \in S} b_{s_0}$, instead of b_s 's. In order to turn the observable (4.5) to an EW, we need to choose its parameters in such a way that it becomes a non-positive operator with positive expectation values in any pure product state $j_i = j_i^{(1)} i \dots j_i^{(n)} i$:

Now it is the time to reduce the problem to LP one. In order to determine the feasible region, we need to know the apexes, namely the extremum points, to construct the hyperplanes surrounding the feasible regions. Suppose that j_i be a pure product state with $j_i^{(k)} i = (j_{i_0}^{(k)}; j_{i_1}^{(k)}; \dots; j_{i_{d_k-1}}^{(k)})^T$ and let

$$P_S = \text{Tr}(\sum_{i=0}^{d_1-1} j_i i h j) = \sum_{i=0}^{d_1-1} j_i^{(1)} i j_i^{(2)} i \dots j_i^{(n)} i^2; \quad j_i^{(k)} = \begin{cases} \geq j_i^{(k)} & \text{if } k \in \text{flg}[S] \\ \geq 1 & \text{if } k \notin \text{flg}[S] \end{cases}$$

$$P_{2;\dots;n} = d_1 \sum_{i=0}^{d_1-1} j_i i j_i^2 = \sum_{i=0}^{d_1-1} j_i^{(1)} i j_i^{(2)} i \dots j_i^{(n)} i^2$$

$$P_S^{(0)} = \text{Tr}(\sum_{i=0}^{d_1-1} j_i i h j);$$

which all lie in the interval $[0;1]$ (see appendix I). The number of P_S is $2^{n-1}-1$ (cardinality of the power set of N^0 excepted $f;g$).

The extremum points or apexes consist of

a. origin: $P_S = 0; P_{S^0} = 0 \ 8 S \in N^0$ which corresponds to the following choice of pure product state

$$j^{(1)}_i = (1 \ 0 \ 0 \ \dots 0)^T$$

$$j^{(k)}_i = (0 \ 1 \ 0 \ \dots 0)^T \quad k \in N^0$$

b. $P_S = 1; P_{S^0} = 1 \ 8 S^0 \in S; P_{N^0} = 0; P_{S^0} = 0 \ 8 S^0 \in N^0 \setminus S; P_S = 0 \ 8 S \in N^0$ which can be reached by choosing the following pure product states

$$j^{(k)}_i = (1 \ 0 \ 0 \ \dots 0)^T \quad k \in \text{flg}[S];$$

$$j^{(k)}_i = (0 \ 1 \ 0 \ \dots 0)^T \quad k \notin \text{flg}[S];$$

Obviously if $P_S = 1$, then for all $S^0 \in S, P_{S^0} = 1$; thus for $S = N^0$, we get the following important apex

c. $P_S = 1; P_{S^0} = 0 \ 8 S \in N^0$:

which can be obtained by choosing

$$j^{(k)}_i = (1 \ 0 \ 0 \ \dots 0)^T \quad 8 k \in \text{flg}[N^0];$$

d. $P_S^0 = 1; P_{S^0} = 1 \ 8 S^0 \in S \setminus N^0$; the others are zero:

The last category arise from pure product state with the components of the form $\begin{smallmatrix} (1) \\ 0 \end{smallmatrix} = \begin{smallmatrix} (k) \\ 0 \end{smallmatrix} = 1$, if $k \in S$ and $\begin{smallmatrix} (k) \\ k-1 \end{smallmatrix} = 1$, if $k \notin S$, i.e.,

$$j^{(k)}_i = (1 \ 0 \ 0 \ \dots 0)^T \quad k \in \text{flg}[S];$$

$$j^{(k)}_i = (0 \ 0 \ \dots 0 \mid \begin{smallmatrix} 1 \\ \{z\} \end{smallmatrix} \ 0 \ \dots 0)^T \quad k \notin \text{flg}[S];$$

$k-1^{\text{th}}$

Regarding the above consideration, we are now ready to state the feasible region.

To this aim we first prove that, $2^{n-1} + m$ extremum points obtained above form the apexes of $(2^{n-1} + m)$ -simplex in Euclidean space of dimension $N = 2^{n-1} + m - 1$. To this purpose we consider the convex hull of these points, i.e. draw $N + 1$ hyperplanes passing through each combination of N points out of $N + 1$ ones ($C_N^{N+1} = N + 1$). Now, we get a bounded region formed from their intersection which is the required feasible region of $W_R^{(n)}$ and it is obviously a $(2^{n-1} + m)$ -simplex. It is strait forward to show that the feasible region can be obtained by taking the expectation values of

$$^{(S)}W_{opt}^{(n)} = a_s @_s + \sum_{s \in S \subseteq N^0} (1)^{\beta_j \beta_{s^0 j}} + d_1 (1)^{\beta_j \beta_{N^0 j}} j_{00 \dots 0} j_{00 \dots 0}^0 A_s S \in N^0 \quad (4.9)$$

in pure product states (see appendix II). Now, in order that $W_R^{(n)}$ to be an EW, the following expectation values

$$F(P_2; P_3; \dots; P_1^0; P_2^0; \dots) = \text{Tr}(W_R^{(n)} j_{ih} j) = \sum_{s \in N^0} b_s P_s + b_{2, \dots, m} P_{2, \dots, m} + \sum_{s \in N^0} a_s^0 P_s^0 \quad (4.10)$$

must be positive. So our task is to solve the following LP problem

$$\begin{aligned} & \text{minimize} \quad \sum_{s \in N^0} b_s P_s + b_{2, \dots, m} P_{2, \dots, m} + \sum_{s \in N^0} a_s^0 P_s^0 \\ & \text{subject to} \quad \begin{aligned} & \geq P_s - P_s^0 + \sum_{s \in S^0} (1)^{\beta_j \beta_{s^0 j}} P_{s^0} = 0 \\ & \geq \sum_{s \in N^0} P_s^0 = 0 \end{aligned} \end{aligned} \quad (4.11)$$

Putting the coordinates of apexes in Eq.(4.10) yields all $a_s = 0$ and $a_s + a_s^0 = 0$. As we stated in previous section, all P_s and P_s^0 lie in the closed interval $[0; 1]$. Now, F is a linear function of P_s and P_s^0 and if we require it to be positive on the apexes (which are extremum points), then it will be positive in the whole feasible region.

At the end we need to know all eigenvalues of $W_R^{(n)}$ which consist of $a_s; a_s + a_s^0; !_1 = a_{N^0}$, b_{N^0} and $!_2 = d_1 a_{N^0} - (d_1 - 1)!_1$. Since $a_s; a_s + a_s^0 = 0$, then one of the remaining eigenvalues: $!_1$ and $!_2$ must be negative to guarantee $W_R^{(n)}$ to be an EW.

5 Optimality of $^{(S)}W_{\text{opt}}^{(n)}$

After determining the feasible regions, one needs to know whether the boundary of EWs is formed by optimal EWs. An EW is optimal if a positive operator P is subtracted from that then it will be no longer an EW [13].

Note that the EWs corresponding to hyperplanes surrounding feasible regions of $W_R^{(n)}$ are optimal since they cover the simplex feasible region in an optimal way (see appendix III). Thus, the structure of the optimal EWs $^{(S)}W_{\text{opt}}^{(n)}$ characterizes the boundary of REWs $W_R^{(n)}$. In fact, from the results of this Section it will become clear that we can restrict ourselves to the structure of the optimal EWs corresponding to hyperplanes surrounding feasible regions. In other words optimal EWs $^{(S)}W_{\text{opt}}^{(n)}$ are tangent to the boundary between separable and non-separable states.

Another advantage of $^{(S)}W_{\text{opt}}^{(n)}$ is that one can rewrite the $W_R^{(n)}$ in terms of positive operators

$\sum_{j_1, \dots, j_n} d_{j_1, \dots, j_n} |j_1 \dots j_n\rangle\langle j_1 \dots j_n|$ and some optimal EWs, i.e.,

$$W_R^{(n)} = \sum_S a_S {}^{(S)}W_{\text{opt}}^{(n)} + \sum_{j_1, \dots, j_n} d_{j_1, \dots, j_n} |j_1 \dots j_n\rangle\langle j_1 \dots j_n| + \sum_S (a_S + a_S^0) \rho_S^0 \quad S \in N^0 \quad (5.12)$$

Therefore for positive d_{j_1, \dots, j_n} , the REWs can be decomposed as

$$W_R^{(n)} = \sum_S {}^{(S)}Q^{T(N^0)} + P \quad (5.13)$$

where ${}^{(S)}Q = {}^{(S)}W_{\text{opt}}^{(n)T(N^0)}$ and P is positive operator and in this case $W_R^{(n)}$ can not detect PPT entangled states (non-separable density matrices with positive partial transpose with respect to all particles).

6 Some special cases of $W_R^{(n)}$

In this section we discuss some special cases of REWs to enlighten the subject.

As we see the number of P_S is three, while the number of optimal points is four. Therefore, there are four hyperplanes surrounding the feasible region, where the hyperplanes pass through each combination of three points out of four ones. (see Fig.1). Thus the problem can be reduced to

$$\begin{aligned}
 & \text{minimize} && \text{Tr}(W_R^{(3)} j_{ih} j) \\
 & && 8 \\
 & && \text{~~~~~} \\
 & \text{subject to} && 1 + P_{2;3} - P_2 - P_3 = 0; \\
 & && P_2 - P_{2;3} = 0; \\
 & && \text{~~~~~} \\
 & && P_3 - P_{2;3} = 0; \\
 & && \text{~~~~~} \\
 & && P_{2;3} = 0;
 \end{aligned} \tag{6.16}$$

where the above given inequalities follow rather easily by taking the expectation value of the following optimal EWs

$$\begin{aligned}
 {}^2W_{\text{opt}}^{(3)} &= a_2 (\begin{smallmatrix} 2 & & \\ & j_{000} & ih_{000} \end{smallmatrix} j) \\
 {}^3W_{\text{opt}}^{(3)} &= a_3 (\begin{smallmatrix} 3 & & \\ & j_{000} & ih_{000} \end{smallmatrix} j) \\
 {}^1W_{\text{opt}}^{(3)} &= a_1 (I_8 \begin{smallmatrix} 2 & & \\ & j_{000} & ih_{000} \end{smallmatrix} j)
 \end{aligned} \tag{6.17}$$

in pure product states, and as usual the optimal EWs can be written as partial transpose of the following positive operators

$$\begin{aligned}
 {}^1W_{\text{opt}}^{(3)T_{23}} &= a_1 (j_{100}i + j_{011}i)(h_{100}j + h_{011}j); \\
 {}^2W_{\text{opt}}^{(3)T_3} &= a_2 (j_{011}i + j_{110}i)(h_{001}j + h_{110}j); \\
 {}^3W_{\text{opt}}^{(3)T_2} &= a_3 (j_{010}i + j_{101}i)(h_{010}j + h_{101}j);
 \end{aligned} \tag{6.18}$$

respectively. Now the EW $W_R^{(3)}$ can be written in terms of ${}^iW_{\text{opt}}$ as

$$\begin{aligned}
 W_R^{(3)} &= a_1 {}^1W_{\text{opt}} + a_2 {}^2W_{\text{opt}} + a_3 {}^3W_{\text{opt}} + 2 !_2 j_{000} ih_{000} j \\
 &= a_1 ({}^{(1)}Q^{T_{(23)}} + a_2 ({}^{(2)}Q^{T_{(3)}} + a_3 ({}^{(3)}Q^{T_{(2)}} + 2 !_2 P
 \end{aligned} \tag{6.19}$$

As mentioned above in section 4, for $!_2$ the EW $W_R^{(3)}$ becomes decomposable and can not detect PPT entangled states.

$$j_{102ih102j} + j_{103ih103j} + j_{120ih120j} + j_{122ih122j} + j_{123ih123j}$$

$$_2^0 = j_{02ih002j} + j_{03ih003j} + j_{112ih112j} + j_{113ih113j}$$

$$_3^0 = j_{20ih020j} + j_{21ih121j}$$

Here all possible $_s^0$ ($S = f_2g; f_3g$ and $;$) can be appear. In this case feasible region lies in a space of dimension six. The coordinates of apexes and relevant pure product states j_i are

$$(P_2; P_3; P_{23}; P_1^0; P_2^0; P_3^0)$$

$j_i =$ <table border="0"> <tr> <td>0</td><td>1</td><td>0</td><td>1</td><td>0</td><td>1</td> </tr> <tr> <td>B</td><td>1</td><td>B</td><td>0</td><td>B</td><td>0</td> </tr> <tr> <td>@</td><td>C</td><td>@</td><td>C</td><td>@</td><td>C</td> </tr> <tr> <td></td><td>A</td><td></td><td>A</td><td></td><td>A</td> </tr> <tr> <td>0</td><td></td><td>0</td><td></td><td>0</td><td></td> </tr> </table>	0	1	0	1	0	1	B	1	B	0	B	0	@	C	@	C	@	C		A		A		A	0		0		0		!	(0;0;0;0;0;0)
0	1	0	1	0	1																											
B	1	B	0	B	0																											
@	C	@	C	@	C																											
	A		A		A																											
0		0		0																												
$j_i =$ <table border="0"> <tr> <td>0</td><td>1</td><td>0</td><td>1</td><td>0</td><td>1</td> </tr> <tr> <td>B</td><td>1</td><td>B</td><td>1</td><td>B</td><td>0</td> </tr> <tr> <td>@</td><td>C</td><td>@</td><td>C</td><td>@</td><td>C</td> </tr> <tr> <td></td><td>A</td><td></td><td>A</td><td></td><td>A</td> </tr> <tr> <td>0</td><td></td><td>0</td><td></td><td>0</td><td></td> </tr> </table>	0	1	0	1	0	1	B	1	B	1	B	0	@	C	@	C	@	C		A		A		A	0		0		0		!	(1;0;0;0;0;0)
0	1	0	1	0	1																											
B	1	B	1	B	0																											
@	C	@	C	@	C																											
	A		A		A																											
0		0		0																												
$j_i =$ <table border="0"> <tr> <td>0</td><td>1</td><td>0</td><td>1</td><td>0</td><td>1</td> </tr> <tr> <td>B</td><td>1</td><td>B</td><td>0</td><td>B</td><td>1</td> </tr> <tr> <td>@</td><td>C</td><td>@</td><td>C</td><td>@</td><td>C</td> </tr> <tr> <td></td><td>A</td><td></td><td>A</td><td></td><td>A</td> </tr> <tr> <td>0</td><td></td><td>0</td><td></td><td>0</td><td></td> </tr> </table>	0	1	0	1	0	1	B	1	B	0	B	1	@	C	@	C	@	C		A		A		A	0		0		0		!	(0;1;0;0;0;0)
0	1	0	1	0	1																											
B	1	B	0	B	1																											
@	C	@	C	@	C																											
	A		A		A																											
0		0		0																												
$j_i =$ <table border="0"> <tr> <td>0</td><td>1</td><td>0</td><td>1</td><td>0</td><td>1</td> </tr> <tr> <td>B</td><td>1</td><td>B</td><td>1</td><td>B</td><td>1</td> </tr> <tr> <td>@</td><td>C</td><td>@</td><td>C</td><td>@</td><td>C</td> </tr> <tr> <td></td><td>A</td><td></td><td>A</td><td></td><td>A</td> </tr> <tr> <td>0</td><td></td><td>0</td><td></td><td>0</td><td></td> </tr> </table>	0	1	0	1	0	1	B	1	B	1	B	1	@	C	@	C	@	C		A		A		A	0		0		0		!	(1;1;1;0;0;0)
0	1	0	1	0	1																											
B	1	B	1	B	1																											
@	C	@	C	@	C																											
	A		A		A																											
0		0		0																												

$$\begin{aligned} {}^2W_{\text{opt}} &= a_2 \left(\begin{array}{cc} 2 & j_{000} \\ 2 & j_{000} \end{array} \begin{array}{c} i h \\ i h \end{array} \begin{array}{c} 0 \\ 2 \end{array} \right); \\ {}^3W_{\text{opt}} &= a_3 \left(\begin{array}{cc} 2 & j_{000} \\ 3 & j_{000} \end{array} \begin{array}{c} i h \\ i h \end{array} \begin{array}{c} 0 \\ 3 \end{array} \right); \\ {}^1W_{\text{opt}} &= a_1 \left(\begin{array}{cc} I_{24} & 2 \\ 2 & j_{000} \end{array} \begin{array}{c} 3 \\ 3 \end{array} + \begin{array}{cc} 2 & j_{000} \\ 2 & j_{000} \end{array} \begin{array}{c} i h \\ i h \end{array} \begin{array}{c} 0 \\ 1 \end{array} \right); \end{aligned} \quad (6.22)$$

Taking the partial transposition of $W_{\text{opt}, i=1;2;3}$ with respect to $f_2;3$ gives yields

$$\begin{aligned} {}^1W_{\text{opt}}^{T_{23}} &= a_1 (j_{100i} + j_{011i}) (h_{100j} + h_{011j}); \\ {}^2W_{\text{opt}}^{T_3} &= a_2 (j_{001i} + j_{110i}) (h_{001j} + h_{110j}); \\ {}^3W_{\text{opt}}^{T_2} &= a_3 (j_{010i} + j_{101i}) (h_{010j} + h_{101j}); \end{aligned} \quad (6.23)$$

respectively. Evidently these EW's are optimal, since these have been written as the partial transposition of pure maximally entangled states.

7 Bell-diagonal EW's by LP methods

Recently multi-qubit Bell decomposable entangled witnesses (BDEW's) [14] have been introduced as

$$W_{\text{BD}} = \sum_{i_1 i_2 \dots i_n = 0;1} a_{i_1 i_2 \dots i_n} j_{i_1 i_2 \dots i_n} i_{i_1 i_2 \dots i_n} j \quad (7.24)$$

where $j_{i_1 i_2 \dots i_n} i$ ($d_i = 2; i = 1; 2; \dots; n$) are n -qubit maximally entangled orthonormal states, i.e.,

$$j_{i_1 i_2 \dots i_n} i = \begin{pmatrix} z \end{pmatrix}^{i_1} \begin{pmatrix} x \end{pmatrix}^{i_2} \dots \begin{pmatrix} x \end{pmatrix}^{i_n} j_{00 \dots 0} i \quad (7.25)$$

where x and z are usual Pauli matrices.

In general it is hard to find the BDEW's with feasible region of simplex type or even polygon type, namely those which can be manipulated by LP method. Here we give two examples which are both set in LP problem, where only one of them (the first example) can be solved exactly by the prescription of this paper.

The first example is EW of the form

$$W_1 = aI_{2^n} + 2(b-a)j_{00 \dots 0} i_{00 \dots 0} j + 2(c-a)j_{00 \dots 01} i_{00 \dots 01} j + (d-a) \quad (7.26)$$

where

$$j_{00 \dots 01} i_{00 \dots 01} j = I_{2^n} - f(j_{011 \dots 110} i_{011 \dots 110} j + j_{11 \dots 110} i_{11 \dots 110} j) + (j_{011 \dots 11} i_{011 \dots 11} j + j_{11 \dots 111} i_{11 \dots 111} j)$$

The eigenvalues of W_1 are $a; 2b - a; 2c - a; d$. This BDEW is similar to the one introduced in [14], where the extra term is added to optimize the EWs corresponding to the boundary plane of feasible region. Suppose that $P_{00\dots0} = 2^{-n} |j\rangle\langle j|$, $P_{00\dots01} = 2^{-n} |j\rangle\langle j|$ and $P = \text{Tr}(|j\rangle\langle j|)$. Then the pure product states which produce the apexes are

and consequently these yield the following hyperplanes surrounding the feasible region (see Fig 2)

The positivity of the last constraint come from the positivity of the expectation value of the following optimal EW

in pure product states ρ_{ij} , since it can be written as the partial transpose of a positive operator with respect to the i th particle as

$$W_{\text{opt}}^{T_1} = 2(j_{11 \dots 10} i_{11 \dots 10} j + j_{11 \dots 11} i_{11 \dots 11} j):$$

Now the remaining task is to solve the following LP problem

$$\begin{aligned}
 & \text{minimize} \quad a + (b-a)P_{00\dots 0} + (c-a)P_{00\dots 01} + (d-a)P_{00\dots 010} \\
 & \text{subject to} \quad P_{00\dots 0} + P_{00\dots 01} + P_{00\dots 010} = 1 \\
 & \quad P_{00\dots 0}, P_{00\dots 01}, P_{00\dots 010} \geq 0
 \end{aligned} \tag{7.27}$$

Thus, above problem is reduced to LP and can be solved by simplex method. Putting the apexes in Eq.(7.26) we deduce that a, b, c, d should be positive. Now, the operator W_1 fulfills the properties of EWs if at least one of its eigenvalues is negative, namely $2b-a < 0$ or $2c-a < 0$.

The second example which sets in LP problem is

$$W_2 = aI_{2^n} + 2^{n-1}(b-a)\sum_{00\dots 0} |0\rangle\langle 0| + 2^{n-1}(c-a)\sum_{10\dots 0} |1\rangle\langle 1|$$

which can not be solved by the prescription of this paper. Its feasible region can be determined by Lagrangian multiplier method, as it is discussed in [14]. Let $P_{00\dots 0} = \sum_{00\dots 0} |0\rangle\langle 0|$, $P_{10\dots 0} = \sum_{10\dots 0} |1\rangle\langle 1|$. There the problem reduces to

$$\begin{aligned}
 & \text{minimize} \quad a + 2^{n-2}(b-a)P_{00\dots 0} + 2^{n-2}(c-a)P_{10\dots 0} \\
 & \text{subject to} \quad \begin{cases} \frac{1}{2^{n-2}}P_{00\dots 0} + (1 - \frac{1}{2^{n-2}})P_{10\dots 0} = 0 \\ \frac{1}{2^{n-2}}P_{10\dots 0} + (1 - \frac{1}{2^{n-2}})P_{00\dots 0} = 0 \\ P_{00\dots 0}, P_{10\dots 0} \geq 0 \end{cases}
 \end{aligned} \tag{7.28}$$

These constraints can not be reached by partial transposition approach, and the feasible region is estimated by convex hull of apexes (see Fig.3).

8 Detecting some entangled states by $W_R^{(n)}$

This section is devoted to some entangled states which can be detected by general $W_R^{(n)}$ and three-qubit REW $W_R^{(3)}$. First we consider some Bell states. All of the Bell states $|j\rangle_{10\dots 0}$ can

be detected by $W_R^{(n)}$, since we have

$$\text{Tr}(W_R^{(n)} \rho_{00\dots 0} \otimes \rho_{00\dots 0}) = \lambda_2;$$

$$\text{Tr}(W_R^{(n)} \rho_{i_1 0\dots 0} \otimes \rho_{i_1 0\dots 0}) = \lambda_1 \quad ; \quad i_1 \neq 0;$$

therefore for $\lambda_2 < 0$ one can detect $\rho_{00\dots 0}$ and for $\lambda_1 < 0$ the others modulated Bell states can be detected by $W_R^{(n)}$. On the other hand, imposing some constraints on operator

$$\rho_{i_1 0\dots 0} = \frac{1}{B \text{Tr}(\rho_s) + D d_1} \left(B \left(\sum_{j=0}^{d-1} \rho_{j 0\dots 0} \otimes \rho_{j 0\dots 0} \right) + D d_1 \rho_{i_1 0\dots 0} \otimes \rho_{i_1 0\dots 0} \right); \quad (8.29)$$

one can get a density matrix which can be detected by $W_R^{(n)}$, where ρ_s denotes the separable state inside the parenthesis on the righthand side. The positivity of $\rho_{i_1 0\dots 0}$ constrains B and D to be positive and in order to detect both $\rho_{0 0\dots 0}$ and $\rho_{i_1 0\dots 0}; i_1 \neq 0$, we must have

$$B \text{Tr}(W_R^{(n)} \rho_s) + D d_1 \lambda_2 < 0; \quad (8.30)$$

$$B \text{Tr}(W_R^{(n)} \rho_s) + D d_1 \lambda_1 < 0; \quad (8.31)$$

respectively. Because of the positivity of D , Eq.(8.30) is satisfied if $\lambda_2 < 0$ and Eq.(8.31) is satisfied if $\lambda_1 < 0$. Now, we can proceed our discussion further to detect PPT entangled states which is useful for determining non-decomposable region. Here we do not discuss the decomposability and non-decomposability issues in detail, since it needs the other opportunity and comes elsewhere. $\rho_{i_1 0\dots 0}$ is PPT states with respect to any subsystems of the particles, if B and D be positive, so the Eq.(8.30) yields

$$1 - \frac{B}{D} < \frac{d_1 \lambda_2}{\text{Tr}(W_R^{(n)} \rho_s)} \Rightarrow \lambda_2 < \$ \quad (8.32)$$

where

$$\$ = \frac{1}{d_1} \text{Tr}(W_R^{(n)} \rho_s) \quad (8.33)$$

The above requirement makes $\rho_{0 0\dots 0}$ a class of PPT entangled state which can be detected by $W_R^{(n)}$. But to detect the other $\rho_{i_1 0\dots 0}$ we must have $\lambda_1 < \$$ which is impossible. This is in

agreement with the discussion made in section 5. Now, combining thus obtained results with those of Eq. (5.12) which implies that $W_R^{(n)}$ is decomposable provided that $!_2$ is positive, one can rather determine the decomposability and non-decomposability of REW's (see Fig.4).

Furthermore one can construct some entangled states which can be detected by particular REW's. As an example consider entangled density matrices

$$\begin{aligned} &= \frac{1}{4B + 2D} (B_{-2} + 2D |j_{000}\rangle \langle j_{000}|) \\ &^0 = \frac{1}{4B + 2D} (B_{-3} + 2D |j_{000}\rangle \langle j_{000}|) \end{aligned}$$

which can be detected by three-qubit REW (6.15), with some constraints. The positivity of these states implies that $B; B + 2D \geq 0$ and the positivity of $T_3; \mathbb{Q}_2$ is achieved if $B \geq D \geq 0$. In order that, $\text{Tr}(W_R^{(3)})$ to be negative we should have $B(a_{2,3} + a_2) + D!_2 < 0$, where we have two possibilities: for $D > 0$ we have $!_2 < 0$ and for $D < 0$ we have $!_2 > 0$.

9 Positive maps

As it is shown in [23], there is a close connection between the positive maps and the entanglement witnesses, i.e., the Jamiołkowski isomorphism

$$d_1 d_2 \dots d_n (I_{d_1 \dots d_n} \otimes E) |j\rangle \langle j| + |h\rangle \langle h| = W_{d_1 d_1^0 \dots d_n d_n^0}^{(1,2;\dots,2n)}; \quad d_i = d_i^0; \quad i = 1, \dots, n \quad (9.34)$$

$$E(\cdot) = \text{Tr}_{(1,3;\dots,2n-1)} W_{d_1 d_1^0 \dots d_n d_n^0}^{(1,2;\dots,2n)} (\text{Tr}_{d_1 d_2^0 \dots d_n^0} I_{d_1 d_2^0 \dots d_n^0}) \quad (9.35)$$

where

$$|j\rangle \langle i| = \frac{1}{d_1 d_2 \dots d_n} \sum_{i_1=0}^{d_1-1} \sum_{i_2=0}^{d_2-1} \dots \sum_{i_n=0}^{d_n-1} |j_{i_1}^{(1)} i_{i_1}^{(2)} i_{i_2}^{(3)} i_{i_2}^{(4)} \dots i_{i_n}^{(2n-1)} i_{i_n}^{(2n)}\rangle \langle i| \quad (9.36)$$

is the maximally entangled state in $H_{d_1}^{(1)} \otimes H_{d_1^0}^{(2)} \otimes \dots \otimes H_{d_n}^{(2n-1)} \otimes H_{d_n^0}^{(2n)}$. Hence the Jamiołkowski isomorphism is a one-to-one mapping between the set of trace preserving quantum operations

$$E: H_{d_1}^{(1)} \otimes H_{d_2}^{(3)} \otimes \dots \otimes H_{d_{n-1}}^{(2n-1)} \rightarrow H_{d_1^0}^{(2)} \otimes H_{d_2^0}^{(4)} \otimes \dots \otimes H_{d_{n-1}^0}^{(2n)} \quad (9.37)$$

and $d_1 = d^0 \dots d_n = d_n^0$ EW s if $d_i = d_i^0$ for $i = 1, 2, \dots, n$.

Now, using the Jamiołkowski isomorphism (9.35), we try to construct the positive maps connected with REW s. Evidently tensor product of some EW s is also an EW in higher dimension.

To be more precise let $W_{d_1 d_1^0 \dots d_n d_n^0}^{(1, 2, \dots, 2n)}$ be an EW acting on $H_{d_1}^{(1)} \otimes H_{d_1^0}^{(2)} \dots \otimes H_{d_n}^{(2n-1)} \otimes H_{d_n^0}^{(2n)}$ then depending on possible partition

$$n = n_1 + n_2 + \dots + n_m; \quad n_i \geq 1 \quad (9.38)$$

one can construct an EW by tensor product of REW s as

$$W_{d_1 d_1^0 \dots d_n d_n^0}^{(1, 2, \dots, 2n)} = W_{d_1 d_1^0 \dots d_{n_1} d_{n_1}^0}^{(1, 2, \dots, 2n_1)} \otimes W_{d_{n_1+1} d_{n_1+1}^0 \dots d_{n_1+n_2} d_{n_1+n_2}^0}^{(2n_1+1, \dots, 2n_1+2n_2)} \dots \otimes W_{d_n d_n^0}^{(2n-2n_m+1, \dots, 2n)}; \quad (9.39)$$

then using Jamiołkowski isomorphism (9.35), one can obtain the corresponding positive map.

For instance, considering the tensor product of n REW s (corresponding to the partition $n = 1 + 1 + \dots + 1$)

$$W_{d_1 d_1^0 \dots d_n d_n^0}^{(1, 2, \dots, 2n)} = \bigotimes_{k=1}^n W_{d_k d_k^0}^{(2k-1, 2k)}$$

with

$$W_{d_k d_k^0}^{(2k-1, 2k)} = a_1^{(2k-1, 2k)} I + (a_2^{(2k-1, 2k)} - a_1^{(2k-1, 2k)}) j_{00}^{(2k-1, 2k)} i_{00}^{(2k-1, 2k)} j + a_1^{(0, 2k-1, 2k)} i_{01}^{(0, 2k-1, 2k)}$$

acting on Hilbert space $H_{d_k}^{(2k-1)} \otimes H_{d_k^0}^{(2k)}$ with $i_{01}^{(0, 2k-1, 2k)}$ given in Eq. (4.7), we get the following positive map

$$E^{(n)}(\cdot) = \sum_{j_1, j_2, \dots, j_n} X_{j_1, j_2, \dots, j_n} O_s$$

where

$$O_s = \sum_{j_1, j_2, \dots, j_n} \left(a_1^{(2j_1-1, 2j_1)} I_{d_{j_1}}^{(j_1+1)} + a_1^{(0, 2j_1-1, 2j_1)} \bigotimes_{k=d_{j_1}}^{d_{j_1}^0-1} j_{00}^{(0, 2j_1-1, 2j_1)} \text{Tr}_{j_1, \dots, j_n}(\cdot) \right)$$

and $(j_1 \dots j_{\beta_j})$ is the ordered of S . Choosing all $a_2 = a_1^0 = 0$ and $a_1 = 1$ yields the generalized reduction map introduced in [21]. As an example for $n = 2$ one can easily verify that

$$\begin{aligned}
 E^{(2)}(\rho) &= \text{Tr}_{1,3} [W_{d_1 d_1 d_2 d_2}^{(1;2;3;4)} (\text{Tr}_{1,3} I_{d_1 d_2}^{(2;4)})] = a_1^{(1;2)} a_1^{(3;4)} \text{Tr}(\rho) I_{d_1}^{(2)} I_{d_2}^{(4)} + a_1^{(1;2)} (a_2^{(3;4)} a_1^{(3;4)}) I_{d_1}^{(2)} \text{Tr}_1(\rho) \\
 &\quad + a_1^{(3;4)} (a_2^{(1;2)} a_1^{(1;2)}) \text{Tr}_3(\rho) I_{d_2}^{(4)} + (a_2^{(1;2)} a_1^{(1;2)}) (a_2^{(3;4)} a_1^{(3;4)}) \\
 &\quad a_1^{(1;2)} a_1^{(3;4)} \text{Tr}(\rho) I_{d_1}^{(2)} \sum_{k=d_2}^{d_1^0-1} j_{k|k} j + (a_2^{(1;2)} a_1^{(1;2)}) a_1^{(3;4)} \text{Tr}_3(\rho) \sum_{k=d_2}^{d_1^0-1} j_{k|k} j \\
 &\quad + a_1^{(1;2)} (a_2^{(3;4)} a_1^{(3;4)}) \sum_{k=d_1}^{d_1^0-1} j_{k|k} j \text{Tr}_1(\rho) + a_1^{(1;2)} a_1^{(3;4)} \text{Tr}(\rho) \sum_{k=d_1}^{d_1^0-1} j_{k|k} j I_{d_2}^{(4)} \\
 &\quad + a_1^{(1;2)} a_1^{(3;4)} \text{Tr}(\rho) \sum_{k=d_1}^{d_1^0-1} j_{k|k} j \sum_{k=d_2}^{d_1^0-1} j_{k|k} j
 \end{aligned}$$

Taking $a_2^{(1;2)} = a_2^{(3;4)} = a_1^{(1;2)} = a_1^{(3;4)} = 0$ and $a_1^{(1;2)} = a_1^{(3;4)} = 1$ yields

$$E^{(2)}(\rho) = \text{Tr}(\rho) I_{d_1}^{(2)} I_{d_2}^{(4)} I_{d_1}^{(2)} \text{Tr}_1(\rho) \text{Tr}_3(\rho) I_{d_2}^{(4)} + \quad (9.40)$$

and choosing $a_2^{(1;2)} = a_2^{(3;4)} = 0$ and $a_1^{(1;2)} = a_1^{(3;4)} = a_1^{(1;2)} = a_1^{(3;4)} = 1$ we reach the new reduction positive map for different dimensions

$$\begin{aligned}
 E^{(2)}(\rho) &= \text{Tr}(\rho) I_{d_1}^{(2)} I_{d_2}^{(4)} I_{d_1}^{(2)} \text{Tr}_1(\rho) \text{Tr}_3(\rho) I_{d_2}^{(4)} + \sum_{k=d_2}^{d_1^0-1} \text{Tr}(\rho) I_{d_1}^{(2)} j_{k|k} j \text{Tr}_3(\rho) \sum_{k=d_2}^{d_1^0-1} j_{k|k} j \\
 &\quad + \sum_{k=d_1}^{d_1^0-1} j_{k|k} j \text{Tr}_1(\rho) \text{Tr}(\rho) \sum_{k=d_1}^{d_1^0-1} j_{k|k} j I_{d_2}^{(4)} + \sum_{k=d_1}^{d_1^0-1} \text{Tr}(\rho) j_{k|k} j \sum_{k=d_2}^{d_1^0-1} j_{k|k} j
 \end{aligned}$$

These examples show that one can construct more positive maps by making tensor product of EW s $W_R^{(n)}$ in arbitrary way provided that the dimensionality condition $d_1 \leq d_1^0$ is satisfied.

10 Conclusion

The generalized reduction type entanglement witnesses with simplex feasible regions are introduced, where the EW s corresponding to hyperplanes surrounding the feasible regions are

optimal. These REWs are of types that their manipulation is reduced to LP problem and can be solved exactly by using the simplex method. As it shown above, the REWs are decomposable in cases if their second eigenvalue, namely λ_2 becomes positive while for negative values of λ_2 , the decomposability or non-decomposability of REWs is still open for debate. Also various other interesting issues remain unsolved, such as keeping the REWs in realm of LP problems despite of adding some other operators or entangled states to them.

Appendix I:

Proof of the inequalities: $0 \leq P_S; P_S^0; P_{2, \dots, n} \leq 1$.

In this appendix we prove that all P_S , P_S^0 and $P_{2, \dots, n}$ take the values between 0 and 1. The inequalities $0 \leq P_S; P_S^0 \leq 1$ can be easily concluded from the following ones

$$0 \leq \text{Tr}(\rho_j^{(k)}) = \sum_{i=1}^n \rho_{ii}^{(k)} = 1;$$

$$0 \leq \text{Tr}(\rho_S^{(k)}) = \sum_{i=1}^n \rho_{ii}^{(k)} = 1;$$

For $P_{2, \dots, n}$, the Cauchy-Schwartz inequality implies that

$$P_{2, \dots, n} = d_1 \sum_{i=0}^{n-1} \rho_{ii}^2 = \sum_{i=0}^{n-1} \rho_{ii}^{(1)} \rho_{ii}^{(2)} \dots \rho_{ii}^{(n)} = \sum_{i=0}^{n-1} \rho_{ii}^{(1)} \rho_{ii}^{(2)} \dots \rho_{ii}^{(n)};$$

where

$$\rho_{ii} = \begin{pmatrix} \rho_{00}^{(1)} & \rho_{00}^{(2)} & \dots & \rho_{00}^{(n)} \\ \rho_{10}^{(1)} & \rho_{10}^{(2)} & \dots & \rho_{10}^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \rho_{(n-1)0}^{(1)} & \rho_{(n-1)0}^{(2)} & \dots & \rho_{(n-1)0}^{(n)} \end{pmatrix};$$

and using the following inequality

$$\rho_{ii}^{(1)} \rho_{ii}^{(2)} \dots \rho_{ii}^{(n)} = \sum_{k=2}^n \rho_{ii}^{(k)} = 1;$$

one can conclude that $0 \leq P_{2,\dots,m} \leq 1$.

Appendix II:

Proof of the feasible region of (4.11).

In order to obtain the feasible region of (4.11), we need to evaluate the expectation value of optimal EWs

$$\langle S \rangle W_{\text{opt}}^{(n)} = a_s \langle \mathcal{O}_s \rangle + \sum_{s \in S \subseteq \mathcal{O}_{NS}^0} (1) \mathcal{P}^{\dagger} \mathcal{P}^{0j}_{s0} + d_1 (1) \mathcal{P}^{\dagger} \mathcal{N}^{0j}_{j_{00:\dots:0} \text{ in }_{00:\dots:0}} \langle \mathcal{A}_s \rangle \leq N^0 \quad (\text{I-1})$$

in pure product states $j \text{ in } j$ where $\langle S \rangle W_{\text{opt}}^{(n)} = \langle 1 \rangle W_{\text{opt}}^{(n)}$. Now by taking the partial transpose of $\langle S \rangle W_{\text{opt}}^{(n)}$ with respect to $(N^0 \text{ in } S)$ we have

$$\langle S \rangle W_{\text{opt}}^{(n)T(N^0 \text{ in } S)} = a_s \sum_{i \in j} \langle S \rangle_{ij} \text{ in } \langle S \rangle_{ij} j \quad (\text{I-2})$$

where

$$j \langle S \rangle_{ij} i = j \langle 1 \rangle_{ij} i - j \langle 2 \rangle_{ij} i \dots j \langle n \rangle_{ij} i + j \langle 1 \rangle_{ij} i - j \langle 2 \rangle_{ij} i \dots j \langle n \rangle_{ij} i; \quad (\text{I-3})$$

$$j \langle k \rangle_{ij} i = \begin{cases} jji & \text{if } k \geq 1 \text{ in } S \\ \geq jji & \text{if } k \neq 1 \text{ in } S \end{cases}; \quad j \langle k \rangle_{ij} i = \begin{cases} jji & \text{if } k \geq 1 \text{ in } S \\ \geq jji & \text{if } k \neq 1 \text{ in } S \end{cases} \quad (\text{I-4})$$

Noting that all of these operators are positive definite and using the relation

$$\text{Tr}(\langle S \rangle W_{\text{opt}}^{(n)T(N^0 \text{ in } S)} j \text{ in } j) = \text{Tr}(\langle S \rangle W_{\text{opt}}^{(n)} (j \text{ in } j)^{T(N^0 \text{ in } S)}) \geq 0$$

yields all feasible regions which are simplices.

Appendix III:

Proof of the optimality of $\langle S \rangle W_{\text{opt}}^{(n)}$.

Here in this appendix we try to prove that witness $\langle S \rangle W_{\text{opt}}^{(n)}$ is optimal, to this aim we give the proof for the special case $\langle 2 \rangle W_{\text{opt}}^{(n)}$, since the proof of general case is rather similar to this

particular one. According to the Reference [13], the EW $^{(2)}W_{\text{opt}}^{(n)}$, is optimal if for all positive operator P and $\epsilon > 0$, the following new Hermitian operator

$$W_{\text{new}} = (1 + \epsilon) ^{(2)}W_{\text{opt}}^{(n)} - \epsilon P \quad (\text{I-5})$$

is not anymore an EW. Suppose that there is a positive operator P and $\epsilon > 0$ such that $W_{\text{new}} = ^{(2)}W_{\text{opt}}^{(n)} - \epsilon P$ is yet an EW. Let the positive operator P be the pure projection operator $|j_i\rangle\langle j_i|$, since an arbitrary positive operator can be written as sum of pure states with positive coefficients as $P = \sum_i |j_i\rangle\langle j_i|$.

Now, one should note that the expectation value of the operator $^{(2)}W_{\text{opt}}^{(n)}$ in pure product states $|j_i\rangle$ will vanish if they satisfy the following equation

$$A_i B_j + A_j B_i = 0 \quad (\text{I-6})$$

with

$$A_i = (a_1)_i (a_2)_i \dots (a_n)_i; \quad B_j = (b_1)_j (b_2)_j \dots (b_n)_j$$

But, it is straightforward to see that, for $A_i, B_j \in \mathbb{R}$, the pure state $|j_i\rangle$ will be similar to one of the $|j_{ij}^{(2)}\rangle$ with $i \in j$ with

$$|j_{ij}^{(2)}\rangle = \frac{1}{\sqrt{d_1}} (|j_i\rangle + |j_i\rangle + \dots + |j_i\rangle) \quad ; \quad j = 0; \dots; d_1 - 1; \quad (\text{I-7})$$

concluding that an arbitrary P has the form: $P = \sum_{i \in j}^{d_1 - 1} a_{ij} |j_{ij}^{(2)}\rangle\langle j_{ij}^{(2)}|$ with $a_{ij} \geq 0$.

Finally, substituting Eq. (I-6) in the following expression

$$\text{Tr}(P |j_i\rangle\langle j_i|) = \sum_{ij} a_{ij} (A_i B_j + A_j B_i)^2 = \sum_{ij} a_{ij} \frac{A_j^2}{B_j} (B_j B_i - B_j B_i)^2 = 0$$

and choosing B_i 's such that $B_j B_i \neq B_j B_i$ yields $a_{ij} = 0$ and consequently one can conclude that $P = 0$.

References

- [1] J. Preskill, The Theory of Quantum Information and Quantum Computation (California Institute of Technology, Pasadena, CA, 2000), <http://www.theory.caltech.edu/people/preskill/ph229/>.
- [2] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge University Press, Cambridge, 2000).
- [3] The Physics of Quantum Information: Quantum Cryptography, Quantum Teleportation and Quantum Computation, edited by D. Bouwmeester, A. Ekert, and A. Zeilinger (Springer, New York, 2000).
- [4] M. Horodecki, P. Horodecki and R. Horodecki, Springer Tracts Mod. Phys. 173, 151 (2001).
- [5] M. Lewenstein, D. Bru, J. I. Cirac, B. Kraus, M. Kus, J. Samsonowicz, A. Sanpera, and R. Tarrach, J. Mod. Opt. 47, 2841 (2000).
- [6] B. M. Terhal, Theor. Comput. Sci. 287, 313 (2002).
- [7] D. Bru, J. Math. Phys. 43, 4237 (2002).
- [8] J. S. Bell, Physics (N.Y.) 1, 195 (1964).
- [9] A. Peres, Phys. Rev. Lett. 77, 1413 (1996).
- [10] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A 223, 1 (1996).
- [11] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A 54, 3824 (1996).
- [12] W. K. Wootters, Phys. Rev. Lett. 80, 2245 (1998).

- [13] M .Lewenstein, B .K raus, J.I.C irac, and P .H orodecki, Phys. Rev. A 62, 052310 (2000);
 ibid. 63, 044304 (2001).
- [14] M .A .Jafarizadeh, M .Rezaee, S.K .A .Seyed Yagoobi, Phys. Rev. A 72, 062106 (2005).
- [15] M .A .Jafarizadeh, M .Rezaee, S.A hadpour, Phys. Rev. A 74, 042335 (2006).
- [16] E .Storm er, Acta M ath. 110, 233 (1963); S.L.W oronow icz, Rep. M ath. Phys. 10, 165
 (1976); M .D .Choi, Proc. Sym pos. Pure M ath. 38, 583 (1982).
- [17] A .C .D oherty, P .A .P arrilo, and F .M .Spedalieri, Phys. Rev. Lett. 88, 187904 (2002).
- [18] A .C .D oherty, P .A .P arrilo, and F .M .Spedalieri, Phys. Rev. A . 69, 022308 (2004).
- [19] R .O .V ianna, A .C .D oherty, eprint quant-ph/0608095 (2006).
- [20] S.Boyd and L.Vandenberghe, Convex Optim ization, C am bridge U niversity P ress, (2004).
- [21] W .Hall, Phys. Rev. A 72, 022311 (2005)
- [22] A .A c n, D .B ru , M .Lewenstein, and A .Sanpera, Phys. Rev. Lett. 87, 040401 (2001).
- [23] A .Jam iolkow ski, Rep. M ath. Phys. 3, 275 (1972).

Figure Captions

Figure-1: 3-simplex displaying the feasible region of three-qubit REW .

Figure-2: 3-simplex displaying the feasible region of multi-qubit BDEW W_1 .

Figure-3: Convex polygon displaying the boundaries of the feasible region for multi-qubit BDEW W_2 .

Figure-4: Decomposable and non-decomposable regions of REW s: for $!_2 \geq 0$ the REW s are decomposable, for $!_2 < 0$ the REW s are non-decomposable and in dashed region, the decomposability or non-decomposability of REW s is still open for debate.




Fig. (1)




Fig. (2)




Fig. (3)




Fig. 4