

Constructing Operator Valued Probability Measures in Phase Space

Demosthenes Ellinas

Department of Sciences, Division of Mathematics,
Technical University of Crete, GR 731 00 Chania Crete Greece

February 12, 2021

Abstract

Probability measures (quasiprobability mass), given in the form of integrals of Wigner function over areas of the underlying phase space, give rise to operator valued probability measures (OVM). General construction methods of OVMs, are investigated in terms of geometric positive trace increasing maps (PTI), for general 1D domains, as well as 2D shapes e.g. circles, disks. Spectral properties of OVMs and operational implications of their constructing PTIs are discussed.

Keywords: Quantum probability, POVM, Wigner function, Phase space Quantum Mechanics

PACS: 02.50.-r, 03.65.Ta, 03.67.-a

An OVM is a map $K : \mathcal{F} \rightarrow L(H)$; from a σ -algebra \mathcal{F} of subsets of a nonempty set X ; to the bounded operators $L(H)$ on a Hilbert space H ; such that for $A \in \mathcal{F}$; and $X \in \mathcal{F}$, the function $\mu(A) = \text{Tr}(K(A))$ is a normalized (i.e. $\mu(X) = 1$), generalized (i.e. negative valued) measure (i.e. additive set function). If $\mu(A) \geq 0$; i.e. $\text{Tr}(K(A)) \geq 0$; $\forall A \in \mathcal{F}$ we have a positive OVM, further if $K(A)^* = K(A)$; and $K(A)^2 = K(A)$; we have a projective OVM, namely an observable (see e.g. [1]). For the case of Wigner function [2][3] $W_{j>}(x) = \int_{\mathbb{R}^2} D(y) D(y-x) dy$; we have $\mathcal{C}; H = \text{span}\{|n\rangle; n = 0, 1, \dots\}$ the Fock space, with N the number operator, $\pi = e^{iN}$ the parity operator, and $D(x) = e^{ax} R^a = e^{(iqP - ipQ)}$ the displacement operator. Integrals of Wigner function [4][5] $\int_X W_{j>}(x) d^2x = \text{Tr}(j \text{ in } K(X))$; physically define the quasi probability mass over X ; in terms of OVM $K(X) = \int_X D(x) D(x)^* d^2x$:

Proposition 1 Let X_q a region in q -axis determined by the characteristic function (cfun), $\chi_q(x); x \in \mathbb{R}$; with Fourier transform (FT) $e(p)$; then the associated OVM is $K_{X_q} = e(P)$: The solution of eigen problem $K_{X_q} j_i = \lambda_j j_i$; determines the eigenvalues $\lambda_j = e(p)$; and the eigenvectors $j_i = |p_i\rangle$

ellinas@science.tuc.gr

pi: If further $\psi(q)$ is L periodic with Fourier series $\psi(q) = \frac{a_0}{2} + \sum_{m=1}^P a_m \cos(\frac{m}{L}q) + b_m \sin(\frac{m}{L}q)$; the OVM becomes

$$K_q = \frac{a_0}{2} \delta(q) + \sum_{m=1}^P r_m e^{i m q} \delta(q) = \frac{m}{L} i \hbar p = \frac{m}{L} j + e^{-i m q} \delta(q) = \frac{m}{L} i \hbar p = \frac{m}{L} j ; \quad (1)$$

where $r_m = \frac{1}{2}(a_m + i b_m)$, and $\theta_m = \arg(a_m + i b_m)$:

Proof: The region operator (convention: alternative name for OVM related to quasi-probability functions, see below), is constructed by smearing along q axis the point at the origin as follows

$$K_q = \int_{-R}^R e^{i q p} e^{-i q p} dq = \int_{-R}^R \psi(q) e^{i q p} dq = \int_{-R}^R \psi(p) \delta(p - q) e^{i q p} dq ; \quad (2)$$

where $\psi(p) = \int_{-R}^R \psi(q) e^{i q p} dq$; is the FT. Straightforward evaluation of FT of the cfun of eq. (3), below leads to the result. The eigen problem is also easily solved utilizing the action of parity operator on momentum state.

Remarks: 1) From FT of a L periodic cfun of some region we obtain the eigenvalues of its associated OVM $K_q = \psi(p)$; where

$$\psi(w) = a_0 \delta(w) + \sum_{m=1}^P a_m \cos\left(w \frac{m}{L}\right) + i b_m \sin\left(w \frac{m}{L}\right) ; \quad (3)$$

If cfun is asymmetric then its FT is complex valued, this implies the region operator would not be a real one.

2) Let e.g. the pure density matrix $\rho = |j\rangle\langle j|$ determined by an even parity state vector i.e. $\langle j|i\rangle = \langle j|i\rangle$; then the occupation probability of the region X_q is

$$p(X_q) = \text{Tr}(K_q \rho) = \int_{-R}^R \psi(p) |j(p)|^2 dp ; \quad (4)$$

If further the region X_q is symmetric with respect to the origin, i.e. its cfun is an even function (i.e. $b_m = 0$); then we obtain the probability

$$p(X_q) = a_0 |j(0)|^2 + 2 \sum_{m=1}^P a_m |j(\frac{m}{L})|^2 ; \quad (5)$$

3) Let us choose the special region for which $a_m = 0$; $b_m = 0$; then the associated region operator becomes

$$K_q = a_0 \delta(q) + \sum_{m=1}^P a_m \cos\left(\frac{m}{L} q\right) \delta(q) = \frac{m}{L} i \hbar p = \frac{m}{L} j + \sum_{m=1}^P a_m \cos\left(\frac{m}{L} q\right) \delta(q) = \frac{m}{L} i \hbar p = \frac{m}{L} j ; \quad (6)$$

This is also expressed as $K_q = \int_{-\infty}^{\infty} \delta(p - 0) \delta(p = 0) \dots$; with μ_q been a PTIM ap is generated by the set of Kraus generators [6] $\mu_q = \int_{-\infty}^{\infty} \delta(p - 0) \delta(p = 0) \dots$. The latter expression of K_q ; if utilized together with state-observable duality (c.f. [7]), would provide, by means of a unitarization of μ_q map; means for some physical implementation of the general region operator K_q (see similar constructions of other region operators in [8], and below).

4) The operator K_q defined on a region of q axis with $\mu_q(p)$, can be rotated by $\frac{1}{2}$ radians to become K_p ; i.e. a similar operator on p -axis which reads

$$K_p = e^{i\frac{1}{2}N} K_q e^{-i\frac{1}{2}N} = e^{i\frac{1}{2}N} e(P) e^{-i\frac{1}{2}N} = e(Q) : \quad (7)$$

Having available operators K_q ; K_p ; been defined along orthogonal axes, we further rotate them clockwise by an angle θ ; to obtain the respective operators along new rotated axes as follows,

$$K_q = e^{i\theta N} K_q e^{-i\theta N} = e^{i\theta N} e(P) e^{-i\theta N} = e(\cos \theta P - \sin \theta Q) ; \quad (8)$$

and

$$K_p = e^{i\theta N} K_p e^{-i\theta N} = e^{i\theta N} e(Q) e^{-i\theta N} = e(\cos \theta Q + \sin \theta P) : \quad (9)$$

These two operators are related by a $\frac{1}{2}$ radians rotation i.e. $K_p = e^{i\frac{1}{2}N} K_q e^{-i\frac{1}{2}N}$:

5) Using properties of Fourier transform we obtain shift transforms of a region operator e.g. $K_q e^{icP} = e(P + c)$ as follows,

$$e^{icP} K_q = e(P + c) ; K_q e^{icP} = e(P - c) ; e^{i\frac{c}{2}P} K_q e^{-i\frac{c}{2}P} = e(P + c) : \quad (10)$$

6) Using the squeezing operator $S(a) = e^{a^2 \alpha^2 - \frac{1}{2} \alpha^2} = e^{\frac{1}{2} r (Q^2 - P^2)}$, with parameter $r = \frac{1}{2} \ln \frac{1}{s}$; we obtain by similarity transformation multiplicative actions on position and momentum operators

$$S(a) Q S(a) = Q e^r ; S(a) P S(a) = P e^{-r} : \quad (11)$$

This gives rise to a region operator with (squeezed) scaled support on q -axis i.e.

$$S(a) K_q S(a) = S(a) e(P) S(a) = e(P e^{-r}) : \quad (12)$$

7) OVM from s-Parametrized Quasiprobability Functions: Working out the power series expression of s-parametrized quasiprobabilities phase space functions $F(\alpha, \beta; s)$; of [9, 3]

$$F(\alpha, \beta; s) = \sum_{k=0}^{\infty} \frac{2^k}{(1-s)^{k+1}} (1)^k \text{Tr}(\rho^k) D(\alpha)^k D(\beta)^k ; \quad (13)$$

we obtain that

$$F(\alpha, \beta; s) = \frac{2}{\pi} \text{Tr}(\rho(\alpha, \beta; s)) = \frac{2}{\pi} \text{Tr}(D(\alpha) (s) D(\beta)^Y) ; \quad (14)$$

where the s -parametrized parity operator $\rho(s) = \frac{(1+s)^N}{(1-s)^{N+1}}$; is now displaced by the coherent state generating operators. Special cases are the Glauber-Sudarshan P function $F(\cdot; s=1) = P(\cdot)$; the Wigner function $F(\cdot; s=0) = W(\cdot)$; and the Q positive function $F(\cdot; s=-1) = Q(\cdot)$: The "point operator" at the origin of phase plane is now the s -parametrized parity operator $\rho(s)$: Applying appropriate geometric PTIMaps $\rho(s)$ as in the present case of Wigner function, i.e. $s=0$; we can derive s -parametrized OVMs.

Examples: We proceed with examples that elucidate the general construction: 1) Let us choose $X_q = \{1, 2, \dots, n\}$; namely the case where the region consists of the first n positive integers. This set with $\rho(x) = 1$ for $x = 1, 2, \dots, n$; and 0 otherwise, leads to region operator

$$K_q = \int_R (x) e^{i2xP} dx = \sum_{m=1}^n e^{i2mP} = \frac{e^{i(2n+1)P} - e^{iP}}{2i \sin P} : \quad (15)$$

As the set is not symmetric with respect to the origin of axes, the resulting region operator is a complex one in this case.

2) Circle OVM $K_C(a)$; construction steps: displace by a along q -axis the 0 point OVM i.e. $\int_0^{2\pi} e^{-iaP} e^{i\theta P} d\theta$; and rotate it around by means of the continuous PTIMap ρ_2 ; with Kraus generators $f e^{iN}$; $0 < 2g$; i.e. $e^{-iaP} e^{i\theta P} \rho_2(e^{-iaP} e^{i\theta P})$: Explicitly,

$$\begin{aligned} & \int_0^{2\pi} e^{-iaP} e^{i\theta P} \rho_2(e^{-iaP} e^{i\theta P}) d\theta = \int_0^{2\pi} e^{-iaP} e^{i\theta P} e^{-iN\theta} e^{i\theta P} e^{iN\theta} d\theta \\ & = \int_0^{2\pi} d e^{iN\theta} e^{2iaP} e^{iN\theta} = \sum_{m,n=0}^{\infty} d e^{iN\theta} j_n i l m j e^{2iaP} j_n i l m j e^{iN\theta} \end{aligned} \quad (16)$$

$$= 2 \sum_{n=0}^{\infty} e^{-a^2=2} L_n(a^2) j_n i l m j K_C(a); \quad (17)$$

where $j_n i l m j = e^{-a^2=2} L_n(a^2)$; L_n been the Laguerre polynomials (result obtained originally in [4]).

3) Disc OVM $K_D(a)$; construction steps: start with OVM $K_L(a)$ of line segment extended along $[-L; L]$; been constructed by means of PTIMap in [8], now rotate it around by means of the continuous PTIMap ρ_2 ; with Kraus

generators $f e^{-i N}; 0 < 2 g$: Explicitly,

$$K_L(a) = \frac{\sin(P a)}{P} \int_{-\infty}^{\infty} dx e^{i N x} K_L(a) e^{i N x} \\ = \int_{-\infty}^{\infty} dx e^{i N x} \frac{\sin(P a)}{P} e^{i N x} = \int_{-\infty}^{\infty} dx e^{i N x} e^{2ixP} e^{i N x} \quad (18)$$

$$= \int_{-\infty}^{\infty} dx e^{i N x} e^{2ixP} e^{i N x} = \int_{-\infty}^{\infty} dx e^{i N x} e^{2ixP} e^{i N x} \quad (19)$$

Operational construction of OVMs: Let $\rho^1 = D(x) D^\dagger(x) \rho^1$; $\rho^2 = 1 D(x) D^\dagger(x)$; be point operators with support on points $x \in \mathbb{C}$; belonging to two respective phase planes. Their sum acting in $H = H^1 \oplus H^2$; is $\rho^1 + \rho^2 = D(x) D^\dagger(x) (\rho^1 + \rho^2) D^\dagger(x) D(x)$; i.e. is the displaced sum of two operators with support on the zero point of the respective phase planes. To construct the latter we resort to the property: the unitary operator $V = \exp(i \frac{1}{2} J)$; where $J = \frac{1}{2i} (a_1^\dagger a_2 - a_2^\dagger a_1)$; serves as the permutation operators between parity operators of two different Hilbert spaces i.e. $\rho^2 = V^2 \rho^1 V^{-2}$. Then $\rho^1 + \rho^2 = \rho^1 + V^2 \rho^1 V^{-2} = \rho^1$; where we have introduced the positive trace increasing map $\rho \rightarrow \rho$; with two Kraus operators $(1; V^2)$. Let the density matrix acting in $H = H^1 \oplus H^2$; and consider the expectation value $\text{Tr}(\rho^1 + \rho^2) = \text{Tr}(\rho^1) = \text{Tr}(\rho^1)$; where the dual PT map $\rho \rightarrow \rho$ has been introduced (see [7]), as $\rho^1 = \rho^1 + V^2 \rho^1 V^{-2}$; obtaining the value $\text{Tr}(\rho^1 + \rho^2)$; requires the operational construction of density matrix ρ^1 ; which amounts to a unitary dilation of ρ^1 : To accomplish this we introduce the auxiliary space $H_A = \text{span}\{|0\rangle, |1\rangle\}$; and the unitary operator

$$W = \begin{pmatrix} 1 & V^2 \\ V^{-2} & 1 \end{pmatrix}; \quad (20)$$

acting on $H_A \oplus H = H^1 \oplus H^2$: Then we obtain $\rho^1 = \text{Tr}_A(W^\dagger (\rho^1 \otimes \rho^1) W)$; the partial tracing map Tr_A signifies operationally an unconditional measurement in H_A [10].

Acknowledgment: Work partially supported by EPEAEK:Pythagoras II.

References

- [1] P. Bush, M. Grabowski and P. Lahti, Operational Quantum Mechanics, (Springer-Verlag, Berlin 1992, 2nd corrected printing)
- [2] W. P. Schleich, Quantum Optics in Phase Space, (Wiley-VCH, Berlin 2001).
- [3] A. Royer, Phys. Rev. A, 15, 449{450 (1977).

- [4] A. J. Bracken, H.-D. Döbner and J. G. Woods, *Phys. Rev. Lett.* **83**, 3758 (1999).
- [5] A. J. Bracken, D. Ellinas and J. G. Woods, *Acta Phys. Hung. B, Quant. Electronics*, **20**, 121 (2004); *J. Phys. A Math. and Gen.* **36**, L297 (2003).
- [6] K. Kraus *States, Effects and Operations*, (Springer, Berlin 1983).
- [7] By means of the cyclic property of trace, the expectation value of an observable X ; viz. $\langle X \rangle_i = \text{Tr}(X \rho_i)$; on a state ρ_i mapped by a CP map \mathcal{E} ; as $\rho_i = \sum_j A_{ij} \rho_j A_{ij}^\dagger$; serves to define the dual map $\mathcal{E}^\dagger(X) = \sum_j A_{ij}^\dagger X A_{ij}$; that satisfy the equation $\langle X \rangle_i = \text{Tr}(\mathcal{E}^\dagger(X) \rho_i)$:
- [8] D. Ellinas and I. Tsochantzjis, *Rep. Math. Phys.* **57**, 69 (2006).
- [9] H. Moya-Cessa and P. L. Knight, *Phys. Rev. A* **48**, 2479 (1993).
- [10] W. M. de Muynck, *Foundations of Quantum Mechanics, an Empiricist Approach*, (Kluwer, Dordrecht, 2002).