

ADDENDUM TO “MULTIPARTITE STATES UNDER LOCAL UNITARY TRANSFORMATIONS”

Sergio Albeverio*

Institut für Angewandte Mathematik, Universität Bonn, Wegelerstr. 6, D-53115 Bonn
e-mail: albeverio@uni-bonn.de

Laura Cattaneo

Institut für Angewandte Mathematik, Universität Bonn, Wegelerstr. 6, D-53115 Bonn
e-mail: cattaneo@wiener.iam.uni-bonn.de

Shao-Ming Fei

Institut für Angewandte Mathematik, Universität Bonn, Wegelerstr. 6, D-53115 Bonn
Department of Mathematics, Capital Normal University, Beijing 100037
Max Planck Institute for Mathematics in the Sciences, Inselstr. 22, D-04103 Leipzig
e-mail: fei@uni-bonn.de

Xiao-Hong Wang

Department of Mathematics, Capital Normal University, Beijing 100037
e-mail: wangxh@mail.cnu.edu.cn

(Received 2005)

In previous work the authors introduced a notion of generic states and obtained criteria for local equivalence of them. Here they introduce the concept of CHG states maintaining the criteria of local equivalence. This fact allows the authors to halve the number of invariants necessary to characterize the equivalence classes under local unitary transformations for the set of tripartite states whose partial trace with respect to one of the subsystems belongs to the class of CHG mixed states.

Keywords: tripartite quantum states, local unitary transformations, en-

SFB 611; IZKS; BiBoS; CERFIM (Locarno); Acc. Arch. USI (Mendrisio)

tanglement, invariants

In the paper [1] we exploited the equivalence criterion for the members of a class of bipartite mixed states constructed in [2, 3] to write an equivalence criterion for the members of a class of pure tripartite states. In this Addendum we show how the conditions assumed in [2, 3, 1] can be relaxed.

We first consider (mixed) states on a bipartite system $H_A \otimes H_B$, where H_A and H_B are finite dimensional Hilbert spaces of dimension N_A and N_B , respectively. Let ρ be a density matrix defined on $H_A \otimes H_B$ with $\text{rank}(\rho) = n \leq N^2$, where $N = \min\{N_A, N_B\}$. ρ can be decomposed according to its eigenvalues and eigenvectors:

$$\rho = \sum_{i=1}^n \lambda_i |\varphi_i\rangle \langle \varphi_i|,$$

where λ_i resp. $|\varphi_i\rangle$, $i = 1, \dots, n$, are the nonzero eigenvalues resp. eigenvectors of the density matrix ρ . $|\varphi_i\rangle$ has the form

$$|\varphi_i\rangle = \sum_{k=1}^{N_A} \sum_{l=1}^{N_B} a_{kl}^i |e_k\rangle \otimes |f_l\rangle, \quad a_{kl}^i \in \mathbb{C}, \quad \sum_{k=1}^{N_A} \sum_{l=1}^{N_B} a_{kl}^i a_{kl}^{i*} = 1, \quad i = 1, \dots, n,$$

where $\{|e_i\rangle\}_{i=1}^{N_A}$ and $\{|f_i\rangle\}_{i=1}^{N_B}$ are orthonormal bases in $\mathcal{H}_A$ and $\mathcal{H}_B$, respectively, and $*$ means complex conjugation. Let A_i denote the matrix given by $(A_i)_{kl} = a_{kl}^i$. We introduce $\{\rho_i\}$, $\{\theta_i\}$:

$$\rho_i = \text{Tr}_B |\varphi_i\rangle \langle \varphi_i| = A_i A_i^\dagger, \quad \theta_i = (\text{Tr}_A |\varphi_i\rangle \langle \varphi_i|)^* = A_i^\dagger A_i, \quad i = 1, \dots, n, \quad (1)$$

with † denoting adjoint. Tr_A and Tr_B stand for the traces over the first and second Hilbert spaces respectively, and therefore, ρ_i and θ_i can be regarded as reduced density matrices. Let $\Omega(\rho)$ and $\Theta(\rho)$ be two “metric tensor” matrices, with entries given by

$$\Omega(\rho)_{ij} = \text{Tr}(\rho_i \rho_j), \quad \Theta(\rho)_{ij} = \text{Tr}(\theta_i \theta_j), \quad \text{for } i, j = 1, \dots, n, \quad (2)$$

and

$$\Omega(\rho)_{ij} = \Theta(\rho)_{ij} = 0, \quad \text{for } N^2 \geq i, j > n.$$

In [3] a mixed state ρ is called *generic* if the corresponding “metric tensor” matrices Ω and Θ satisfy

$$\det(\Omega(\rho)) \neq 0 \quad \text{and} \quad \det(\Theta(\rho)) \neq 0. \quad (3)$$

We shall say here that a mixed state ρ is *high generic* if

$$\det(\Omega(\rho)) \neq 0 \quad \text{or} \quad \det(\Theta(\rho)) \neq 0. \quad (4)$$

If we add conditions:

$$[\rho_i, \rho_j] = 0, \quad [\theta_i, \theta_j] = 0 \quad (5)$$

and ρ_i is a full rank matrix. We call a mixed state is a *comm-high generic* or *CHG* state if it is a high generic one and also satisfy the above two conditions. Condition (5) assures that $A_i A_i^\dagger$ and $A_i^\dagger A_i$ have common eigenvectors.

Similarly we also introduce trilinear expressions $X(\rho)$ and $Y(\rho)$ as

$$X(\rho)_{ijk} = \text{Tr}(\rho_i \rho_j \rho_k), \quad Y(\rho)_{ijk} = \text{Tr}(\theta_i \theta_j \theta_k), \quad i, j, k = 1, \dots, n. \quad (6)$$

PROPOSITION 1. *Two CHG density matrices with non-degenerate Ω (or non-degenerate Θ) are equivalent under local unitary transformations if and only if there exists an ordering of the corresponding eigenstates such that the following invariants have the same values for both density matrices:*

$$J^s(\rho) = \text{Tr}_B(\text{Tr}_A \rho^s), \quad s = 1, \dots, n \\ \Omega(\rho), X(\rho) \quad (\text{or } \Theta(\rho), Y(\rho)) \quad (7)$$

Proof: In [2] it was proved that two generic states such that Ω and Θ are both non-degenerate are equivalent under local unitary transformations if and only if there exists an ordering of the corresponding eigenstates such that the invariants $J^s(\rho)$, $s = 1, \dots, n$, $\Omega(\rho)$, $\Theta(\rho)$, $X(\rho)$, and $Y(\rho)$ take the same values for both density matrices. In particular (see [2, eq.(14)] or [3, eq.(15)]), from the conditions $\Omega(\rho)$ non-degenerate, $\Omega(\rho) = \Omega(\rho')$, and $X(\rho) = X(\rho')$ follows that

$$\rho'_i = u \rho_i u^\dagger, \quad (8)$$

for some $u \in \mathcal{U}_A$, where $\mathcal{U}_A$ denotes the space of all unitary matrices on $\mathcal{H}_A$. So we have $A'_i A_i'^\dagger = u A_i A_i^\dagger u^\dagger$. Thus A_i and A'_i have the same singular

values. The *Singular value decomposition* of matrices (see, e.g., [4]) and (5) assure the existence of unitary matrices U, V, U', V' such that

$$UA_iV = \text{diag}(s_1(A_i), \dots, s_n(A_i)), \quad U'A'_iV' = \text{diag}(s_1(A'_i), \dots, s_n(A'_i)),$$

where $s_j(A_i)$ and $s_j(A'_i)$ represent the j -th singular value of A_i and A'_i , respectively, and diag means the principal diagonal (of a non necessarily diagonal matrix). Because of $s_j(A'_i) = s_j(A_i)$ for all i , we have $A'_i = u_1 A_i w_1$, and $|\varphi'_i\rangle = u_1 \otimes w_1 |\varphi_i\rangle$, $i = 1, \dots, n$, where $u_1 = U'^\dagger U$, $w_1 = V'^* V^T$.

From the genericity condition $\Theta(\rho)$ non-degenerate, and from $\Theta(\rho) = \Theta(\rho')$ and $Y_{ijk}(\rho) = Y_{ijk}(\rho')$ we can similarly deduce that

$$\theta'_i = w^\dagger \theta_i w \quad (9)$$

for some $w \in \mathcal{U}_B$. Again from the *Singular value decomposition* of matrices, we have $|\varphi'_i\rangle = u \otimes w |\varphi_i\rangle$, $i = 1, \dots, N^2$, and $\rho' = u \otimes w \rho u^\dagger \otimes w^\dagger$. Hence ρ' and ρ are equivalent under local unitary transformations: the common u and v for different $|\varphi_i\rangle$'s can be obtained in two ways, either from the condition $\Omega(\rho)$ non-degenerate, $\Omega(\rho) = \Omega(\rho')$ and $X_{ijk}(\rho) = X_{ijk}(\rho')$, or from $\Theta(\rho)$ non-degenerate, $\Theta(\rho) = \Theta(\rho')$ and $Y_{ijk}(\rho) = Y_{ijk}(\rho')$. Therefore, it is sufficient to consider only $\Omega(\rho)$ and $X(\rho)$ (or $\Theta(\rho)$ and $Y(\rho)$). ■

Consider now tripartite pure states of a system $H_A \otimes H_B \otimes H_C$, where H_i , $i \in \{A, B, C\}$, is a finite dimensional Hilbert space of dimension N_i . By using the results of Proposition 1, Proposition 2 in [1] can be rewritten as follows.

PROPOSITION 2. *Let $|\psi\rangle$ and $|\psi'\rangle$ be two pure states of $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ and assume that $\rho = \text{Tr}_A(|\psi\rangle\langle\psi|)$ is a CHG mixed state. $|\psi\rangle$ is equivalent to $|\psi'\rangle$ under local unitary transformations if and only if*

$$I_{\alpha,\beta}^{A,s}(|\psi\rangle) = I_{\alpha,\beta}^{A,s}(|\psi'\rangle) \quad (10)$$

for $s \in \{B, C\}$, $\alpha = 1, \dots, \min\{N_B^2, N_C^2\}$, $\beta = 1, \dots, N_r$, where $r \in \{B, C\}$ but is different from s , and for $\rho' = \text{Tr}_A(|\psi'\rangle\langle\psi'|)$

$$\begin{aligned} \Omega(\rho)_{jk} &= \Omega(\rho')_{jk}, & X(\rho)_{jkl} &= X(\rho')_{jkl}, \\ (\text{or } \Theta(\rho)_{jk} &= \Theta(\rho')_{jk}, & Y(\rho)_{jkl} &= Y(\rho')_{jkl}), \end{aligned}$$

for the j, k such that $\lambda_j = \lambda_k$.

The weaker conditions allow us to halve the number of invariants to consider, as compared to the statements of [1].

Acknowledgments

L. Cattaneo gratefully acknowledges the financial support by the Lise Meitner Fellowship of the Land Nordrhein-Westfalen. X.-H. Wang gratefully acknowledges the support provided by the China-Germany Cooperation Project 446 CHV 113/231, “Quantum information and related mathematical problems”.

REFERENCES

- [1] S. Albeverio, L. Cattaneo, S.M. Fei, and X.H. Wang: Multipartite states under local unitary transformations, *Rep. Math. Phys.* **56** (2005), 341.
- [2] S. Albeverio, S.M. Fei, and X.H. Wang: Equivalence of bipartite quantum mixed states under local unitary transformations, in *Proc. First Sino-German Meeting on Stochastic Analysis - Satellite Conference to the ICM 2002*, Edts. S. Albeverio, Z.M. Ma, and M. Röckner, Beijing 2002.
- [3] S. Albeverio, S.M. Fei, P. Parashar, and W.-L. Yang: Nonlocal properties and local invariants for bipartite systems, *Phys. Rev. A* **68** (2003), 010313.
- [4] G.H. Golub and C.F. Van Loan: *Matrix Computations*, 3rd ed., Johns Hopkins University Press, Baltimore 1996.