Exact Solutions of Schrodinger Equation for a Ring-Shaped Potential

O zlem Yesiltas¹, Ram azan Sever²

¹GaziUniversity, Faculty of Arts and Sciences, Department of Physics, 06500, Ankara Turkiye

²M iddle East Technical University, Department of Physics, 06531 Ankara, Turkiye

M arch 29, 2024

Abstract

The exact solutions of Schrodinger equation are obtained for a noncentral potential which is a ring-shaped potential. The energy eigenvalues and corresponding eigenfunctions are obtained for any angular momentum `. Nikiforov-U varov method is used in the computations.

PACS numbers: 03.65 Db, 03.65 Ge

Keywords:Nikiforov-Uvarov, Coulomb, ring shaped

Corresponding Author: sever@metu.edu.tr

1 Introduction

E xact solutions of the fundam ental wave equations of the non-relativistic quantum mechanics have attracted much attention in recent years. So far many e orts have been made to solve the time-independent Schrödinger equation for non-central potentials in three and two dimensions such as the C oulombic ring-shaped potential [1-5] and deformed ring-shaped potential [6], the H ellmann potential [7]. In quantum mechanics, A haronov-Bohm (AB) e ect is an another popular non-central potential type [8]. It is also studied through the wide areas such as path integral solutions[9], in non-commutative quantum mechanics [10], a D irac particle in the A haronov-Bohm potential [11-13] and scattering [14]. On the other hand, AB potential is also known as ring shape potential [6].

Recently, an alternative and simple method called as Nikiforov-Uvarov Method [15] (NUmethod) has been introduced for solving the Schrodinger equation for some well known potentials [16-18]. It is also used to solve the Klein-Gordon and Dun-Kemmer-Petiau wave equations in the presence of exponential type potentials such as standard W oods-Saxon [19], Poschl-Teller [20] and Hulthen potentials [21-23]. The Nikiforov-Uvarov method [15] is used in the computations to obtain energy eigenvalues and the corresponding eigenfunctions. Recently A lhaidari et al have studied D irac and Klein G ordon equation for the C oulom b, oscillator and H artm ann potentials [24].

R ing shaped potential has an application eld in quantum chem istry as a model for the Benzene molecule given as [1,4,9]

$$V_{q}(\mathbf{r}) = {}^{2}_{0} \frac{2a_{0}}{\mathbf{r}} \frac{a_{0}^{2}}{\mathbf{r}^{2}\sin^{2}}$$
(1)

where $a_0 = \frac{-2}{e^2}$; $_0 = \frac{1}{2} \frac{e^4}{-2}$, Bohr radius and the ground state energy of the hydrogen atom, respectively, is the particle mass, and are dimensionless positive real parameters which range from about 1 to 10 [9].

The paper is arranged as follows. In section II, the N ikiforov-U varov M ethod is given brie y. In section III, NU m ethod is applied to three dimensional Schrödinger equation with the ring shaped potential. Results are discussed in section IV.

2 The Nikiforov-U varov M ethod

The NU-m ethod developed by N ikiforov and U varov is based on reducing the second order differential equations (ODEs) to a generalized equation of hypergeom etric type [15]. It is provided us an analytic solution of Schrodinger equation for certain kind of potentials. This is based on the solutions of general second order linear di erential equation with orthogonal functions [15]. For a given appropriate potential, the one-dimensional Schrodinger equation is reduced to a generalized equation of hypergeom etric type with a suitable coordinate transform ation s = s(r). Then, the equation has the form,

$$\infty^{(3)}(s) + \frac{\sim (s)}{(s)}^{(3)}(s) + \frac{\sim (s)}{2(s)}(s) = 0$$
 (2)

where A (s) = $\frac{\sim (s)}{(s)}$ and B (s) = $\frac{\sim (s)}{2(s)}$. In Eq.(2), (s) and \sim (s) are polynomials at most second degree. \sim (s) is a polynomial with at most rst degree [15]. The wave function is constructed as a multiple of two independent parts,

$$(s) = (s)y(s);$$
 (3)

and Eq.(2) becomes [15]

$$(s)y^{\omega}(s) + (s)y^{\circ}(s) + y(s) = 0;$$
 (4)

where

$$(s) = (s) \frac{d}{ds} (\ln (s));$$
 (5)

and

$$(s) = \sim (s) + 2 \quad (s)$$
: (6)

is de ned as

$$_{n} + n^{\circ} + \frac{[n(n-1)^{\circ}]}{2} = 0; n = 0; 1; 2; ...$$
 (7)

determine (s) and by de ning

$$\mathbf{k} = \mathbf{k}^{\circ}(\mathbf{s}) :$$
 (8)

and the function (s) becomes

(s) =
$$(\frac{2}{2})$$
 $(\frac{2}{2})^2 \sim + k$ (9)

The polynom ial (s) with the parameter s and prime factors show the dierentials at rst degree. Since (s) has to be a polynom ial of degree at most one, in Eq.(9) the expression under the square root must be the square of a polynom ial of rst degree [15]. This is possible only if its discriminant is zero. A fter de ning k, one can obtain (s), (s), (s) and . If we look at Eq.(5) and the Rodrigues relation

$$y_n(s) = \frac{B_n}{(s)} \frac{d^n}{ds^n} [(s) (s)];$$
 (10)

where C_n is normalization constant and the weight function satisfy the relation as

$$\frac{d}{ds}[(s)(s)] = (s)(s):$$
 (11)

where

$$\frac{(s)}{(s)} = \frac{(s)}{(s)};$$
(12)

3 Solutions of The Ring-Shaped Potential

Let us start with the Schrodinger equation in three dimensions written as

$$\frac{r^2}{2} - \frac{1}{r^2} \frac{\theta}{\theta r} + \frac{r^2}{\theta r} - \frac{1}{r^2} - \frac{1}{\sin \theta} \frac{\theta}{\theta} - \sin \theta - \frac{\theta}{\theta} + \frac{1}{\sin^2 \theta} \frac{\theta^2}{\theta'^2} + V(r; r) = 0$$
(13)

where V (r;) is taken as below

$$V(r;) = \frac{A}{r} + \frac{B}{r^2 \sin^2}$$
 (14)

A and B parameters are used instead of the parameters dened in Eq.(1) for simplicity. If we take the spherical wave function as (r; ;') = R(r)Y(;') in order to separate the Schrodinger equation into variables, we obtain

$$\frac{1}{R(\mathbf{r})}\frac{\theta}{\theta \mathbf{r}} r^{2}\frac{\theta R(\mathbf{r})}{\theta \mathbf{r}} + \frac{2 A r}{\sim^{2}} + \frac{2 E r^{2}}{\sim^{2}} + \frac{1}{Y(\mathbf{j'})}\frac{1}{\sin \theta} \frac{\theta}{\theta} \sin \frac{\theta Y(\mathbf{j'})}{\theta}$$
(15)

$$\frac{2 B}{\sim^2 \sin^2} + \frac{1}{Y(j')} \frac{1}{\sin^2} \frac{\theta^2 Y(j')}{\theta'^2} = 0$$
(16)

Using Y (;') = H () (') to separate the Schrödinger equation into variables, the following three equations are obtained as:

$$\frac{d^{2}R(r)}{dr^{2}} + \frac{2}{r}\frac{dR(r)}{dr} + \frac{2}{\sim^{2}} E + \frac{A}{r} \frac{\sim^{2}}{2r^{2}} R(r) = 0; \qquad (17)$$

$$\frac{d^{2}H()}{d^{2}} + \cot \frac{dH()}{d} + \frac{m^{2}}{\sin^{2}} - \frac{2B}{\sim^{2}\sin^{2}} H() = 0; \quad (18)$$

and

$$\frac{d^2}{d'^2} + m^2 (') = 0:$$
(19)

where m^2 and are separation constants. The solution of Eq.(18) is

$$_{\rm m} = {\rm Ae}^{\rm im'}; \ {\rm m} = 0; \ 1; \ 2; ...$$
 (20)

We will rst solve the radial part of the Schrödinger equation. If we put R (r) = $\frac{G(r)}{r}$ which is bounded as r ! 0 and Eq.(16) becomes

$$G^{\infty}(\mathbf{r}) + \frac{1}{\mathbf{r}^2} \quad \mathbf{"}^2 \mathbf{r}^2 \quad \mathbf{b}^2 \mathbf{r} \quad \mathbf{(+1)} \quad \mathbf{G}(\mathbf{r}) = 0:$$
 (21)

where $u^2 = \frac{2 E}{r^2}$, $b^2 = \frac{2 A}{r^2}$ and = (+1).

$$\sim (\mathbf{r}) = 0;$$
 $(\mathbf{r}) = \mathbf{r}; \sim (\mathbf{r}) = \mathbf{r}^{2} \mathbf{r}^{2} \mathbf{b}^{2} \mathbf{r} (+1)$ (22)

U sing Eq.(9), (r) is found in four possible values as

$$(\mathbf{r}) = \frac{1}{2} \quad \stackrel{\text{k}}{:} \quad \stackrel{\text{p}}{\frac{\mathbf{r}}{\mathbf{r}^2}} \mathbf{r} + \stackrel{\text{$+\frac{1}{2}$}}{:} ; \quad \mathbf{k}_1 = \mathbf{b}^2 + 2^{\mathbf{p}} \overline{\mathbf{r}^2} (\stackrel{\text{$+\frac{1}{2}$}}{:} ; \\ \stackrel{\text{p}}{\frac{\mathbf{r}}{\mathbf{r}^2}} \mathbf{r} \stackrel{\text{$\frac{1}{2}$}}{:} ; \quad \mathbf{k}_2 = \mathbf{b}^2 - 2^{\mathbf{p}} \overline{\mathbf{r}^2} (\stackrel{\text{$\frac{1}{2}$}}{:} ; \\ \stackrel{\text{$\frac{1}{2}$}}{:} ; \quad \mathbf{k}_2 = \mathbf{b}^2 - 2^{\mathbf{p}} \overline{\mathbf{r}^2} (\stackrel{\text{$\frac{1}{2}$}}{:} ; \\ \stackrel{\text{$\frac{1}{2}$}}{:} ; \\ \stackrel{\text{$\frac{1}{2}$}}{:} ; \quad \mathbf{k}_2 = \mathbf{b}^2 - 2^{\mathbf{p}} \overline{\mathbf{r}^2} (\stackrel{\text{$\frac{1}{2}$}}{:} ; \\ \stackrel{\text{$\frac{1}{2}$}}{:} ; ; \\ \stackrel{\text{$\frac{1}{2}$}}{:} ; ; \\ \\ \stackrel{\text{$\frac{1}{2}$}}{:} ; ; \\ \\ \stackrel{\text{$\frac{1}$$

where $k_{1,2}$ is determined by means of the same procedure as in [15]. From Eq.(6) we obtain

$$(\mathbf{r}) = \begin{pmatrix} \mathbf{p} & \mathbf{n} & \mathbf{r} \\ 1 + 2 \begin{pmatrix} \mathbf{p} & \mathbf{n} & \mathbf{r} \\ \mathbf{n} & \mathbf{r} \\ 1 & 2 \begin{pmatrix} \mathbf{p} & \mathbf{n} & \mathbf{r} \\ \mathbf{n} & \mathbf{r} \\ \mathbf$$

Imposing (r) < 0 [15], for $k_2 = b^2 2^{p - \frac{1}{2}} (r + \frac{1}{2})$ we obtain energy eigenvalues as

$$E_{n_1}; = \frac{A^2}{2^2} \frac{1}{(n_1 + 1)^2}$$
(25)

which agrees with the results [5]. Following the same steps, we nd the weight function as

$$(s) = r^{2^{\nu}+1} e^{\frac{p_{\pi^2}}{r}}$$
(26)

 y_{n_1} is obtained by using Eq.(10)

$$y_{n_1}$$
, (r) = $L_{n_1}^{(2^{\prime}+1)}$ (r) (27)

and also other part of the wave function

$$(r) = r'^{+1} e^{p_{\overline{n^2}r}}$$
(28)

nally, the radial part solution is obtained as

$$G_{n_1;}(\mathbf{r}) = A_{n_1;} \mathbf{r}^{2'+1} e^{p_{\overline{n_2}r}} L_{n_1}^{2'+1}(\mathbf{r})$$
(29)

where A_{n} , is a normalization constant

$$A_{n_1} = \frac{s}{\frac{n_1!}{2(n_1+2!+1)(n_1+2!+1)!}}$$

Hence, these radial solutions are consistent with refs. [5,17].

Now we solve angular part of the Schrödinger equation given in Eq.(17). Dening a new variable as

$$\mathbf{x} = \cos \tag{30}$$

then, Eq.(17) takes the form

$$\int_{\infty}^{\infty} (\mathbf{x}) = \frac{2\mathbf{x}}{1 + \mathbf{x}^2} + \frac{1}{1 + \mathbf{x}^2} + \frac{1}{1$$

To apply the NU method, we de ne the polynom ials

$$\sim = 2x; = 1 x^{2}; \sim = x^{2} + (m^{2})$$
 (32)

where $=\frac{2 B}{r^2}$ and (x) becomes

$$(x) = \begin{cases} 8 & p \\ < & (m^2 + m^2); k_1 = m^2; \\ p & (m^2 + m^2); k_2 = m^2 \end{cases}$$
(33)

where $k_{1,2}$ is determined by means of the same procedure as in [15]. We obtain as

$$(x) = \begin{cases} 2x + 2^{p} + m^{2}; k_{1} = , \\ 2x + 2^{p} + m^{2}; k_{1} = , \\ 2x + 2^{p} + m^{2}; k_{1} = , \\ 2x + 2^{p} + m^{2}x; k_{2} = m^{2} , \\ 2x + 2^{p} + m^{2}x; k_{2} = m^{2} . \end{cases}$$
(34)

We impose $^{\circ}(x) < 0$ because of the physical solutions. For $k_2 = m^2$, we use negative sign of in this case. Thus we obtain

$$r = n_2 + \frac{p_m^2}{m^2 + m_m^2}$$
 (35)

where we use = ('+1). In order to get the wave function of the polar angle of Schrödinger equation, we use Eqs.(3) and (10-12) and one obtains,

$$= (1 \quad x^2)^{\frac{1}{2}} p_{\frac{m^2}{m^2}}$$
(36)

and

$$= (1 \quad x^2)^{p_{\frac{m^2}{m^2}}}$$
(37)

$$y_n = B_n (1 x^2)^p \frac{d^n}{dx^n} ((1 x^2)^n)^p \frac{d^n}{dx^n}$$
 (38)

where $B_{\,n}\,$ is a norm alization constant and the wave function is given as

$$n_{2};$$
 () = $C_{n_{2}} (\sin)^{m^{0}} P_{n_{2}}^{(m^{0},m^{0})} (\cos)$ (39)

where $m^{\circ} = \frac{p}{m^2 + \cdots}$ and C_{n_2} is a norm alization constant. Eq.(39) agrees with the results [5]. C_{n_2} is given as:

$$C_{n_{2}} = \frac{s}{\frac{(2n_{2} + 2m^{\circ} + 1)(n_{2} + 1)(n_{2} + 2m^{\circ} + 1)}{2^{2m^{\circ} + 1}(2n_{2} + m^{\circ} + 1)(n_{2} + m^{\circ} + 1)}}$$

Therefore, the total energy of the system in Eq.(13) is given as (in a.u: $= e = - = 1; a_0 = 1; a_0 = 1=2$)

$$E_{(n_1,n_2),m} = \frac{p_1^2}{2(m^2 + m_1 + n_2 + 1)^2}$$
(40)

which compares with ref.[1].

4 Conclusions

We have solved Schrödinger equation for the ring shaped potential. The energy eigenvalues and the corresponding eigenfunctions are obtained exactly by using NU -m ethod. E igenfunctions are expressed in terms of Laguerre and Jacobi polynom ials for radial and angular parts respectively. It is seen that NU m ethod is an applicable tool for not only central potentials but also noncentral and com bined potentials. Results are in good agreem ent with the earlier works [1,5,6]. Som e num erical results are given in Table 1. It is pointed out that for the values of B = 0, the results are in good agreem ent with ref. [6,25].

5 A cknow ledgem ents

This research was partially supported by the Scienti c and Technological Research Council of Turkey.

References

- [1] Kibler M., Winternitz P., J. Phys. A 20 4097 (1987).
- [2] C.Quesne, J.Phys. A 21 3093 (1988).
- [3] A.A.Makarov et al., Nuovo Cimento A 52 1061 (1967).
- [4] H. Hartmann, D. Schuch, Int. J. Quantum Chem. 18 125 (1980); H. Hartmann, Theor.
 Chim. Acta 24 201 (1972).
- [5] F.Yasuk et al, J.Phys.A: Math.Gen. 38 6579-6586 (2005).
- [6] M Aktas R. Sever, Journal of M ath. Chem. 37 139 (2005).
- [7] S.M. Ikhdair, R. Sever, ArX iv/quant-ph/0603205.
- [8] Y.Aharonov, D.Bohm, Phys. Rev. 115 485491 (1959).
- [9] I. Sokmen, Physics Letters A 132 65 (1988); G.G.Blado, Theo. Chim. Acta, 94 53-66 (1996); B.M. Mandal, Int. J. of M od. Phys. A 15 1225 (2000).
- [10] K.Li, S.Dulat, Eur. Phys. J.C 46 825-828 (2006).
- [11] M S.Shikakhwa, N K Pak, Phys. Rev. D 67 105019 (2003).
- [12] A.D.A Ihaidari, Annals Phys. 320 453 (2005); Erratum -ibid. 321 1524-1525 (2006).
- [13] A.D.A lhaidari, Phys. Lett. 18 651-664 (2005).
- [14] A.D.Alhaidari, J.Phys.A 38 3409 (2005).
- [15] A F.N ikiforov and V B.Uvarov, Special Functions of M athem atical Physics (B irkhauser, Basel, 1988).
- [16] O.Yesiltas, M.Simsek, R.Sever, C.Tezcan, Physica Scripta 67 472 (2003).
- [17] M.Aktas, R.Sever, J. of Mol. Struct.: THEOCHEM 71 223-228 (2004).

- [18] C. Berkdem ir, A. Berkdem ir, R. Sever, Phys. Rev. C 72 027001 (2005).
- [19] A.Berkdem ir, C.Berkdem ir, R.Sever, Mod.Phys.Lett. A 21 (27) 2087-2097 (2006).
- [20] O.Yesiltas, Phys. Scr. 75 (2007) 4146.
- [21] F.Yasuk, C.Berkdem ir, A.Berkdem ir, C.Onem, Physica Scripta 71 (4) 340 (2005).
- [22] H.Egrifes, R.Sever, Phys.Lett.A 344 117-126 (2005).
- [23] M. Sim sek, H. Egrifes, J. Phys. A: Math. Gen. 37 4379-4393 (2004).
- [24] A D. A lhaidari, H. Bahlouli, A. Al Hasan, Phys. Lett. A 349 8797 (2006).
- [25] A.Guha, S.Mukherjee, J.Math.Phys., 28 (1987) 340.

n_1	n ₂	m	E
1	1	0	-0.849763
2	2	0	-0.377661
		1	-0.330452
3	3	2	-0.178302
		1	-0.192028
		0	-0,212431
4	4	3	-0.112357
		2	-0.118038
		1	-0.125353
		0	-0.135954
5	5	4	-0.0776005
		3	-0.0804445
		2	-0.0838663
		1	-0.0882161
		0	-0.0944122

Table 1: Energy levels for $^2 = Z$, = 1