

Selective continuous quantum measurements: Restricted path integrals and wave equations *

Lajos Diósi [†]

KFKI Research Institute for Particle and Nuclear Physics
H-1525 Budapest 114, POB 49, Hungary

bulletin board ref.: quant-ph/9501009

Abstract

We discuss both the restricted path integral (RPI) and the wave equation (WE) techniques in the theory of continuous quantum measurements. We intend to make Mensky's fresh review complete by transforming his "effective" WE with complex Hamiltonian into Ito-differential equations.

* *submitted to Phys. Lett. A*

[†]E-mail: diosi@rmki.kfki.hu

1 Introduction

Quite recently, a short review on two mathematical techniques of continuous quantum measurements has been presented by Mensky [1]. The basic ideas and formal elements of the restricted path integral (RPI) method (Sect.2) and the master equation (ME) method (Sect.3), respectively, have been explained. Selective (Sect.4) and non-selective (Sect.5) measurements have been discussed. The former, being the more complicated one, only has been given a very brief outline anticipating an efficient method of Schrödinger-equation replacing circuitous path integrals. In the present Letter we bring out this efficient wave equation (WE) which has emerged from a great body of earlier works.

The idea of RPI to model continuous measurements originates from 1979 [2]. Elaborating of the *statistical* theory within RPI have been promoted basically by Refs. [3, 4, 5], in addition to the papers cited in Ref. [1]. The proof of the correspondence between the RPI and the Ito-stochastic WE formalism (5ab) was given first in 1988 [4].

2 Selective measurements

Let us start with recapitulating Mensky's RPI formalism [1]. In his notations, the state vector of the system under continuous measurement evolves as $\psi_t^\alpha = U_t^\alpha \psi_0$, where the propagator is expressed by the following RPI:

$$U_t^\alpha(q'', q') = \int_\alpha d[p]d[q] \exp \left(\frac{i}{\hbar} \int_0^t [p\dot{q} - H(p, q, t)] dt \right). \quad (1)$$

The output of continuous measurement is labelled by α , and its probability distribution is given by

$$P(\alpha) = \langle \psi_t^\alpha | \psi_t^\alpha \rangle. \quad (2)$$

(Interestingly, neither this statement nor the fact that ψ_t^α is unnormalized do appear in Ref. [1] explicitly.) For the simple case of continuous monitoring an observable $A(p, q, t)$, the propagator's RPI reduces to unrestricted path integral:

$$\begin{aligned} & U_t^{[a]}(q'', q') \\ &= \int d[p]d[q] \exp \left(\frac{i}{\hbar} \int_0^t [p\dot{q} - H(p, q, t)]dt - \kappa \int_0^t [A(p, q, t) - a(t)]^2 dt \right) \mathcal{B} \end{aligned}$$

A comparison of this path integral with Feynman's standard ones leads to the naive conjecture that "one may forget about any path integrals and reduce the problem to solving the Schrödinger-equation with a complex Hamiltonian" [1]:

$$\frac{\partial}{\partial t} \psi_t = \left(-\frac{i}{\hbar} H(p, q, t) - \kappa [A(p, q, t) - a(t)]^2 \right) \psi_t \quad (4)$$

where a functional dependence of ψ_t on $[a] \equiv \{a(t') | 0 \leq t' \leq t\}$ is understood though has not been denoted explicitly.

It is inevitable to note that this Schrödinger-equation is unconventional: it is *not* linear, norm-conserving, deterministic, and regular either. Its stochasticity is obvious from the fact that the effective Hamiltonian depends on the measurement output record $[a]$ whose probability distribution still depends on the state vector via the Eq. (2): $P[a] = \|\psi_t\|^2$. *Nevertheless, these coupled functional equations can be disentangled into two separate*

stochastic equations: one for the state vector and another for the measurement output [4]. It turns out that both ψ_t and $a(t)$ are Wiener-processes rather than regular functions of t :

$$\frac{\partial}{\partial t}\psi_t = \left(-\frac{i}{\hbar}H(p, q, t) - \frac{\kappa}{2}[A(p, q, t) - \langle A \rangle_t]^2 + \sqrt{\kappa}[A(p, q, t) - \langle A \rangle_t]\dot{\xi}_t \right) \psi_t \quad (5a)$$

$$a(t) = \langle A \rangle_t + \frac{1}{2\kappa}\dot{\xi} \quad (5b)$$

where $\langle A \rangle_t$ denotes the expectation value of the observable $A(p, q, t)$ in the quantum state ψ_t . The "function" ξ_t is the standard Wiener-process whose time-derivative $\dot{\xi}_t$ is the standard white-noise with $\delta(t)$ as auto-correlation.

The Eqs. (5ab) provide a radical improvement as compared to the naive WE (4); explicit solutions become available for certain special cases like, e.g., for continuous position measurement of free particles [4].

3 Concluding remarks

It should be noted that, alternatively to RPI, a theory of quantum filtering has been devised to model continuous measurement and resulted in equations mathematically equivalent to (5ab) [6]. Furthermore, investigating the so-called quantum measurement problem have led to important components of the formalism explained in Ref. [1] and in the present Letter. Simple stochastic WE of structure (5a) was found heuristically [7] without any underlying model like e.g. RPI. The RPI technique itself was used independently e.g. in Refs. [8, 9, 10]. In the context of quantum measurement

theory, there is a fruitful co-existence of (a specific version of) RPI and WE techniques, as pointed out recently [11].

This work was supported by the grant OTKA No. 1822/1991.

References

- [1] M Mensky, Phys. Lett. A 196 (1994) 159.
- [2] M.B. Mensky, Phys. Rev. D 20 (1979) 384.
- [3] A. Barchielli, L. Lanz and G.M. Prosperi, Nuovo Cim. 72 B (1982) 79;
C.M. Caves and G.J. Milburn, Phys. Rev. A 36 (1987) 5543.
- [4] L. Diósi, Phys. Lett. A 129 (1988) 419; A 132 (1989) 233.
- [5] L. Diósi, Phys. Rev. A 42 (1990) 5086.
- [6] V.P. Belavkin, Phys. Lett. A (1989) 335; A. Barchielli and V.P. Belavkin, J. Phys. A (1991) 1495; V.P. Belavkin and P. Staszewski, Phys. Rev. A 45 (1992) 1347.
- [7] N. Gisin, Phys. Rev. Lett. 52 (1984) 1567.
- [8] L. Diósi, Phys. Rev. A 40 (1989) 1165.
- [9] G.C. Ghirardi, R. Grassi and P. Pearle, Phys. Rev. A 42 (1990) 1057.
- [10] J. B. Hartle, Phys. Rev. D 44 (1991) 3173.

- [11] L. Diósi, N. Gisin, J. Halliwell and I.C. Percival, Phys. Rev. Lett. (to appear)