

Sub-femtosecond determination of transmission delay times for a dielectric mirror (photonic bandgap) as a function of angle of incidence

Aephraim M. Steinberg and Raymond Y. Chiao

Department of Physics, U.C. Berkeley, Berkeley, CA 94720

Internet: aephraim@physics.berkeley.edu

(Preprint quant-ph/9501013; Received Phys. Rev. 17 August, 1994)

Using a two-photon interference technique, we measure the delay for single-photon wavepackets to be transmitted through a multilayer dielectric mirror, which functions as a "photonic bandgap" medium. By varying the angle of incidence, we are able to confirm the behavior predicted by the group delay (stationary phase approximation), including a variation of the delay time from superluminal to subluminal as the band edge is tuned towards to the wavelength of our photons. The agreement with theory is better than 0.5 femtoseconds (less than one quarter of an optical period) except at large angles of incidence. The source of the remaining discrepancy is not yet fully understood.

PACS numbers: 03.65.Bz, 42.50.Wm, 73.40.Gk

In recent years, there has been a great deal of interest in two related topics: tunneling times [1{4] and photonic bandgaps [5{7]. A standard quarter-wave-stack dielectric mirror is in fact the simplest example of a one-dimensional photonic bandgap, and consequently may be thought of as a tunnel barrier for photons within its "stop band." The periodic modulation of the refractive index is analogous to a periodic Kronig-Penney potential in solid-state physics, and leads to an imaginary value for the quasimomentum in certain frequency ranges (that is, to an exponentially decaying field envelope within the medium, and high reflectivity due to constructive interference (Bragg reflection). We have exploited this analogy to perform the first measurement of the single-photon tunneling delay time [8{11], using as our barrier an 11-layer mirror of alternating high ($n=2.22$) and low ($n=1.41$) index quarter-wave layers, with minimum transmission of about 1% at the center of the bandgap. We confirmed the striking prediction which drives the tunneling time controversy: in certain limits, a transmitted wave packet peak may appear on the far side of the barrier faster than if the peak had traversed the barrier at the vacuum speed of light c . Meanwhile, several microwave experiments have investigated other instances of superluminal propagation including electromagnetic analogies to tunneling [12{17].

While in itself, this anomalous peak propagation does not constitute a violation of Einstein causality [18{27], it certainly leads one to ask whether there may exist another, longer timescale in tunneling, with more physical significance than the group (i.e., peak) delay. After all, in a certain sense, the bulk of the transmitted wave originates in the leading edge of the incident wave packet, not near the incident peak [28,23,29]. Many theories have been propounded to describe the duration of the tunneling interaction, and the leading contenders involve studying oscillating barriers [1,30] or Larmor precession of a tunneling electron in a barrier with a confined magnetic field [31{33]. It should be stressed that these theories are not intended to describe the propagation of wave packets, but rather the dynamical timescale of the tunneling process; several experiments have supported their predictions [34].

Nevertheless, there is a popular misconception that these times (and in particular the Buttiker-Landauer time in its "semiclassical" or WKB limit \hbar/Γ , where Γ represents the evanescent decay constant inside the barrier, i.e., the magnitude of the imaginary wavevector) predict the arrival time of wave packets. In [8], we were able to exclude the semiclassical time, but not Buttiker's version of the Larmor time, as describing peak propagation. Furthermore, some workers have expressed concern about the paucity of data supporting the superluminality of the group delay, in spite of our finding of a seven-standard-deviation effect. Microwave experiments have also traditionally been met with skepticism (see, for example, [35].) In light of these objections, we have extended the earlier experiment to study the delay time as a function of angle-of-incidence. As the angle is changed, the frequency and the width of the bandgap change as well, so this is essentially a way to study the energy-dependence of the tunneling time.

Our apparatus is shown in Fig. 1. As the technique and the sample have both been described at length elsewhere [36,37,8,38,39], we will content ourselves with an abbreviated sketch of the method. A crystal with an optical⁽²⁾ nonlinearity is pumped by a cw ultraviolet laser, and in the process of spontaneous parametric down-conversion emits simultaneous pairs of horizontally-polarized infrared photons. The two photons in each pair leave the crystal on opposite sides of the ultraviolet pump, conserving momentum. They are correlated in time to within their reciprocal

As of 1995, at National Institute of Standards and Technology, Phys A 167, Gaithersburg, MD 20899

bandwidth of about 15 fs. They are also correlated in energy, their frequencies summing to that of the (narrow-band) 351 nm pump. When the two photons arrive simultaneously at a beam splitter, there is no way to distinguish the two Feynman paths leading to coincidences between detectors placed at the beam splitter's two output ports: both photons being transmitted, and both photons being reflected. This leads to an interference effect in which the coincidence rate is suppressed (the two photons tending to head off to the same detector). By contrast, if the photons arrive at the beam splitter at different times (on the scale of their 15 fs correlation time), coincidence counts occur half the time. Thus by placing a dielectric mirror in one arm of the interferometer and adjusting the external path length to minimize the coincidence rate, we can measure the delay experienced by the photon wavepackets which are transmitted through this barrier. We find that near the transmission minimum, the photons travel through the mirror faster than they travel through an equivalent length of air, whereas when the mirror is angled to bring the band-edge closer to the photons' wavelength, they travel slower than through air, as one would expect. Figure 2 shows sample data for these two situations, where the sign change can be clearly seen.

In [8], our results were consistent with the group delay predictions, and also with Buttiker's proposed Lamor time [33], but not with the "semiclassical" time. The measured times exceeded the predictions by approximately 0.5 fs, but this result was at the borderline of statistical significance, and not discussed. Since then, further data taken at various angles of incidence have continued to show a discrepancy, ranging from an excess of 0.5 fs near normal incidence to a deficit of over 1 fs at large angles of incidence. At the same time, the data are in close agreement with the group delay, and appear to rule out identification of the Lamor theory with a peak propagation time. Our attempts to eliminate systematic effects and characterize those which remain were described in [8]. Since then, unable to find any other sources of error to explain the discrepancy, we are convinced that it is a property of the sample under study, and not of the interferometer used for the measurements. We therefore obtained a second dielectric mirror of design parameters identical to the first, to see whether the errors could be attributed to deviations from the ideal quarter-wave-stack structure. As can be seen from Figure 3, both mirrors show quite similar behavior. Both are 11-layer quarter-wave stacks as described above. Mirror 1 shows a transmission minimum at 692 nm, while mirror 2's minimum is at 688 nm; this difference is insignificant on the scale of the bandgap, which extends from 600 nm to 800 nm. We conclude that some real effect is at work, modifying the stationary-phase prediction. In principle, frequency-dependent transmission could lead to such an effect, as does second-order group-velocity dispersion; both effects are much too small to explain the present discrepancy. As discussed in [40], attempts to numerically model dielectric mirrors with small, random fluctuations in layer thicknesses were able to produce deviations on the right order, but in general they did not lead to deviations of the form we observed experimentally. It is conceivable that loss or scattering in the dielectrics could also help explain the effect, and we are beginning to investigate this possibility [41]; see also [42,43].

Theoretical curves are plotted along with the data in Figures 3 and 4. The group delay is calculated by the method of stationary phase. The transmission phase of the 11-layer structure is calculated numerically, and differentiated first with respect to angle of incidence to give the transverse shift and then with respect to incident frequency to give the time delay, according to the formulas $y = \partial T / \partial k_y = \partial T / \partial (k \sin \theta) = (k \cos \theta) \partial T / \partial \theta$ and $\tau_g = \partial T / \partial \omega = (y/c) \sin \theta$, where T is the transmission phase [44]. Buttiker's Lamor time [33] is equal to the magnitude of the complex time [45,46] $\tau_c = i \partial (\ln t) / \partial \omega = \tau_L$, where t is the complex transmission amplitude, and τ_L the Lamor frequency. For our optical structure, an effective Lamor frequency τ_L corresponds to a uniform (over the barrier region) scaling of the local index of refraction by a factor of $1 + \tau_L = 1$. Since in the limit of interest, the "in-plane portion" of the Lamor time (i.e., the real part of the complex time) differs little from the group delay, we take them to be equal in order to include the effects of the transverse shift in the Lamor theory. The "out-of-plane portion" (or imaginary part) is calculated numerically, and added in quadrature to the group delay in order to generate the Lamor time. Since our measurements compare the transit time through the barrier with that through air, we subtract the time parallel wavefronts propagating at c would take to reach a point on the far side of the barrier (with a transverse shift of y) from both the group delay and the Lamor time, so as to facilitate comparison with the experimental data.

At the moment, more work (both experimental and theoretical) is needed to understand the discrepancy. We are therefore planning to repeat this experiment with s-polarized light (by introducing half-wave plates before and after the sample being studied), which has very different transmission characteristics as a function of angle (also leading to a larger difference between the group delay and the Lamor theories). As shown in Figure 4, our preliminary data are again consistent with the group delay and not with the Lamor time, but due to the lower transmission for this polarization, we need to improve our signal-to-noise ratio before reaching any definitive conclusions.

The superluminality of the barrier traversal near mid-gap is now well supported by the data, and the group delay (stationary-phase) theory can be seen to be relatively accurate for a variety of angles of incidence, but there is a residual discrepancy on the order of 0.5 femtoseconds, which is not yet fully understood.

This work was supported by the U.S. Office of Naval Research under grant N00014-90-J-1259. We would like to thank P. G. Kwiat, M. M. Mitchell, B. Johnson, G. McKinney, and J. Holden for their assistance; W. Heilmann for discussions about data analysis; and R. Landauer, J. D. Jackson, and E. Yablonovitch for useful disagreements.

Notes added during revision: Since the submission of this manuscript, a paper has appeared [47] extending our previous experimental results to barriers of varying thicknesses (and transmission as low as 0.01%) near normal incidence, using classical femtosecond pulses. It reports general agreement with the group delay theory, aside from a discrepancy on the order of 1.5 fs. Two papers have also appeared discussing the effects of dissipation on tunneling times [48,49].

-
- [1] M. Buttiker and R. Landauer, Phys. Rev. Lett. 49, 1739 (1982).
 - [2] E. H. Hauge and J. A. Stvneng, Rev. Mod. Phys. 61, 917 (1989).
 - [3] M. Buttiker, In "Electronic Properties of Multilayers and Low Dimensional Semiconductors," J.M. Chamberlain, L. Eaves, and J.C. Portal, eds., Plenum, New York (1990).
 - [4] R. Landauer and Th. Martin, Rev. Mod. Phys. 66, 217 (1994).
 - [5] E. Yablonovitch, J. Opt. Soc. Am. B 10 (2), 283 (1993), and references therein, and other articles in this special issue on photonic bandgaps.
 - [6] S. John, Physics Today 44, 32 (1991).
 - [7] C.M. de Sterke and J.E. Sipe, Phys. Rev. A 38, 5149 (1988).
 - [8] A.M. Steinberg, P.G. Kwiat, and R.Y. Chiao, Phys. Rev. Lett. 71, 708 (1993).
 - [9] R.Y. Chiao, P.G. Kwiat, and A.M. Steinberg, "Faster Than Light?" Scientific American 269, 52 (1993).
 - [10] R. Landauer, Nature 365, 692 (1993).
 - [11] J.A. Stvneng and E.H. Hauge, Phys. World 6, 23 (1993).
 - [12] T.K. Ishii and G.C. Giakos, Microwaves and RF 30, 114 (1991) and references therein.
 - [13] A. Ranfagni, D. Mugnai, P. Fabeni, and G.P. Pazzi, Appl Phys Lett. 58, 774 (1991).
 - [14] A. Enders and G. Nintz, J. Phys. I France 3, 1089 (1993).
 - [15] A. Ranfagni, P. Fabeni, G.P. Pazzi, and D. Mugnai, Phys. Rev. E 48, 1453 (1993).
 - [16] G. Nintz, A. Enders, and H. Spieker, J. Phys. I France 4, 565 (1994).
 - [17] A.M. Steinberg, J. Phys I France 4, 1813 (1994).
 - [18] L. Brillouin, Wave Propagation and Group Velocity. Academic, New York (1960).
 - [19] C.G.B. Garrett and D.E. McCumber, Phys. Rev. A 1, 305 (1970).
 - [20] S. Chu and S. Wong, Phys. Rev. Lett. 48, 738 (1982).
 - [21] S. Chu and S. Wong, Phys. Rev. Lett. 49, 1292 (1982).
 - [22] E.L. Bolda, J.C. Garrison, and R.Y. Chiao, Phys. Rev. A 49, 2938 (1994).
 - [23] J.M. Deutsch and F.E. Low, Ann. Phys. 228, 184 (1993).
 - [24] Y. Japha and G. Kurizki, "Unified Causal Theory of Photonic Superluminal Delays in Dielectric Structures," submitted to Phys. Rev. Lett., 1994.
 - [25] G. Kurizki and Y. Japha, "Causality and Interference in Electron and Photon Tunneling," to appear in Quantum Interferometry, proceedings of the Adriatico Research Workshop on Quantum Interferometry (Trieste, March 1993), organized by F. de Martini and A. Zeilinger.
 - [26] K. Hass and P. Busch, Phys. Lett. A 185, 9 (1994).
 - [27] Mark Ya. Azbel, Sol. St. Comm. 91, 439 (1994).
 - [28] C.R. Leavens and G.C. Aers. In "Scanning Tunneling Microscopy III," R. Wiesendanger and H.-J. Guntherodt, eds., Springer-Verlag, Berlin (1993).
 - [29] A.M. Steinberg, P.G. Kwiat, and R.Y. Chiao, Found. Phys. Lett. 7, 223 (1994).
 - [30] T. Martin and R. Landauer, Phys. Rev. A 47, 2023 (1993).
 - [31] A.I. Baz', Sov. J. Nucl. Phys. 5, 161 (1967).
 - [32] V.F. Rybachenko, Sov. J. Nucl. Phys. 5, 635 (1967).
 - [33] M. Buttiker, Phys. Rev. B 27, 6178 (1983).
 - [34] R. Landauer, Nature 341, 567 (1989).
 - [35] K.D. Stephan, IEEE Antennas and Propagation Magazine 35, 13 (1993).
 - [36] C.K. Hong, Z.Y. Ou, and L.M.andel, Phys. Rev. Lett. 59, 2044 (1987).
 - [37] A.M. Steinberg, P.G. Kwiat, and R.Y. Chiao, Phys. Rev. A 45, 6659 (1992).
 - [38] J. Jeers and S.M. Barnett, Phys. Rev. A, 47, 3291 (1993).
 - [39] J.H. Shapiro and K.-X. Sun, J. Opt. Soc. Am. B 11, 1130 (1994).
 - [40] A.M. Steinberg, When Can Light Go Faster Than Light? The tunneling time and its sub-femtosecond measurement via quantum interference, Ph.D. thesis, University of California at Berkeley, 1994.
 - [41] A.M. Steinberg, "Conditional probabilities in quantum theory, and the tunneling time controversy," submitted to Phys. Rev. A (1994).


- [42] A. Ranfagni, D. Mugnai, P. Fabeni, and G. P. Pazzi, *Physica Scripta* 42, 508 (1990).
- [43] D. Mugnai, A. Ranfagni, R. Ruggeri, and A. Agresti, *Phys. Rev. Lett.* 68, 259 (1992).
- [44] A. M. Steinberg and R. Y. Chiao, *Phys. Rev. A* 49, 3283 (1994).
- [45] D. Sokolovski and J. N. L. Connor, *Phys. Rev. A* 47, 4677 (1993).
- [46] A. M. Steinberg, "How much time does a tunneling particle spend in the barrier region?", to appear in *Phys. Rev. Lett.* (1995).
- [47] Ch. Spielmann, R. Szepcs, A. Stingl, and F. Krausz, *Phys. Rev. Lett.* 73, 2308 (1994).
- [48] G. Nimtz, H. Spieker, and H. M. Brodowsky, *J. Phys. France* 4, 1379 (1994).
- [49] F. Raciti and G. Salesi, *J. Phys. France* 4, 1783 (1994).

Figure Captions

1. Experimental setup for determining single-photon propagation times through a multilayer dielectric mirror. By translating the sample, we can observe the interference dip for photons tunneling through the 1.1 μm barrier or for photons traversing an equal thickness of air. We can thus compare arrival times for tunneling and freely propagating wave packets.
2. Coincidence rate versus trombone prism position (see Fig. 1) for p-polarized photons travelling through the reflective coating as well as for those travelling through an equal thickness of air, for (a) normal incidence and (b) 55-degree incidence.
3. Left axis: measured delay for mirror one (squares) and mirror two (circles) as a function of angle of incidence, to be compared with the theoretical group delay and the Lamor interaction time proposed by Buttiker. Right axis: transmission versus angle of incidence. All curves are for p-polarization.
4. Same as Fig. 3, but for s-polarization. Due to the much lower transmission, only one preliminary data point is shown, but the different characteristics of the theoretical curves for both transmission and delay suggest that upon improvement of our signal-to-noise ratio, further work in this direction may help elucidate the discrepancy between experiment and theory.

This figure "fig1-1.png" is available in "png" format from:

<http://arxiv.org/ps/quant-ph/9501013v1>



This figure "fig2-1.png" is available in "png" format from:

<http://arxiv.org/ps/quant-ph/9501013v1>

This figure "fig3-1.png" is available in "png" format from:

<http://arxiv.org/ps/quant-ph/9501013v1>

This figure "fig4-1.png" is available in "png" format from:

<http://arxiv.org/ps/quant-ph/9501013v1>

This figure "fig5-1.png" is available in "png" format from:

<http://arxiv.org/ps/quant-ph/9501013v1>

