

Physical Review Letters 74 (1995) 4571; 76 (1996) 2205 (erratum)

## Nonlocal effects in Fock space

Asher Peres

Department of Physics, Technion | Israel Institute of Technology,  
32 000 Haifa, Israel

25 January 1995

### Abstract

If a physical system contains a single particle, and if two distant detectors test the presence of linear superpositions of one-particle and vacuum states, a violation of classical locality can occur. It is due to the creation of a two-particle component by the detecting process itself.

PACS: 03.65.Bz

Electronic address: peres@photon.technion.ac.il

It has been known for a long time that quantum systems consisting of two [1] or more [2, 3] distant particles display remarkable nonlocal effects. Recently, a similar nonlocal effect was predicted by Hardy [4] for a quantum system involving no more than one photon. Hardy considered a state

$$|j\rangle_i = p |j\rangle_{i_a} |0\rangle_{i_b} + q |0\rangle_{i_a} |j\rangle_{i_b} + r |0\rangle_{i_a} |0\rangle_{i_b}; \quad (1)$$

and various tests performed by two distant observers, Alice and Bob, who find a violation of classical locality. In the above equation,  $|0\rangle_{i_a}$  and  $|j\rangle_{i_a}$  denote the vacuum and one-particle states in a beam directed toward Alice;  $|0\rangle_{i_b}$  and  $|j\rangle_{i_b}$  likewise refer to a beam directed toward Bob; and  $p, q, r$  are numerical coefficients, none of which is zero. Hardy gave explicit instructions on how to actually perform these experiments, by means of beam splitters and parametric down conversion processes. The abundance of technical details is helpful for convincing the reader that the experiment is indeed feasible, but it somewhat obscures the origin of the nonlocality.

As shown below, the latter is simply due to the creation of a component  $|j\rangle_{i_a} |j\rangle_{i_b}$  by the detecting process itself. To simplify the discussion, I shall restrict it to the case where

$$|j\rangle_i = (|j\rangle_{i_a} |0\rangle_{i_b} - |0\rangle_{i_a} |j\rangle_{i_b}) / \sqrt{2}; \quad (2)$$

is a pure one-particle state, without vacuum component. Such a state could also be written without invoking Fock space notations, since it involves only ordinary quantum mechanics, with a given number of particles (one). However, it is impossible to repeat Hardy's argument by using a first-quantized formalism, for reasons that will soon be clear.

Note that the right hand side of Eq. (2) has the same structure as the singlet state of a pair of particles of spin  $\frac{1}{2}$ , if we reinterpret  $|0\rangle_{i_a}$  and  $|j\rangle_{i_a}$  as

representing particle a with spin up and down, respectively, and likewise for particle b. The route to nonlocality is now obvious.

Both Alice and Bob have a choice of two different experiments. One is to test the mere presence of a particle, by measuring the projection operators  $P_a$  and  $P_b$  on the one-particle states  $|j_i\rangle_a$  and  $|j_i\rangle_b$ , respectively. Alice can also opt to test the projection operator  $P_{a^0}$  on the state  $\frac{1}{2}(|j_i\rangle_a + \sqrt{3}|0\rangle_a)$ , namely a coherent superposition of one-particle and vacuum states. Independently of her decision, Bob can choose to test  $P_{b^0}$ , the projection operator on  $\frac{1}{2}(|j_i\rangle_b + \sqrt{3}|0\rangle_b)$ . There are therefore four different experiments, and quantum theory makes the following predictions, for the state  $|j_i\rangle$  in Eq. (2):

$$\langle P_{a^0} \rangle = \langle P_{b^0} \rangle = 0.5; \quad (3)$$

$$\langle P_a P_b \rangle = 0; \quad (4)$$

$$\langle P_a P_{b^0} \rangle = \langle P_{a^0} P_b \rangle = \langle P_{a^0} P_{b^0} \rangle = 0.375; \quad (5)$$

These results violate the Clauser-Horne inequality [5] (a variant of Bell's inequality), namely

$$0 \leq \langle P_{a^0} + P_{b^0} - P_{a^0} P_{b^0} - P_{a^0} P_b - P_a P_{b^0} + P_a P_b \rangle \leq 1; \quad (6)$$

For the given  $|j_i\rangle$ , the actual value of the above expression is 0.125.

Obviously, the total number of particles is not conserved when we measure  $P_{a^0}$  or  $P_{b^0}$ , since these operators do not commute with the number operator. The apparatuses used by Alice and Bob must be able to create new particles, or supply some of their own. Nonlocal effects may thus appear for an initial state that contains a single particle, provided that the final state may contain two. (I did not include in this discussion the numerous auxiliary particles in the two measuring apparatuses, as Hardy did in ref. [4], because quantum

mechanical probabilities do not depend on the detailed structure of these apparatuses, and it is both customary [13] and legitimate [6] to ignore the latter.)

This work was supported by the Gerard Swope Fund, and the Fund for Encouragement of Research.

1. J. S. Bell, *Physics* 1, 195 (1964).
2. D. M. Greenberger, M. Home, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, ed. by M. Kafatos, Kluwer, Dordrecht (1989) p. 69.
3. N. D. Mermin, *Am. J. Phys.* 58, 731 (1990).
4. L. Hardy, *Phys. Rev. Letters* 73, 2279 (1994).
5. J. F. Clauser and M. A. Home, *Phys. Rev. D* 10, 526 (1974).
6. A. Peres, *Quantum Theory: Concepts and Methods*, Kluwer, Dordrecht (1993), Chapt. 12.