

## Conditional probabilities in quantum theory, and the tunneling time controversy

Aephraim M. Steinberg

Department of Physics, U.C. Berkeley, Berkeley, CA 94720

(Preprint quant-ph/9502003; Submitted to Phys. Rev. A on July 19, 1994)

It is argued that there is a sensible way to define conditional probabilities in quantum mechanics, assuming only Bayes's theorem and standard quantum theory. These probabilities are equivalent to the "weak measurement" predictions due to Aharonov et al., and hence describe the outcomes of real measurements made on subensembles. In particular, this approach is used to address the question of the history of a particle which has tunneled across a barrier. A gedankenexperiment is presented to demonstrate the physically testable implications of the results of these calculations, along with graphs of the time-evolution of the conditional probability distribution for a tunneling particle and for one undergoing allowed transmission. Numerical results are also presented for the effects of loss in a bandgap medium on transmission and on reflection, as a function of the position of the lossy region; such loss should provide a feasible, though indirect, test of the present conclusions. It is argued that the effects of loss on the pulse delay time are related to the imaginary value of the momentum of a tunneling particle, and it is suggested that this might help explain a small discrepancy in an earlier experiment.

PACS numbers: 03.65.Bz, 73.40.Gk

The question of how much time a tunneling particle spends in the barrier region has long been controversial, in part because it requires one to discuss not an entire ensemble of identically prepared particles, but only the subset of particles which are later found to have tunneled [1-13]. The absence of an unambiguous prescription in quantum theory for dealing with such questions has compounded the more technical problems, such as the negative kinetic energy of a tunneling particle and the superluminal effects arising from the asymptotic independence of the wave packet delay on barrier thickness [14-20].

In essence, what is desired is a conditional probability distribution. We know how to ask what the probability is for a particle from a given ensemble to be at point  $x$  at time  $t$ . Now we wish to ask for that same probability, conditioned on the additional information that at  $t = 1$ , the particle is found to have been entirely transmitted. Conventional wisdom holds that such distributions do not exist in quantum mechanics; we can never have more information about a particle than its wave function. Let us proceed nevertheless.

Consider the joint probability for two propositions  $A$  and  $B$ ,  $P(A \& B)$ . This can be written as the probability of  $B$ ,  $P(B)$ , multiplied by the probability of  $A$  given  $B$ ,  $P(A|B)$ . This is known as Bayes's theorem, and will serve as our definition of conditional probabilities:

$$P(A|B) = \frac{P(A \& B)}{P(B)} : \quad (1)$$

Now in quantum mechanics, the probability for a particle to be in a given state can be expressed as the expectation value of the projector onto that state;

$$\begin{aligned} P(A) &= \langle \text{proj}(A) \rangle = \langle \hat{A} \rangle \\ &= \langle \hat{A} \rangle = \langle \hat{A} \rangle : \end{aligned} \quad (2)$$

For example, the probability that a particle is at point  $x$  at time  $t$  is given by  $\langle \text{proj}_x(t) \rangle = \langle \delta(x - \hat{x}(t)) \rangle$ . From a statistics perspective, we are looking for the average value of a Boolean variable (the projector, whose eigenvalues are 0 and 1), and interpreting the result as a frequency, or a probability. If we wish to calculate the joint probability of  $A$  and  $B$ , it suffices to find the average value of the product of these two Boolean variables:

$$P(A \& B) = \langle \text{proj}(B) \text{proj}(A) \rangle = \langle \hat{B} \hat{A} \rangle \quad (3)$$

---

Internet aephraim@physics.berkeley.edu; Current address: Atomic Physics Division, National Institute of Standards and Technology, Gaithersburg, MD 20899

(where we have placed B to the left of A to indicate that it is to be determined after A in time). This is where quantum mechanics diverges from standard probability theory, for the product of two Hermitian operators need not be Hermitian (in other words,  $P(A \& B)$  is in general complex). While it has been noted before that there may be advantages to extending the domain of probabilities beyond the region between 0 and 1, most physicists are understandably reluctant to take such a step [21-24]. It is not coincidental that the situation here is analogous to the existence of imaginary momenta in tunneling. In that context, a quasiclassical approximation would suggest that the traversal time  $t = d/v = md/p$ , becoming imaginary as well; certain more careful calculations have also yielded complex values for the traversal time. It is natural to object that an imaginary number is unsatisfactory for describing a time; as Landauer has put it, "Has anyone seen a stop-watch with complex numbers on its dial?" In a similar vein, no one has rolled double-sixes an imaginary number of times. However, quantum mechanics is at bottom not a theory of probabilities, but rather one of wave functions. A wave function, even the wave function of the hand of a stopwatch, can be complex. It is only at the level of interpretation that these functions are turned into probabilities. In measurement theory, the justification for this is that an "ideal" measurement leads to complete decoherence between the various possible outcomes, leaving only the (real) diagonal elements of the density matrix to describe the system. But quantum mechanically, the state of a clock's hand may be  $\exp[(x - x_0)^2/4\tau^2]$ , where  $x_0$  need not be real. It is in this sense that a quantum-mechanical clock may indeed indicate a complex time. At the level of observation, if we choose to measure the expectation value  $\langle x \rangle$ , we will find a real number:  $\text{Re } x_0$ . If we wish to find  $\text{Im } x_0$ , we must instead look at the momentum of the clock hand, whose expectation value is  $\text{Im } x_0 = 2\tau^2$ .

We therefore suspend our disbelief in complex probabilities, and continue calculating the conditional probability distribution for the position of a tunneling particle,  $P(x; t_{\text{trans}})$ , where "trans" indicates that the particle is found on the far side of the barrier (assumed to extend from  $x = -d/2$  to  $x = +d/2$ ) at late times. From Eq. (1), we write this as

$$\begin{aligned} P(x; t_{\text{trans}}) &= \frac{P(x; t_{\text{trans}})}{P(\text{trans})} \\ &= \frac{\langle \psi_{\text{trans}} | P_{\text{trans}} | \psi_{\text{trans}} \rangle}{\langle \psi_{\text{trans}} | \psi_{\text{trans}} \rangle} \\ &= \frac{\langle \psi_{\text{trans}} | \hat{H} | \psi_{\text{trans}} \rangle}{\langle \psi_{\text{trans}} | \psi_{\text{trans}} \rangle} \\ &= \frac{\langle \psi_{\text{trans}} | \hat{H} | \psi_{\text{trans}} \rangle}{\langle \psi_{\text{trans}} | \psi_{\text{trans}} \rangle} \\ &= \frac{1}{T} \langle \psi_{\text{trans}} | \hat{H} | \psi_{\text{trans}} \rangle \end{aligned} \quad (4)$$

where  $\psi_{\text{trans}}$  is the initial state in which the particle is prepared (e.g., a packet incident from  $x = -1$  at  $t = -1$ ),  $\psi_{\text{trans}}$  is the state of a transmitted particle (i.e., at late times simply  $\psi_{\text{trans}}$  projected onto the region on the far side of the barrier), and  $T = \langle \psi_{\text{trans}} | \psi_{\text{trans}} \rangle$  is the transmission amplitude. For a symmetric barrier and symmetric initial conditions,  $\psi_{\text{trans}}$  is simply  $\psi_{\text{trans}}$  flipped about  $x = 0$  and about  $t = 0$ , since it is defined to consist of a packet heading towards  $x = +1$  as  $t \rightarrow +1$ , just as  $\psi_{\text{trans}}$  is defined to come entirely from  $x = -1$  as  $t \rightarrow -1$ . In more practical terms, we can define  $\psi_{\text{trans}}(x; t) = T \psi_{\text{trans}}(x; t) + R \psi_{\text{trans}}(x; t)$  (where  $R$  is the reflection amplitude), so that at late times  $\psi_{\text{trans}}(x = 0; t = 0) / T = T + R = 1$  and  $\psi_{\text{trans}}(x = 0; t = 0) / T = R + T = 0$ . Strictly speaking,

$$\psi_{\text{trans}}(x; t) = \int_{-\infty}^{\infty} \frac{dk}{2\pi} \phi(k) \psi_{\text{trans}}(x; t) \quad (5)$$

$\phi(k)$  is a bandwidth function, and the eigenstates

$$\psi_{\text{trans}}(x; t) = \begin{cases} e^{ikx} + R(k)e^{-ikx} e^{-i(k)t} & \text{for } x < -d/2 \\ T(k)e^{ikx} e^{-i(k)t} & \text{for } x > +d/2 \end{cases} \quad (6)$$

are orthogonal for different  $k$ . Now,

$$\begin{aligned} \psi_{\text{trans}}(x; t) &= \int_{-\infty}^{\infty} \frac{dk}{2\pi} \phi(k) [T(k) \psi_{\text{trans}}(x; t) + R(k) \psi_{\text{trans}}(x; t)]; \text{ thus} \\ \psi_{\text{trans}}(x < -\frac{d}{2}; t) &= \int_{-\infty}^{\infty} \frac{dk}{2\pi} \phi(k) [T(k)e^{ikx} + T(k)R(k)e^{ikx} + R(k)T(k)e^{ikx} e^{-i(k)t}] \\ &= \int_{-\infty}^{\infty} \frac{dk}{2\pi} \phi(k) T(k)e^{ikx} e^{-i(k)t} \\ &= \psi_{\text{trans}}(x; t) \text{ for real } t; \text{ and} \end{aligned} \quad (7)$$

$$\begin{aligned}
\psi(x > \frac{d}{2}; t) &= \int_{-\infty}^{\infty} \frac{dk}{2\pi} [T(k) e^{ikx} + R(k) e^{-ikx}] e^{-iE(k)t} \\
&= \int_{-\infty}^{\infty} \frac{dk}{2\pi} [T(k) e^{ikx} + R(k) e^{-ikx}] e^{-iE(k)t} = \frac{1}{2} \\
&= \psi(x; t) \text{ for real.}
\end{aligned} \tag{9}$$

(The assumption of  $\psi$ 's reality assures the time-symmetry of the incident state, yielding the parity relationship between the various states.) So long as  $\psi(k)$  is restricted to a bandwidth over which the transmission and reflection coefficients vary negligibly, we may approximate the latter as constants, and Eq. 9 does indeed correspond to the projection of  $\psi_i$  onto the far side of the barrier at late times (for a more careful discussion of the effects of finite bandwidth on the closely related Larmor clock, see [25]). It is also simple to confirm that

$$\langle \psi | j | \psi \rangle = \int_{-\infty}^{\infty} dk j^2 T(k) |T(k)|^2 \tag{10}$$

To find a "conditional expectation value" for an observable  $R$  in general, for a particle prepared in state  $i$  and later found to be in state  $f$ , we sum over  $R$ 's eigenvalue spectrum:

$$\langle R \rangle_{fi} = \sum_j P(R_j | f) r_j = \frac{\langle \psi_f | R | \psi_i \rangle}{\langle \psi_f | \psi_i \rangle} \tag{11}$$

where  $R_j$  and  $r_j$  are the eigenstates and corresponding eigenvalues of  $R$ . The question to be answered is whether this expression, which follows from blindly applying probability rules to quantum mechanical expressions which need not be real-valued, has any physical meaning. Indeed, Aharonov et al. have arrived at precisely this expression on entirely physical grounds [26,27]. They reexamined von Neumann's theory of measurement, considering a new limit which they termed "weak measurement." In this regime, the measuring device is prepared in a state which will disturb the system to be measured as little as possible. As a result of the uncertainty principle, each individual measurement of this type is extremely imprecise. Aharonov et al. asked nevertheless what the mean result would be for a large number of such measurements, if one only examined the measuring device on those occasions on which the system being studied was found to be in the desired final state  $f$  (after the period during which the measurement interaction was on).

They considered a measuring apparatus or "pointer" whose position and conjugate momentum we shall term  $Q$  and  $P$ . A measurement of a given observable  $R$  results from a time-dependent interaction

$$H_{int} = g(t) P \otimes R \tag{12}$$

since  $P$  is the generator of translations for the pointer, the mean position of the pointer after the interaction will have shifted by an amount proportional to the expectation value  $\langle R \rangle_i$ . In an ideal measurement, the relative shifts corresponding to different eigenvalues of  $R$  are large compared with the initial uncertainty in the pointer's position, and the resulting lack of overlap between the final states leads to the effective decoherence (or irreversible "collapse") between different eigenstates of  $R$ . In [26,27], this approach is modified in that the initial position of the pointer has a large uncertainty, so that the overlap between the pointer states which become entangled with the state of the particle remains close to unity, and hence that the measurement does not constitute a collapse. Seen another way, this means that the pointer momentum  $P$  may be very well-defined, and therefore need not impart an uncertain "kick" to the particle; the measurement is "weak" in that it disturbs the state of the particle as little as possible between the state preparation and the post-selection. If the initial state of the pointer is  $\exp[-Q^2/4\sigma^2]$ , then after the measurement it will, for suitably normalized  $g(t)$ , be in the state  $\exp[-(Q - \langle R \rangle_{fi})^2/4\sigma^2]$ . As discussed earlier, the real part of  $\langle R \rangle_{fi}$  corresponds to the mean shift in the pointer position, while the imaginary part constitutes a shift in the pointer momentum. This latter effect is a reflection of the back-action of a measurement on the particle. It obviously does have physical significance, but since it does not correspond to a spatial translation of the pointer, should not be thought of as part of the measurement outcome. Furthermore, unlike the spatial translation  $Q = \text{Re} \langle R \rangle_{fi}$ , this effect is sensitive to the initial state of the pointer:  $P = \text{Im} \langle R \rangle_{fi} = 2\sigma^2$ . As  $\sigma$  becomes large, the measurement becomes very weak, and the momentum shift of the pointer (like the back-action on the particle) vanishes, while the spatial shift remains constant.

It is already quite remarkable that the same expressions for these conditional probabilities or "weak values" arise from general probability arguments as well as from careful consideration of measurement interactions. These expressions have many other attractive properties which make it tempting to consider them "elements of reality." In

many respects, they behave in a more intuitive fashion than do wavefunctions themselves. For example, in [26] it is pointed out that if a "weak" measurement of an operator  $R + S$  is made on a particle after it is prepared in an eigenstate of  $R$  with eigenvalue  $r$  and before it is detected in an eigenstate of  $S$  with eigenvalue  $s$ , the result will be simply  $r + s$ . This holds whether or not  $R$  and  $S$  commute, and even if  $r + s$  is outside the eigenvalue spectrum of  $R + S$ ; hence such a simple rule could not be obeyed by standard quantum measurements (ones which are precise, or "strong," enough to disturb the time-evolution between  $r$  and  $s$ ). More generally, weak values are noncontextual and additive;  $\langle R + S \rangle_{f,i} = \langle R \rangle_{f,i} + \langle S \rangle_{f,i}$ . When averaged over an orthonormal set of final states, they reproduce the usual expectation value, since  $P(A) = \sum_f P(f)P(A|f)$ . These conditional probabilities are also easily shown to obey a chain rule,  $P(A \& B|f) = P(B|f)P(A|B)$ . In sum, there are many reasons to ascribe a certain level of reality to these conditional probability distributions.

In [28], it was argued that one could use this formalism to calculate the "dwell time" for transmitted or reflected particles individually. In the time-independent case treated in that paper, the dwell time is defined as the number density integrated over the barrier region (the expectation value of the projector onto the barrier region,  $\int_{d=2}^{d=2} (x + d=2) (x - d=2)$  for a barrier extending between  $d=2$ ) divided by the incident flux. It was shown for a rectangular barrier that these conditional dwell times were equal to the usual dwell time calculated for the ensemble as a whole, plus an imaginary part corresponding to the back-action of a measuring device on the tunneling particle. For the transmitted particles, however, the real part of this dwell involved equal contributions from regions near either edge of the barrier, while for reflected particles the dominant contribution came from the region near the entrance face only. Here I would like to expand on the utility of this approach to tunneling times, by presenting complete probability distributions for the position of a particle at a given time, conditioned upon reflection or transmission. These can be related back to the conditional dwell times by

$$Z_{f,i} = \int_{d=2}^{d=2} dt \int_{d=2}^{d=2} dx P(x; t|f); \quad (13)$$

where  $f$  indicates the final state under consideration and  $d=2$  are the edges of the barrier.

To understand the meaning of these conditional probability distributions, let us first consider an explicit gedanken-experiment. In [28], it was seen that the Larmor times [29][31] are simply one instance of a "weak measurement" of the dwell time. The two components of the Larmor time turn out (as in the closely related analysis by Sokolovskii et al. [24,32][35]; see also [36][39]) to be the real and imaginary parts of the conditional dwell time. By considering the effects of preparing the measuring device (in that case, the spin of the tunneling particle itself) in a state with great "pointer position" uncertainty, it was found that only the real part (representing the mean shift in pointer position, and neglecting the pointer momentum) had physical significance independent of the details of the measurement. However, the fact that both the "position" and the conjugate "momentum" of that pointer are simply spin angles makes the distinction a subtle one. In addition, that experiment fails to show how the conditional probability distributions for transmitted and reflected particles differ, measuring only the integrated dwell time, which is the same for both subensembles. Instead of considering spin, let us imagine a scenario as in Fig. 1 [40]. A heavy charged particle such as a proton is tunneling in one dimension. It passes through a series of parallel conducting plates with small holes. The plates may be held at a large positive voltage in order to form a tunnel barrier for the proton. (Due to the attraction between the proton and its image in each plate, it also experiences a periodic effective potential; in principle, if the plates were separated by one-half the proton's de Broglie wavelength, they would therefore form a "bandgap" for the proton even if no external potential were applied.) We will be concerned with the "opaque" limit, i.e., the case where the transmission probability is small because the evanescent decay constant  $\kappa = \sqrt{k_0^2 - k^2}$  (where  $k$  is the incident wavevector and  $\hbar^2 k_0^2 = 2mV$  is the height of the barrier) is much greater than the reciprocal of the barrier width. Between each pair of plates, but far from the proton's trajectory, is an electron constrained to move parallel to the plates. Due to the shielding of the proton's field by the plates, each electron only feels a significant Coulomb force while the proton is between the same pair of plates as that electron. Thus each electron serves as a test particle; if it begins at rest, its final momentum serves as a record of how long the proton spent in the region from which it is not shielded. Note that the electron momentum is thus the "pointer position," and the conjugate "pointer momentum" is the physical position of the electron. This follows from the form of the Coulomb interaction, which in the presence of the conducting plates can be written

$$H_{int} = -e^2 g(x_p) y_e; \quad (14)$$

which is approximately linear in electron transverse position  $y_e$  (so long as  $y_e \ll \hbar y_e$ ) and is proportional to a function of the proton's longitudinal position,  $g(x_p)$ , which is close to zero except in the region between the two plates. Let us repeat protons through this apparatus one by one, and first examine the electrons only on those occasions

when a proton is reflected. (The electrons are all "reset" between shots.) The momentum shift of the electron is now proportional to the "weak value" of the time spent by the proton in its region of sensitivity. In other words, it measures the time integral of the conditional probability that the electron was in its region, given that the proton was to be reflected:

$$p_e / \int_{-\infty}^{\infty} dt \int_{-\infty}^{\infty} dx g(x) P(x; t_{\text{refl}}); \quad (15)$$

where  $g(x)$  describes the force on the electron for a proton at position  $x$ , and is reasonably well confined to the region between the pair of plates surrounding the electron in question. The arrows in the figure indicate the nominal momentum of the electron. What we find is perhaps not unexpected. The proton's wave function decays exponentially inside the barrier, and so does  $P(x; t_{\text{refl}})$ ; only the electrons closest to the entrance face accumulate a significant momentum kick. Since nearly all the particles are reflected, this conditional probability is nearly the same as the unconditioned probability  $j(x)^2$ , and thus has a negligible imaginary part; none of the electrons undergoes a significant position shift aside from the time-dependent one due to their momenta.

But now what if we consider only events where the proton is transmitted? Here we find that  $P(x; t_{\text{trans}})$  is essentially an even function of  $x$ , as can be seen by examining Eq. (4) and recalling that  $t_i$  and  $t_r$  are related by a parity  $\hat{p}$  (along with a time-reversal). Using our more practical definitions of  $t_i$  and  $t_r$ , we have

$$\begin{aligned} P(x; t_{\text{trans}}) &= \frac{1}{T} t_i(x; t) i(x; t) = j_i(x; t)^2 + \frac{R}{T} i(-x; t) i(x; t) \\ P(x; t_{\text{refl}}) &= \frac{1}{R} r_i(x; t) i(x; t) = j_i(x; t)^2 + \frac{T}{R} i(-x; t) i(x; t); \end{aligned} \quad (16)$$

One can see clearly from this that  $\int j_i^2 P(x; t_{\text{trans}}) + \int j_i^2 P(x; t_{\text{refl}}) = \int j_i(x; t)^2$  as expected, which leads directly to the well-known relation between transmission, reflection, and (full-ensemble) dwell times  $\int j_i^2 T + \int j_i^2 R = d$ . One can also see that for  $R \gg T$  as in the opaque limit,  $P(x; t_{\text{refl}})$  is essentially equal to the absolute square of the incident wave function (decaying exponentially into the barrier), while  $P(x; t_{\text{trans}})$  is dominated by a term which is an even function of  $x$ . After more careful consideration, one also notes that since  $R$  and  $T$  are 90° out of phase [25,41,42] and since  $i_i$  is dominated in the barrier region by real exponential decay,  $P(x; t_{\text{trans}})$  is mostly imaginary. Only near the two extremes of the barrier, where the differing phases of the evanescent and anti-evanescent waves become important, does the real part become significant. As shown in [28],

$$\begin{aligned} T &= \frac{m}{\hbar k} \frac{1}{T} (B^2 + C^2)d + (BC + CB) \frac{\sinh d}{\cosh d} \\ R &= \frac{m}{\hbar k} \frac{1}{R} (BC + CB)d + (B^2 + C^2) \frac{\sinh d}{\cosh d} \\ d &= \frac{m}{\hbar k} (BC + CB)d + B^2 + C^2 \frac{\sinh d}{\cosh d}; \end{aligned} \quad (17)$$

where  $B$  and  $C$  are the coefficients of the evanescent and anti-evanescent waves, respectively, and satisfy  $B = C e^{-d}$ . In Fig. 1b, the effect of this is seen. The overall momentum transferred to the set of electrons is the same as in the case of a reflected proton, but it is now split evenly between the electrons within an exponential decay length of either edge of the barrier; when a particle is transmitted, it spends as much time by the exit face of the barrier as by the entrance face. The electrons in the center of the barrier still exhibit essentially no momentum shift! It is as though the proton simply "hopped" from one edge to the other, spending negligible time in the barrier. This is related to the well-known fact [9] that the wave packet delay time in opaque tunneling is independent of the barrier thickness. However, we must recall that the imaginary part of  $P(x; t_{\text{trans}})$  is significant over the entire barrier. This manifests itself as a shift in the mean position of all the electrons (which we have been describing as the pointer "momentum," and which is never affected by "ideal" quantum measurements). This is simple to understand; the electrons have some uncertainty in their position to begin with. Due to the attractive Coulomb interaction, the closer a given electron is to the proton, the smaller the potential barrier the proton has to traverse. Thus by selecting protons which succeeded in tunneling, we are post-selecting states where the electrons were nearby to begin with. The constancy of this back-action across the length of the barrier reflects the fact that within the WKB approximation, the transmission is  $\exp[-\int_{-\infty}^{\infty} g(x) dx]$ , i.e., equally sensitive to a change in the potential at any point in the barrier. Unlike the momentum kick (the measurement outcome itself) this effect is entirely dependent on the initial uncertainty in the electron position. If the initial states of the electron are very well localized in space, they are hardly shifted at all

by this effect. On the other hand, the uncertainty principle then requires their initial momenta to be poorly defined, "weakening" the resolution of the measurement. Over many trials, however, the mean value of the momentum shift will be unaffected by the choice of initial uncertainty.

As discussed in [28], this pointer position shift corresponds to the in-plane portion of the Lambor time, while the pointer momentum shift corresponds to minus the out-of-plane portion [31,32]. Both time scales are meaningful, but their meanings are distinct. (Furthermore, the suggestion that one should pay attention to  $j_T j$  rather than to  $\tau$  itself seems odd, if only because unlike the real and imaginary parts individually, this fails to satisfy the stipulation that the dwell in a large region be equal to the sum of the dwell in a set of smaller regions which make it up.) The real part of  $\tau$  indicates the magnitude of the effect our tunneling particle would have on test particles. It also describes the amount of absorption or gain the tunneling particle(s) would suffer if tunneling through an active medium. For  $k > 0$ , this time also approaches the group delay time for the peak of the tunneling packet. On the other hand, the imaginary part indicates the magnitude of the back-action on the tunneling particle due to the measurement. This is the timescale which emerges from consideration of the effects of an oscillating barrier on a tunneling particle, for example [7,43]. Since unlike the real part, it grows proportionally with barrier thickness, it dominates in the opaque limit, where it reduces to  $m d = \hbar$ . (It is interesting to note that to good approximation, the "weak value" thus follows what one might expect in the WKB limit, and yields  $\hbar \tau_i = m d = \hbar \pi$ , even when  $\hbar \pi$  is imaginary; this is another example where the behavior of weak values obeys simple rules even in regimes where we expect these rules to break down; see also [44].) Importantly, the back-action due to this imaginary part depends strongly on the initial state of the measuring device; in this sense, it is not a characteristic of the tunneling particle itself. Of course, in real devices, there will be an interplay between these two timescales. The tunneling particle will affect nearby particles, which may in turn modify the tunneling characteristics. The former effect, however, can now be seen to be limited to the extremities of the barrier region, while the latter can be arbitrarily small if the nearby particles are arranged in such a way that they do not create a large uncertainty in the potential seen by the tunneling particle.

In Figs 2 and 3, we can observe the time-evolution of the conditional probability distributions. At early times, the distribution is real, and mimics the initial wave packet; at late times, it is also real, and mimics the final wave packet. This seems to be in contrast to the claim that in order to reconcile superluminal peak propagation with causality, we must consider all the transmitted particles to have originated near the leading edge of the incident wave packet [12,45,46]. Figure 2 presents the real and imaginary parts of  $P(x|trans)$  for a wavepacket incident from the left on a barrier which extends from  $x = -5$  to  $x = +5$ . Also shown is the full-ensemble distribution  $P(x) = \sum_j P(x|j)$ . The wave packets are constructed according to Eq. 5, with a real Gaussian bandwidth function whose parameters are given in the figure captions. For the parameters chosen, the transmissivity is about  $5.5 \times 10^{-6}$ , so  $P(x|refl)$  would be essentially indistinguishable from  $P(x)$ . As advertised, the real part of  $P(x|trans)$  is exponentially suppressed near the center of the barrier, while the imaginary part becomes large and essentially constant across the barrier region during the tunneling event. Oscillations in both parts are seen to either side of the barrier, due to the self-interference of the wave packet near the potential step. As discussed in [28], these extra-barrier oscillations average out to zero when integrated over space; the total time spent by a particle in a long region  $L$  to either side of the barrier is simply  $m L = \hbar k$ . The regions of negative conditional probability have a clear physical meaning; in the experiment of Figure 1, for example, they predict that a test electron would experience a repulsive momentum transfer rather than an attractive one due to the proton's Coulomb field. Similar predictions apply to any other von Neumann-style measurement one might contemplate. For this reason, although we sympathize with those who consider negative times "unphysical," we do not see a better definition than that provided by measurement outcomes; to deny this definition, one would either have to give up the expression for the Coulomb force or the idea that momentum transfer is the force integrated over the interaction time. From the figures, we can also see that as the wavepacket overcomes the barrier, the peaks in  $P(x|trans)$  decrease in magnitude on the left side of the barrier, and grow on the right side, without ever traversing the center. This is what was meant after Eq. 17 by the particle "hopping" across the barrier. It is in a sense "nonlocality of a single particle." While it is well known that despite the nonlocality inherent in quantum mechanics, no expectation values can ever depend on choices made at spacelike separated points (and thus that nonlocality does not violate Einstein causality), this shows that a single particle can affect expectation values of (weak) measuring devices at two spacelike separated positions (see related discussions of the "reality of the wave function" [47{49}]).

It is worth asking whether the "duration" of the tunneling process is best defined by Eq. (13), or rather by the length of time over which  $P(x^2 | d=2; d=2; t|trans)$  remains close to its maximum value. As can be seen from the figures and from Eq. (4), the latter quantity is simply the length of time during which the wave function is significant at the location of the barrier. Since in tunneling, the group delay time for traversal is small compared with the temporal width of the freely propagating wave packet width (the contrary would imply  $\tau_g = 2m = \hbar k > (1/k) = (\hbar k = m)$ , or  $k > 0$ , leading to  $(k + k)^2 > k^2 + k^2 > k^2 + 0 = k_0^2$ , i.e., a significant portion of the wave packet having

enough energy to traverse the barrier without tunneling), this time scale is dominated by the duration of the wave packet, approximately  $m = \hbar k$ . It should be borne in mind that the same would be true for free propagation. If a wave packet of length  $L$  traverses an empty region of width  $d < L$ , the length of time during which a change in the potential in the region could significantly affect the transmission probability is  $L/v$ ; nonetheless, for most intents and purposes we would think of an individual particle as spending only  $d/v$  in that region. Obviously, the relevant time scale will depend on the precise experimental question one wishes to address.

Figure 3 shows the case for a particle which is incident with enough energy to traverse the barrier. The reflection probability is about 3.5%, and both  $P(x)_{\text{trans}}$  and  $P(x)_{\text{refl}}$  are displayed, along with  $P(x)$ . The latter is generally obscured by  $P(x)_{\text{trans}}$ , but in those regions where it can be seen, it is clearly the weighted average of the reflection and transmission distributions, as expected. In this figure, we find the transmitted portion traversing the barrier relatively smoothly, albeit with some oscillations. It has a very small imaginary part (as follows for  $\mathcal{J} = R_j - 5$ ). The reflected portion, on the other hand, undergoes violent oscillations in both its real and imaginary parts. At late times, these die away, and a real wave packet propagating back to the left is all that remains.

One might think that the real and imaginary parts of this weak value, one part describing the effect of a particle on a measuring device and the other describing the back-action of the measuring device on the particle, should be equal in the "classical" limit, that is, for allowed transmission. Inspecting the figures, however, one sees that in the region of allowed propagation, the imaginary part tends to zero. This can be understood as follows: an infinitesimal perturbation in the potential leads to a reflection amplitude proportional to the perturbation. If prior to the perturbation, the reactivity was zero, the reflection probability is quadratic in the perturbation and hence vanishes in the limit of a gentle measurement. If on the other hand, there is a non-zero reactivity prior to the perturbation, then the reflection probability  $j_0 + r_j^2$  grows linearly with the perturbation, indicating a finite back-action.

As discussed in [28], another interesting feature of these weak values is that  $P(x > d = 2j_{\text{refl}})$  need not vanish. This implies that a particle incident from the left and ultimately reflected does spend some time to the right of the barrier, at least within one wave packet width of the exit face. Although this can be understood as an effect due to the possibility of coherent reflection off the measuring device itself, the fact that the calculated time is independent of the type or strength of the measurement interaction may be taken to ascribe a certain level of reality to this time regardless of how or even whether it is observed. The argument is essentially the same one made in favor of negative times. It should be noted that as the strength of the measurement interaction is lowered, the probability of a particle being reflected by the measuring device falls as the square of the potential, while the "pointer position" shift is linear in the interaction strength.

Although it is always possible to argue about definitions of words like "interaction time," the advantage of the weak measurement approach is that it offers a simple and intuitive formalism to treat a broad variety of experimental predictions in a unified manner. While some will not want to call a negative or a complex number a "time," the words are relatively unimportant; the features shown in Figures 3 and 4 are in principle experimentally testable. In particular, the result shown schematically in Fig. 1, that transmitted particles have equal effects near both sides of a barrier, while reflected particles affect only the region around the entrance face, is to my knowledge a new prediction. The gedankenexperiment of that figure is of course unrealistic, but it is conceivable that in the future an analogous experiment could be performed. For instance, one might consider Rydberg atoms travelling through two or three successive micromasers and leaving partial information in the stored fields.

At the present time, there is an indirect test which should be feasible and in fact rather straightforward. As demonstrated in [14,20], a multilayer dielectric mirror possesses a photonic bandgap [50] and may be used as a tunnel barrier. These mirrors are made of alternating high- and low-index quarter-wave layers. Typically, each dielectric layer has very low loss, and the low transmissivity is due to reflection rather than to absorption. If one layer is now doped with an absorbing material, in general both the transmission and the reflection should decrease; the amount of the decrease can be thought of as an indication of the time spent by the transmitted or reflected photons in the layer in question. Figure 4 contains numerical results for an 11-layer mirror. The ratio of the transmission (and reflection) in the presence of one absorbing layer to the transmission (and reflection) in the absence of absorption was calculated. The logarithm of this ratio (measured in units of the single-layer attenuation, so that in semiclassical terms what is being plotted might be thought of as the number of passes through the layer) is plotted versus the position of the lossy layer. Fig. 4(a) is for light incident in the center of the bandgap, with transmission of about 1.2% (in the absence of any absorber). The similarity of these curves to those of Fig. 2 should be evident. In Fig. 4(b), the light is incident outside the bandgap, near the first resonant transmission point (where the entire mirror may be thought of as a single Kronig-Penney crystal, and multiple reflections between the opposite edges of the mirror interfere constructively, leading to near-100% transmission). In this regime, the mirror is essentially a low-Q Fabry-Perot operating in its fundamental mode, and the transmission is most sensitive to an absorber near the center of the structure, where the mode has its maximum. The reflection is close to zero in the absence of an absorber, so the addition of absorption

can only lead to effective gain in the reflection channel; this is the meaning of the negative value of the curve. In Fig. 4(c), the light is still outside the bandgap, but now at a transmission minimum of about 66%. One can again observe the mode structure within the barrier, and complementary oscillations for the transmitted and reflected parts, reminiscent of Fig. 3.

The effect of absorption offers another way of understanding the meaning of complex times. If each quarter-wave layer has an amplitude transmissivity of  $1/\sqrt{2}$ , we may think of this as an amplitude of  $\exp[-\sqrt{2}/2]t$  to survive a time  $t$  spent in the barrier. In the tunneling regime, the time becomes predominantly imaginary;  $t \rightarrow i t_{BL}$ , where  $t_{BL}$  is the Buttiker-Landauer time ( $m d = \hbar$  for a massive particle). Thus the attenuation factor becomes a phase shift rather than absorption. This is connected to the weak measurement idea in that if absorption is used as a clock, the "pointer" is essentially the photon number. The conjugate "momentum" is hence the optical phase. Yet another way of understanding this is to recall that the imaginary part of the "conditional" dwell time, like the out-of-plane Larmor rotation, can be related to an energy-derivative of the magnitude of the transmission amplitude, while the real part, like the usual dwell time, the group delay, and the in-plane Larmor rotation, can be related to an energy-derivative of the phase of the transmission amplitude [31,28]. Since absorption (or gain, which will behave in precisely the same way) can be expressed as an imaginary contribution to an effective Hamiltonian, we see by analytic continuation that the real part of the conditional dwell time describes the amount of attenuation caused by an absorber, and the imaginary part describes the absorber's effect on the phase. An interesting corollary to this is that loss within the tunnel barrier may increase the physical delay (cf. [51]{54}) by introducing a frequency-dependent phase shift, in principle without a large impact on the transmission and reflection probabilities themselves. Indeed, in [14], the measured delay time exceeded the group delay prediction by about 0.4 fs, but this was at the borderline of statistical significance. More recently, we have found this discrepancy to be statistically significant and to persist for two different barriers of identical design [20]. It is possible that a portion of this effect could be due to absorption or scattering in the dielectric layers at the several percent level, although further work is necessary to see if such a model can be tailored to agree with the observed reflection characteristics.

The present results may also be of use in constructing high-reactivity dielectric mirrors, whose characteristics are ultimately limited by the (small) losses in the dielectrics. It is intuitively clear that the reactivity is sensitive mostly to losses near the entrance face of the mirror, but one might have expected the transmissivity to be equally sensitive to losses anywhere in the structure. The present results show instead that losses near the middle of the mirror can be tolerated without having a significant impact on the transmission, aside from introducing an additional time delay. (It is interesting to note that gain, on the other hand, could shorten the time delay without having a significant effect on the transmission probability; cf. [55]{58}.) If a "smooth" barrier were constructed, so that the WKB approximation held, the conditional dwell time would become pure imaginary between the classical turning points, and the transmission would (in the limit) become completely insensitive to loss in this region.

In conclusion, we have shown that a straightforward definition of conditional probabilities in quantum theory, equivalent to Aharonov et al.'s idea of "weak measurements," allows one to discuss the history of a particle which has tunneled. This approach makes clear the meaning of imaginary and complex times, and can describe the results of a broad range of hypothetical experiments. It is found that tunneling particles spend equal amounts of time near the entrance and exit faces of the barrier, but vanishingly little in the center; by contrast, reflected particles spend most of their time near the entrance face only. One "clock" considered is that of absorption in some region within the barrier. The anomalously small dwell time is shown to lead to anomalously small losses, while the imaginary part of the "conditional" dwell time may lead to an additional time delay in the presence of absorption.

This work was supported by the U.S. Office of Naval Research under grant N00014-90-J-1259. I would like to thank R.Y. Chiao for years of discussions without which this work would have been either impossible or completed much sooner. I would also like to acknowledge many useful comments made by R. Landauer, M. Mitchell, G. Kurizki and Y. Japha.



## Figure Captions

1. A gedankenexperiment using distant electrons to measure how much time a tunneling proton spends in each of several shielded regions of space. The proton is tunneling along  $x$ , through a series of plates held at some repulsive potential  $V$ . While the proton is between a given pair of plates, only the electron between the same pair of plates feels a significant Coulomb attraction. After the proton has stopped interacting with the barrier region, the momentum of the electrons serves as a record of how much time the proton has spent in each sub-region. If we examine the electrons only after detecting a reflected proton or only after detecting a transmitted proton, we may observe the time-integrated conditional probability distributions discussed in the text. (If time-dependent shutters were added before the electrons, the distributions could even be measured as a function of time.) In (a) we see the final state of the electrons for cases where the proton is reflected: only those in the first evanescent decay length of the tunnel barrier feel the proton's Coulomb potential and acquire a consequent momentum kick, indicated by arrows. In (b) we see what happens if the proton is transmitted: electrons near both edges of the barrier acquire a momentum kick, but those near the center do not. The tunneling proton seems not to have spent any time in the center of the barrier. However, the position of the electrons gets shifted when the proton is transmitted, as indicated by the redrawn electron wave packets. As explained in the text, this is indicative of the back-action of the measuring electrons on the tunneling particle.

2. The heavy solid curves show the real part of the "conditional probability" for a tunneling particle to be at position  $x$ , for several different times. This represents the force which an electron at a given  $x$  would experience in the example of Figure 1. The dashed curve represents the back-action, i.e., the position shift an electron would experience. For comparison, the light solid curve shows the probability distribution for the ensemble as a whole, without distinguishing between transmitted and reflected particles. With  $\hbar = m = 1$ , the parameters for these curves are incident wave vector (and hence initial velocity)  $k = 0.5$ , barrier height in wave vector units  $k_0 = 0.75$ , rms wave vector uncertainty  $\Delta k = 0.03$ , and barrier thickness  $d = 10$  (centered at  $x = 0$ ). The transmission probability is thus  $5.5 \times 10^{-5}$ , and the group delay for both transmission and reflection is  $7.1$ , making the effective traversal speed about three times the free propagation velocity. For thick enough barriers, this effective velocity could even exceed the speed of light. Note that the real part of the conditional probability distribution mimics the incident wave packet at early times and the transmitted packet at late times, thus travelling anomalously fast as well; it appears to do this by "skipping" the region in the middle of the barrier entirely (although while tunneling, a large imaginary part builds up in that region). Despite the fact that for certain parameters, the emerging peak may only be causally connected to the leading edge of the incident packet, the (measurable) conditional probabilities as defined here do not support the identification of the transmitted particles with the the leading edge of the incident packet.


3. Same as Figure 2, but for allowed transmission. In addition, the light dashed and dotted curves show the conditional probability distribution for reflected particles. The parameters are  $k = 0.75$ ,  $k_0 = 0.5$ ,  $\Delta k = 0.02$ , and  $d = 10$ , for a reflection probability of  $0.035$  and a group delay of  $18.2$ . In this case, aside from self-interference terms, the transmitted-particle distribution traverses the barrier relatively smoothly, while the reflected-particle distribution displays rapid oscillations in and to both sides of the barrier region, only reconstructing a smooth reflected peak when the wave packet has left the barrier region.

4. Numerical calculation of the effect of introducing a 5% amplitude loss into one of the 11 dielectric layers of the mirror studied in [14]. As explained in the text, the consequent attenuation of the transmitted and reflected beams may be thought of as a measure of the time spent in the layer in question by transmitted and reflected particles, respectively. The logarithm of this attenuation is plotted against the position of the lossy layer. In (a), the incident light is near the center of the bandgap. In (b), it is near the first resonant transmission point,  $k=k_0 + 1.2$ . In (c), it is still in a regime of allowed transmission, but with Fabry-Pérot-like interference lowering the transmission closer to 66%. Here  $k=k_0 + 1.3$ .


- 
- [1] E. U. Condon, Rev. Mod. Phys. 3, 43 (1931); see pp 74 .
  - [2] L. A. McColl, Phys. Rev. 40, 621 (1932).
  - [3] L. Eisenbud, Ph.D. Thesis, Princeton University (1948).
  - [4] E. P. Wigner, Phys. Rev. 98, 145 (1955).
  - [5] T. E. Hartman, J. App. Phys. 33, 3427 (1962).
  - [6] K. W. H. Stevens, Eur. J. Phys. 1, 98 (1980).
  - [7] M. Buttiker and R. Landauer, Phys. Rev. Lett. 49, 1739 (1982).
  - [8] S. Collins, D. Lowe, and J. R. Barker, J. Phys. C 20, 6213 (1987).
  - [9] E. H. Hauge and J. A. Stvneng, Rev. Mod. Phys. 61, 917 (1989).
  - [10] M. Buttiker, In \Electronic Properties of Multilayers and Low Dimensional Semiconductors," J. M. Chamberlain, L. Eaves, and J. C. Portal, eds., Plenum, New York (1990).
  - [11] V. S. Okhovsky and E. Recami, Phys. Rep. 214, 339 (1992).
  - [12] C. R. Leavens and G. C. Aers. In \Scanning Tunneling Microscopy III," R. Wiesendanger and H.-J. Guntherodt, eds., Springer-Verlag, Berlin (1993).
  - [13] R. Landauer and Th. Martin, Rev. Mod. Phys. 66, 217 (1994).
  - [14] A. M. Steinberg, P. G. Kwiat, and R. Y. Chiao, Phys. Rev. Lett. 71, 708 (1993).
  - [15] A. Enders and G. Nimtz, J. Phys. I France 3, 1089 (1993).
  - [16] A. Ranfagni, P. Fabeni, G. P. Pazzi, and D. Mugnai, Phys. Rev. E 48, 1453 (1993).
  - [17] Ch. Spielmann, R. Szpoc, A. Stingl, and F. Krausz, Phys. Rev. Lett. 73, 2308 (1994).
  - [18] G. Nimtz, A. Enders, and H. Spieker, J. Phys. I France 4, 565 (1994).
  - [19] A. M. Steinberg, J. Phys I France 4, 1813 (1994).
  - [20] A. M. Steinberg and R. Y. Chiao, \Sub-femtosecond determination of transmission delay times for a dielectric mirror (photonic bandgap) as a function of angle of incidence," preprint quant-ph/9501013, to appear in Phys. Rev. A (1995).
  - [21] W. Mueckenheim et al., Phys. Rep. 133, 337 (1986).
  - [22] S. Youssef, Mod. Phys. Lett. 6, 225 (1991).
  - [23] M. O. Scully, H. Walther, and W. Schleich, Phys. Rev. A 49, 1562 (1994).
  - [24] D. Sokolovski and J. N. L. Connor, Phys. Rev. A 44, 1500 (1991).
  - [25] J. P. Falck and E. H. Hauge, Phys. Rev. B 38, 3287 (1988).
  - [26] Y. Aharonov and L. Vaidman, Phys. Rev. A 41, 11 (1990).
  - [27] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. 60, 1351 (1988).
  - [28] A. M. Steinberg, \How much time does a tunneling particle spend in the barrier region?," preprint quant-ph/9501015, to appear in Phys. Rev. Lett. (1995).
  - [29] A. I. Baz', Sov. J. Nucl. Phys. 5, 161 (1967).
  - [30] V. F. Rybachenko, Sov. J. Nucl. Phys. 5, 635 (1967).
  - [31] M. Buttiker, Phys. Rev. B 27, 6178 (1983).
  - [32] D. Sokolovski and L. M. Baskin, Phys. Rev. A 36, 4604 (1987).
  - [33] D. Sokolovski and J. N. L. Connor, Phys. Rev. A 42, 6512 (1990).
  - [34] D. Sokolovski and J. N. L. Connor, Phys. Rev. A 47, 4677 (1993).
  - [35] D. Sokolovski and J. N. L. Connor, Sol. St. Comm. 89, 475 (1994).
  - [36] H. A. Fertig, Phys. Rev. Lett. 65, 2321 (1990).
  - [37] H. A. Fertig, Phys. Rev. B 47, 1346 (1993).
  - [38] J. G. Muga, S. Brouard, and R. Sala, Phys. Lett. A 167, 24 (1992).
  - [39] S. Brouard, R. Sala, and J. G. Muga, Phys. Rev. A 49, 4312 (1994).
  - [40] A. M. Steinberg, \Tunneling times and weak measurements," to appear in Ann. N.Y. Acad. Sci., proceedings of the conference on Fundamental Problems in Quantum Theory (June, 1994), D. Greenberger et al., eds.
  - [41] Z. Y. Ou and L. Mandel, Am. J. Phys. 57, 66 (1989).
  - [42] A. M. Steinberg and R. Y. Chiao, Phys. Rev. A 49, 3283 (1994).
  - [43] T. Martin and R. Landauer, Phys. Rev. A 47, 2023 (1993).
  - [44] Y. Aharonov, S. Popescu, D. Rohrlich, and L. Vaidman, Phys. Rev. A 48, 4084 (1993).
  - [45] J. M. Deutsch and F. E. Low, Ann. Phys. 228, 184 (1993).
  - [46] A. M. Steinberg, P. G. Kwiat, and R. Y. Chiao, Found. Phys. Lett. 7, 223 (1994).
  - [47] Y. Aharonov and L. Vaidman, Phys. Lett. A 178, 38 (1993).
  - [48] Y. Aharonov, J. Anandan and L. Vaidman, Phys. Rev. A 47, 4616 (1993).
  - [49] W. Guhr, \The reality and measurement of the wave function," preprint hep-th/9308061, Phys. Rev. A 50, 882 (1994).
  - [50] E. Yablonovitch, J. Opt. Soc. Am. B 10 (2), 283 (1993), and references therein, and other articles in this special issue on photonic bandgaps.

- [51] A. Ranfagni, D. Mugnai, P. Fabeni, and G. P. Pazzi, *Physica Scripta* 42, 508 (1990).
- [52] D. Mugnai, A. Ranfagni, R. Ruggeri, and A. Agresti, *Phys. Rev. Lett.* 68, 259 (1992).
- [53] F. Raciti and G. Salesi, *J. Phys. I France* 4, 1783 (1994).
- [54] G. Nintz, H. Spieker, and H. M. Brodowsky, *J. Phys. I France* 4, 1379 (1994).
- [55] R. Y. Chiao, *Phys. Rev. A* 48, R34 (1993).
- [56] E. L. Bolda, J. C. Garrison, and R. Y. Chiao, *Phys. Rev. A* 49, 2938 (1994).
- [57] A. M. Steinberg and R. Y. Chiao, *Phys. Rev. A* 49, 2071 (1994).
- [58] R. Y. Chiao, *Quantum Optics* 6, 359 (1994).


time = -70.000000.




time = -50.000000.




time = -30.000000.




time = -10.000000.




time = 10.000000.




time = 30.000000.




time = 50.000000.




time = 70.000000.




time = -80.000000.




time = -60.000000.




time = -40.000000.




time = -20.000000.




time = 0.000000.




time = 20.000000.



time = 40.000000.



time = 60.000000.



time = 80.000000.

