

# Husimi Parametric Oscillator in Frame of Symplectic Group and Q-oscillators

V. I. Man'ko

Lebedev Physics Institute, Leninsky pr., 53  
Moscow 117924, Russia

## Abstract

Time-dependent integrals of motion which are linear forms in position and momentum are discussed for Husimi parametric forced oscillator. Generalization of these integrals of motion for q-oscillator is presented. Squeezing and quadrature correlation phenomena are discussed on the base of Schrodinger uncertainty relation. The properties of the generalized correlated states, squeezed states, even and odd coherent states (the Schrodinger cat states) are reviewed. The relation of the constructed nonclassical states to representations of the symplectic symmetry group and finite symmetry groups is discussed.

## 1 Q-invariant of Parametric Oscillator

Husimi [1] has considered forced parametric oscillator described by the Schrodinger equation

$$i\hbar \frac{\partial \psi(x;t)}{\partial t} = \frac{\hbar^2}{2m} \frac{\partial^2 \psi(x;t)}{\partial x^2} + \frac{1}{2} \omega^2(t) x^2 \psi(x;t) - f(t) x \psi(x;t); \quad (1)$$

where we take  $\hbar = m = \omega(0) = 1$ . In the references the Gaussian packets and analogs of oscillator number states have been found as well as Green function and transition amplitudes between the energy levels in a form of series. In [2], [3] the integrals of motion which are linear forms in position and momentum operators have been found and used to rederive the Husimi solutions. The review of the approach has been given in [4], [5].

In this talk for the Husimi parametric oscillator we find integrals of motion which satisfy the commutation relations of q-oscillator [6], [7], i.e. the quantum Heisenberg (Weyl) group is shown to be the dynamical group of this oscillator. In [8], [9] the possibilities to interpret the q-oscillator as nonlinear oscillator with specific exponential dependence of its frequency on the amplitude has been suggested. Using the construction of even and odd coherent states [10] interpreted in quantum optics as Schrodinger cat states [11] we discuss the properties of these states for the Husimi parametric oscillator.

For simplicity of presentation we take  $f(t) = 0$ . Then the time-dependent operator

$$A = \frac{i}{2} [\mu(t)p - \nu(t)x]; \quad (2)$$

where

$$\mu(t) + \nu^2(t)\mu(t) = 0; \quad \mu(0) = 1; \quad \nu(0) = i \quad (3)$$

is the integral of motion [3] satisfying the commutation relation

$$[A, Ay] = 1; \quad (4)$$

One can easily check that the time-dependent operator

$$A_q = A \exp \left( \frac{\sinh^{-1} A y A}{A y A \sinh} \right); \quad (5)$$

where  $\ln q$ , is the integral of motion of the parametric oscillator satisfying the commutation relation

$$A_q A_q^\dagger - A_q^\dagger A_q = q^{-1} A_q^\dagger A_q : \quad (6)$$

It means that operators  $A_q$  ( $A_q^\dagger$ ) corresponding to  $q$ -oscillators due to these commutation relations determine the dynamical symmetry of the parametric oscillator and all the states of the oscillator are the basis of the irreducible representation of  $q$ -deformed Heisenberg-Weyl group.

## 2 Correlated and Squeezed State

It was shown in [3] that some Husimi packet solutions may be interpreted as coherent states [12] since they are eigenstates of the operator  $A$  (2) of the form

$$(\mathbf{x}; t) = \psi_0(\mathbf{x}; t) \exp \left\{ -\frac{j^2}{2} - \frac{\omega^2(t)}{2\omega(t)} + \frac{p}{2} \frac{x}{\omega} g \right\}; \quad (7)$$

where

$$\psi_0(\mathbf{x}; t) = \exp \left\{ -\frac{1}{4} \omega(t) - \frac{1}{2} \exp \frac{i\omega(t)x^2}{2\omega(t)} \right\} \quad (8)$$

is analog of the ground state of the oscillator and  $\omega$  is a complex number. The variances of the position and momentum of the parametric oscillator in the state (8) are

$$\sigma_x = \frac{j^2(t)}{2}; \quad \sigma_p = \frac{j^2(t)}{2}; \quad (9)$$

and the correlation coefficient of the position and momentum has the value corresponding to minimization of the Schrodinger uncertainty relation [13]

$$\sigma_x \sigma_p = \frac{1}{4} \frac{1}{\omega^2}; \quad (10)$$

Due to this the state (7) is a correlated coherent state introduced in [14]. For different dependences of frequency on time there may appear phenomenon of squeezing, i. e.

$$\sigma_x < 1/2; \quad (11)$$

The orthonormal number states of parametric oscillator

$$\psi_n(\mathbf{x}; t) = \psi_0(\mathbf{x}; t) \frac{1}{n!} \left( \frac{\omega}{2} \right)^{n/2} H_n \left( \frac{x}{j(t)} \right) \quad (12)$$

are eigenstates of the integral of motion  $A_q$ . For the Husimi oscillator the  $q$ -coherent states [6] may be easily constructed as normalized eigenstates of the integral of the motion  $A_q$  (5) in the form

$$\psi^{(q)} = N_q \sum_{n=0}^{\infty} \frac{q^{n^2}}{[n]!} \psi_n(x;t); \quad (13)$$

where  $[n]! = [n][n-1]\dots[1]$ ;  $[0]! = 1$  and  $[n] = \sinh^{-1} n = \sinh^{-1} \frac{1}{q}$ . The normalization constant is

$$N_q = \left( \sum_{n=0}^{\infty} \frac{q^{n^2}}{[n]!} \right)^{-1/2} \quad (14)$$

The function (13) satisfies the Schrodinger equation (1), and it is eigenfunction of the  $q$ -oscillator operator  $A_q$ . In [8], [9] it was shown that if the  $q$ -oscillator describes a nonlinearity of classical electromagnetic field vibrations ( $q$ -nonlinearity) at high amplitudes the blue shift effect must exist. It is the effect of shifting the light frequency versus the light intensity which may serve for evaluating an upper limit for the parameter of nonlinearity  $q$ :

### 3 Squeezed Schrodinger Cats

Another normalized solution to the Schrodinger equation (1)

$$\psi_m(x;t) = 2N_m \psi_0(x;t) \exp \left\{ -\frac{j^2}{2} - \frac{\omega(t)^2}{2\omega(t)} g \cosh \frac{p}{2} x \right\}; \quad (15)$$

where

$$N_m = \frac{\exp(j^2/2)}{2 \cosh j^2} \quad (16)$$

is the even coherent state [10] (the Schrodinger cat male state). The odd coherent state of the parametric oscillator (Schrodinger cat female state)

$$\psi_f(x;t) = 2N_f \psi_0(x;t) \exp \left\{ -\frac{j^2}{2} - \frac{\omega(t)^2}{2\omega(t)} g \sinh \frac{p}{2} x \right\}; \quad (17)$$

where

$$N_f = \frac{\exp(j^2/2)}{2 \sinh j^2} \quad (18)$$

satisfies the Schrodinger equation (1) and is the eigenstate of the integral of motion  $A^2$  (as well as the even coherent state) with the eigenvalue  $-j^2$ . The even and odd coherent states are orthogonal to each other. A multimode generalization of the even and odd coherent states has been introduced in [15]. The properties of multimode Schrodinger cat states, the coherent components of which are subjected to parametric excitation, are discussed in [16]. The even and odd coherent states are the partial case of a set of nonclassical states discussed in [17]. They may be used as alternative to squeezed vacuum states in gravitational wave antenna to improve its sensitivity [18].

The multimode generalization of the Husimi parametric oscillator has been considered in [19]. The linear in position and momentum integrals of motion have been constructed

and generalized correlated states [20] in the form of Gaussian packets have been found explicitly. It is interesting to note that for multidimensional parametric oscillator the linear integrals of motion of the type A (2) and their quadratic forms constitute the Lie algebra of the inhomogeneous symplectic group. Thus, the solutions to Schrodinger equation for the multimode parametric oscillator form the basis for irreducible representation of this symplectic group. The remark made for one-mode case on the existence of q{deformed dynamical algebra may be extended to the multimode parametric oscillator, too. Thus, the solutions of the Schrodinger equation for multimode parametric oscillator may be also considered as a basis of the irreducible representation for a q{deformed symplectic group.

The construction of the even and odd coherent states made in [10] used the representation of the inversion group for which even states realize the symmetric irreducible representation and odd states realize the antisymmetric one. Generalization of this construction to higher crystallographic groups for introducing the crystallized Schrodinger cat states is done in [21].

## 4 Multivariable Hermite Polynomials

For Husimi parametric forced oscillator the transition amplitude between its energy levels has been calculated as overlap integral of two generic Hermite polynomials with a Gaussian function (Frank-Condon factor) and expressed in terms of Hermite polynomials of two variables [3]. For N-mode parametric oscillator the analogous amplitude has been expressed in terms of Hermite polynomials of 2N variables, i. e. the overlap integral of two generic Hermite polynomials of N variables with a Gaussian function (Frank-Condon factor for a polyatomic molecule) has been evaluated in [19]. The corresponding result uses the formula

$$\int_{-\infty}^{\infty} H_n^{fRg} H_m^{fRg} (x + d) \exp(-x^T M x + c^T x) dx = \frac{1}{\sqrt{\det M}} \exp\left(\frac{1}{4} c^T M^{-1} c\right) H_{m-n}^{fRg}(y); \quad (19)$$

where the symmetric  $2N \times 2N$  matrix

$$M = \begin{pmatrix} R_1 & R_{12} \\ R_{12}^T & R_2 \end{pmatrix} \quad (20)$$

with  $N \times N$  blocks  $R_1$ ;  $R_2$ ;  $R_{12}$  is expressed in terms of  $N \times N$  matrices  $R$ ;  $r$ ;  $m$  and a  $N \times N$  matrix  $e$  in the form

$$\begin{aligned} R_1 &= R - \frac{1}{2} R m^{-1} R \\ R_2 &= r - \frac{1}{2} r m^{-1} r^T \\ R_{12} &= \frac{1}{2} r m^{-1} R : \end{aligned} \quad (21)$$

Here the matrix  $e$  is transposed matrix and  $R_{12}^T$  is transposed matrix  $R_{12}$ . The  $2N$ -vector  $y$  is expressed in terms of  $N$ -vectors  $c$  and  $d$  in the form

$$y = \begin{pmatrix} c \\ d \end{pmatrix}; \quad (22)$$

where the  $N$  {vectors  $y_1$  and  $y_2$  are

$$\begin{aligned} y_1 &= \frac{1}{4} (R m^{-1} + m^{-1} R) c \\ y_2 &= \frac{1}{4} (r m^{-1} + m^{-1} \tilde{r}) c + d: \end{aligned} \quad (23)$$

The multivariable Hermite polynomials describe the photon distribution function for the multimode mixed and pure correlated light [22], [23], [24]. The nonclassical state of the light may be created due to nonstationary Casimir effect [25], and the multimode Husimi oscillator is the model to describe the behaviour of the squeezed and correlated photons.

## References

- [1] K. Husimi, Progr. Theor. Phys. 9 (1953) 238; *ibid.* 381.
- [2] I. A. Malkin, V. I. Man'ko, and D. A. Trifonov, Phys. Lett. A 30 (1969) 414; Phys. Rev. D 2 (1970) 1371.
- [3] I. A. Malkin and V. I. Man'ko, Phys. Lett. A 32 (1970) 243.
- [4] I. A. Malkin and V. I. Man'ko, Dynamical Symmetries and Coherent States of Quantum Systems (Nauka Publishers, Moscow, 1979) [in Russian].
- [5] V. V. Dodonov and V. I. Man'ko, Invariants and Evolution of Nonstationary Quantum Systems, Proc. of Lebedev Physics Institute 183, ed. M. A. Markov (Nova Science, Commack, N. Y., 1989).
- [6] L. C. Biedenharn, J. Phys. A 22 (1989) L873.
- [7] A. J. Macfarlane, J. Phys. A 22 (1989) 4581.
- [8] V. I. Man'ko, G. Marmo, S. Solimeno, and F. Zaccaria, Int. J. Mod. Phys. A 8 (1993) 3577.
- [9] V. I. Man'ko, G. Marmo, S. Solimeno, and F. Zaccaria, Phys. Lett. A 176 (1993) 173.
- [10] V. V. Dodonov, I. A. Malkin, and V. I. Man'ko, Physica 72 (1974) 597.
- [11] B. Yurke and D. Stoler, Phys. Rev. Lett. 57 (1986) 13.
- [12] R. J. Glauber, Phys. Rev. Lett. 10 (1963) 84.
- [13] E. Schrodinger, Ber. Kgl. Akad. Wiss. Berlin 24 (1930) 296.
- [14] V. V. Dodonov, E. V. Kurmyshev, and V. I. Man'ko, Phys. Lett. A 79 (1980) 150.
- [15] N. A. Ansari and V. I. Man'ko, Phys. Rev. A 50 (1994) 1942.

- [16] V . V . Dodonov, D . E . Nikonov, and V . I. Man'ko, Even and odd coherent states (Schrodinger cat states) for multimode parametric systems, Preprint INFN-TH-NA-93/49, DSF-T-93/49 (1993); Phys. Rev. (submitted).
- [17] M . M . Nieto and D . R . Truax, Phys. Rev. Lett. 71 (1993) 2843.
- [18] N . A . Ansari, L . Di Fiore, M . A . Man'ko, V . I. Man'ko, R . Romano, S . Solimeno, and F . Zaccaria, Phys. Rev. A 49 (1994) 2151.
- [19] I . A . Malkin, V . I. Man'ko, and D . A . Trifonov, J. Math. Phys. 14 (1973) 576.
- [20] E . C . G . Sudarshan, Uncertainty relations, zero point energy and the linear canonical group, Proc. of Second International Workshop on Squeezed States and Uncertainty Relations, Moscow, 25-29 May, 1992, eds. D . Han, Y . S . K im , and V . I. Man'ko (NASA Conference Publication 3219, 1993) p. 241.
- [21] O . Costas, R . Lopez-Pena, and V . I. Man'ko (this Proceedings).
- [22] V . V . Dodonov, O . V . Man'ko, and V . I. Man'ko, Phys. Rev. A 50 (1994) 813.
- [23] V . V . Dodonov and V . I. Man'ko, J. Math. Phys. 35 (1994) 4277.
- [24] V . V . Dodonov, J. Phys. A 27 (1994) 6191.
- [25] V . I. Man'ko, J. of Soviet Laser Research (Plenum Press, N . Y .) 12 (1991) N5.