

Feynman-Kac Kernels in Markovian Representations of the Schrodinger Interpolating Dynamics

Piotr Garbaczewski and Robert Ołkiewicz
Institute of Theoretical Physics, University of Wrocław,
PL-50 204 Wrocław, Poland

February 15, 2022

PACS numbers: 02.50-r, 05.40+j, 03.65-w

Abstract

Probabilistic solutions of the so called Schrodinger boundary data problem provide for a unique Markovian interpolation between any two strictly positive probability densities designed to form the input-output statistics data for the process taking place in a finite-time interval. The key issue is to select the jointly continuous in all variables positive Feynman-Kac kernel, appropriate for the phenomenological (physical) situation. We extend the existing formulations of the problem to cases when the kernel is not a fundamental solution of a parabolic equation, and prove the existence of a continuous Markov interpolation in this case. Next, we analyze the compatibility of this stochastic evolution with the original parabolic dynamics, while assumed to be governed by the temporally adjoint pair of (parabolic) partial differential equations, and prove that the pertinent random motion is a diffusion process. In particular, in conjunction with Born's statistical interpretation postulate in quantum theory, we consider stochastic processes which are compatible with the Schrodinger picture quantum evolution.

I. Motivation: Schrodinger's interpolation problem through Feynman-Kac kernels

The issue of deriving a microscopic dynamics from the (phenomenologically or numerically motivated, by approximating the frequency distributions) input-output

statistics data was addressed, as the Schrodinger problem of a probabilistic interpolation, in a number of publications [1]–[10]. We shall consider Markovian propagation scenarios so remaining within the well established framework, where for any two Borel sets $A, B \subset \mathbb{R}^n$ on which the respective strictly positive boundary densities $\rho(x; 0)$ and $\rho(x; T)$ are defined, the transition probability $m(A; B)$ from the set A to the set B in the time interval $T > 0$ has a density given in a specific factorized form :

$$\begin{aligned} m(x; y) &= \int_{\mathbb{R}^n} f(x) k(x; 0; y; T) g(y) dy \\ m(A; B) &= \int_A dx \int_B dy m(x; y) \\ \rho(x; y) &= \rho(x; 0); \quad dx m(x; y) = \rho(y; T) \end{aligned} \quad (1)$$

Here, $f(x); g(y)$ are the a priori unknown functions, to come out as solutions of the integral (Schrodinger) system of equations (1), provided that in addition to the density boundary data we have in hands any strictly positive, continuous in space variables function $k(x; 0; y; T)$. Our notation makes explicit the dependence (in general irrelevant) on the time interval endpoints. It anticipates an important restriction we shall impose, that $k(x; 0; y; T)$ must be a strongly continuous dynamical semigroup kernel: it will secure the Markov property of the sought for stochastic process.

It is the mathematical discovery [4] that, without the semigroup assumption but with the prescribed, nonzero boundary data $\rho(x; 0); \rho(y; T)$ and with the strictly positive continuous function $k(y; 0; x; T)$, the Schrodinger system (1) of integralequations admits a unique solution in terms of two nonzero, locally integrable functions $f(x); g(y)$ of the same sign (positive, everything is up to a multiplicative constant).

If $k(y; 0; x; T)$ is a particular, con ned to the time interval endpoints, form of a concrete semigroup kernel $k(y; s; x; t); 0 \leq s \leq t \leq T$, let it be a fundamental solution associated with (5) (whose existence a priori is not granted), then there exists [5, 8, 9, 10, 7] a function $p(y; s; x; t)$:

$$p(y; s; x; t) = k(y; s; x; t) \frac{\rho(x; t)}{\rho(y; s)} \quad (2)$$

where

$$\begin{aligned} \rho(x; t) &= \int_{\mathbb{R}^n} dy k(x; t; y; T) g(y) \\ \rho(y; s) &= \int_{\mathbb{R}^n} dx k(x; 0; y; s) f(x) \end{aligned} \quad (3)$$

which implements a consistent propagation of the density $\rho(x;t) = \rho(x;t) \rho(x;t)$ between its boundary versions, according to:

$$\rho(x;t) = \int_0^Z p(y;s;x;t) \rho(y;s) dy \quad (4)$$

$$0 \leq s \leq t < T$$

For a given semigroup which is characterized by its generator (Hamiltonian), the kernel $k(y;s;x;t)$ and the emerging transition probability density $p(y;s;x;t)$ are unique in view of the uniqueness of solutions $f(x); g(y)$ of (1). For Markov processes, the knowledge of the transition probability density $p(y;s;x;t)$ for all intermediate times $0 \leq s < t \leq T$ suffices for the derivation of all other relevant characteristics.

In the framework of the Schrodinger problem the choice of the integral kernel $k(y;0;x;T)$ is arbitrary, except for the strict positivity and continuity demand. As long as there is no "natural" physical motivation for its concrete functional form, the problem is abstract and of no direct physical relevance.

However, in the context of parabolic partial differential equations this "natural" choice is automatically settled if the Feynman-Kac formula can be utilized to represent solutions. Indeed, in this case an unambiguous strictly positive semigroup kernel which is a continuous function of its arguments, can be introduced for a broad class of (admissible [11]) potentials. Time dependent potentials are here included as well [12, 13]. Moreover, in Ref. [8] we have discussed a possible phenomenological significance of the Feynman-Kac potentials, as contrasted to the usual identification of Smoluchowski drifts with force fields affecting particles (up to a coefficient) in the standard theory of stochastic diffusion processes.

In the existing probabilistic investigations [5, 6, 14, 8, 9], based on the exploitation of the Schrodinger problem strategy, it was generally assumed that the kernel actually is a fundamental solution of the parabolic equation. It means that the kernel is a function with continuous derivatives: first order with respect to time, second order with respect to space variables. Then, the transition probability density defined by (2) is a fundamental solution of the Fokker-Planck (second Kolmogorov) equation in the pair $x;t$ of variables, and as such is at the same time a solution of the backward (first Kolmogorov) equation in the pair $y;s$. This feature was exploited in [8, 9].

There is a number of mathematical subtleties involved in the fundamental solution notion, since in this case, the Feynman-Kac kernel must be a solution of the parabolic equation itself. In general, Feynman-Kac kernels may have granted the existence status, even as continuous functions [11, 13, 15], but may not be differentiable, and need not to be solutions of any conceivable partial differential equations.

To our knowledge, this complication in the study of Markovian representations of the Schrodinger interpolating dynamics (and the quantum Schrodinger picture dynamics in particular) has never been addressed in the literature. Moreover, it is far from being obvious that this Markovian interpolation actually is a diffusion process.

II. Schrodinger's interpolation problem : general derivation of the stochastic evolution

1. The Schrodinger system of integral equations

We shall complement our previous analysis [8, 9] by discussing the issue in more detail. It turns out the crucial step lies in a proper choice of the strictly positive and continuous function $k(y;s;x;t); s < t$ which, if we want to construct a Markov process, has to satisfy the Chapman-Kolmogorov (semigroup composition) equation. To proceed generally let us consider a pair of partial differential equations for real functions $u(x;t)$ and $v(x;t)$:

$$\partial_t u(x;t) = -\frac{1}{2} \Delta u(x;t) - c(x;t)u(x;t) \quad (5)$$

$$\partial_t v(x;t) = -\frac{1}{2} \Delta v(x;t) + c(x;t)v(x;t)$$

where, we have eliminated all unnecessary dimensional parameters.

Usually, [15, 11], $c(x;t)$ is assumed to be a continuous and bounded from below function. We shall adopt weaker conditions. Namely, let us decompose $c(x;t)$ into a sum of positive and negative terms: $c(x;t) = c_+(x;t) - c_-(x;t)$; $c_- \geq 0$ where (a) $c_-(x;t)$ is bounded, while (b) $c_+(x;t)$ is bounded on compact sets of $R \times [0;T]$. It means that $c(x;t)$ is bounded from below and locally bounded from above. Clearly, $c(x;t)$ needs not to be a continuous function and then we encounter weak solutions of (5) which admit discontinuities.

With the first (forward) equation (5) we can immediately associate an integral kernel of the time-dependent semigroup (the exponential operator should be understood as the time-ordered expression):

$$k(y;s;x;t) = \left[\exp \left(-\frac{1}{2} \int_s^t H(\cdot) d\tau \right) \right](y;x) \quad (6)$$

where $H(\cdot) = -\frac{1}{2} \Delta + c(\cdot)$. It is clear, that for discontinuous $c(x;t)$, no fundamental solutions are admitted by (5).

By the Feynman-Kac formula, [13, 12], we get

$$k(y; s; x; t) = \int_s^T \exp\left[-\int_s^t c(X_t(\omega); \omega) d\omega\right] d\omega^{(y;s)}_{(x;t)}(\omega) \quad (7)$$

where $d^{(y;s)}_{(x;t)}(\omega)$ is the conditional Wiener measure over sample paths of the standard Brownian motion.

It is well known that k is strictly positive in case of $c(x; t)$ which is continuous and bounded from below; typical proofs are given under an additional assumption that c does not depend on time [15]. However, our assumptions about $c(x; t)$ were weaker, and to see that nonetheless k is strictly positive we shall follow the idea of Theorem 3.3.3 in [15]. Namely, the conditional Wiener measure $d^{(y;s)}_{(x;t)}$ can be written as follows

$$d^{(y;s)}_{(x;t)} = [4(t-s)]^{-1/2} \exp\left[-\frac{(x-y)^2}{4(t-s)}\right] d^{(y;s)}_{(x;t)} \quad (8)$$

where $d^{(y;s)}_{(x;t)}$ is the normalised Wiener measure [11]. We can always choose a certain number $r > 0$ to constrain the event (sample path) set

$$(r) = \{\omega : X_s(\omega) = y; X_t(\omega) = x; \sup_{s \leq \tau \leq t} |X(\tau)| \leq r\} \quad (9)$$

It comprises these sample trajectories which are bounded by r on the time interval $[s; t]$. In the above, $X_t(\omega)$ is the value taken by the random variable $X(t)$ at time t , while a concrete ω -th path is sampled. By properly tuning r , we can always achieve

$$\int_{(r)} d^{(y;s)}_{(x;t)} \geq \frac{1}{2} \quad (10)$$

which implies that

$$k(y; s; x; t) \geq \frac{1}{2} [4(t-s)]^{-1/2} \exp\left[-\frac{(x-y)^2}{4(t-s)}\right] \exp[-(t-s)C] > 0 \quad (11)$$

$$C = \sup_{s \leq \tau \leq t} \int_{(r)} c_+(X(\tau); \omega) d\omega$$

where, by our assumptions, c_+ is bounded on compact sets. Consequently, the kernel k is strictly positive.

With the Schrodinger boundary data problem on mind, we must settle an issue of the continuity of the kernel. To this end, let us invoke a well known procedure of rescaling of path integrals [11, 16]: by passing from the "unscaled" sample paths $X(t)$ over which the conditional Wiener measure integrates, to the "scaled" paths of the Brownian bridge, the $(y; x)$ conditioning can be taken away from the measure.

Then, instead of sample paths ω connecting points y and x in the time interval $t - s > 0$, we consider the appropriately "scaled" paths of the Brownian bridge connecting the point 0 with 0 again, in the (scaled) time 1. It is possible, in view of the decomposition [11, 16]:

$$\omega(t) = \left(\frac{t}{t-s} - \frac{s}{t-s}\right)y + \left(\frac{s}{t-s} - \frac{t}{t-s}\right)x + \sqrt{\frac{t-s}{t}} \tilde{\omega}\left(\frac{s}{t-s}\right) \quad (12)$$

where $\tilde{\omega}$ stands for the "scaled" Brownian bridge. Then, we can write

$$k(y; s; x; t) = [4(t-s)]^{-1/2} \exp\left[-\frac{(x-y)^2}{4(t-s)}\right] d(\omega) \exp\left[\int_s^t c\left(\frac{t}{t-s}y + \frac{s}{t-s}x + \sqrt{\frac{t-s}{t}}\tilde{\omega}\left(\frac{s}{t-s}\right); \omega\right) d\omega\right] \quad (13)$$

where $d(\omega) = d_{(0;1)}^{(0;0)}(\omega)$ is the normalized Wiener measure integrating with respect to the "scaled" Brownian bridge paths, which begin and terminate at the origin 0 in-between "scaled time" instants: 0 corresponding to $\omega = s$ and 1 corresponding to $\omega = t$.

This representation of k , if combined with the assumption that $c(x; t)$ is a continuous function, allows to conclude, [11], that the kernel is continuous in all variables. However, our previous assumptions were weaker, and it is instructive to know that through suitable approximation techniques, Theorem B.7.1 in Ref.[13] proves that the kernel is jointly continuous in our case as well.

It is also clear that $k(y; s; x; t)$ satisfies the Chapman-Kolmogorov composition rule. So, the first equation (5) can be used to define the Feynman-Kac kernel, appropriate for the Schrodinger problem analysis in terms of a Markov stochastic process.

Let us consider an arbitrary (at the moment) pair of strictly positive, but not necessarily continuous, boundary densities $\rho_0(x)$ and $\rho_T(x)$. By Jamison's principal theorem [4] there exists a unique pair of strictly positive, locally (i.e. on compact sets) integrable functions $f(x)$ and $g(x)$ solving the Schrodinger system (1), e.g. such that $\rho_0(x) = f(x) \int_{\mathbb{R}} k(x; 0; y; T) g(y) dy$ and $\rho_T(x) = g(x) \int_{\mathbb{R}} k(y; 0; x; T) f(y) dy$ with the kernel $k(y; s; x; t)$ given by (7).

Let us define:

$$g(x; t) = \int_{\mathbb{R}} k(x; t; y; T) g(y) dy; \quad f(x; t) = \int_{\mathbb{R}} k(y; 0; x; t) f(y) dy \quad (14)$$

The above integrals exist at least for almost every x so that there appears the problem of the existence of a unique and continuous transition probability density

$p(y; s; x; t)$, (2). We shall assume that the function $g(y)$ is bounded at infinity. This means that there exists a constant $C > 0$ and a compact set $K \subset \mathbb{R}$ such that $g(y) \leq C$ for all $y \in \mathbb{R} \setminus K$. Then, for all $t < T$ and any sequences $h_n \rightarrow 0; s_n \rightarrow 0$, as $n \rightarrow \infty$ we get (lim stands for $\lim_{n \rightarrow \infty}$):

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}} |g(x + h_n; t + s_n) - g(x; t)| & \int_K |k(x + h_n; t + s_n; y; T) - k(x; t; y; T)| g(y) dy + \\ & \lim_{n \rightarrow \infty} \int_{\mathbb{R} \setminus K} |k(x + h_n; t + s_n; y; T) - k(x; t; y; T)| g(y) dy \quad (15) \\ & \limsup_{y \in \mathbb{R} \setminus K} |k(x + h_n; t + s_n; y; T) - k(x; t; y; T)| \int_K g(y) dy + \\ & C \lim_{n \rightarrow \infty} \int_{\mathbb{R} \setminus K} |k(x + h_n; t + s_n; y; T) - k(x; t; y; T)| dy \end{aligned}$$

The first term tends to zero because k is jointly continuous and g is locally integrable. The second one tends to zero because of the Lebesgue bounded convergence theorem. Consequently, our assumption succeeds to make $g(x; t)$ continuous on $\mathbb{R} \times [0; T)$. Similarly, we can prove that $g(x; t)$ is bounded.

Now, we can set according to (2), $p(y; s; x; t) = k(y; s; x; t)g(x; t) = g(y; s)$. Then, $p(y; s; x; t)$, $0 \leq s < t \leq T$ becomes a transition probability density of a Markov stochastic process with a factorized density $\rho(x; t) = f(x; t)g(x; t)$. Clearly, this stochastic process interpolates between the boundary data ρ_0 and ρ_T as time continuously varies from 0 to T . Notice that (15) implies the continuity of p in the time interval $[0; T)$.

Although $p(y; s; x; t)$ is continuous in all variables, we cannot be sure that the interpolating stochastic process has continuous trajectories, and no specific (e.g. Fokker-Planck) partial differential equation can be readily associated with this dynamics. Therefore, we must explicitly verify whether the associated process is stochastically continuous. If so, we should know whether it is continuous (i.e. admits continuous trajectories). Eventually, we should check the validity of conditions under which the investigated interpolation can be regarded as a diffusion process. The subsequent analysis will prove that this ultimate goal results only due to the gradual strengthening of conditions imposed on the parabolic system (5).

2. Stochastic continuity of the process

Apart from the generality of formulation of the Schrodinger interpolation problem which appears to preclude an unambiguous identification (diffusion or not) of the constructed stochastic process, we can prove in the present case, a fundamental

property of a stochastic dynamics called a stochastic continuity of the process. In this connection, compare e.g. [5, 17] and [18], where this property is linked to the uniqueness of the corresponding Markov semigroup generator. The stochastic continuity property is a necessary condition for the process to admit continuous trajectories.

The stochastic process is stochastically continuous, if for the probability of the occurrence of sample paths ω , such that the random variable values $X_t(\omega)$ along the trajectory obey $|X_t(\omega) - X_s(\omega)| \rightarrow 0$; $s < t$, the following limiting behaviour is recovered

$$\lim_{t \downarrow s} P[|\omega| : |X_t(\omega) - X_s(\omega)| \geq \epsilon] = 0 \quad (16)$$

for every positive ϵ . This demand can be written in a more handy way in terms of the transition probability density $p(y; s; x; t)$ and the density $\rho(x; t)$ of the process:

$$\lim_{t \downarrow s} \int_{|x-y| \geq \epsilon}^{\infty} dy \int_{-\infty}^{\infty} p(y; s; x; t) dx = 0 \quad (17)$$

So, for the transition density to be stochastically continuous, it suffices that

$$\lim_{t \downarrow s} \int_{|x-y| \geq \epsilon}^{\infty} p(y; s; x; s + \epsilon) dx = 0 \quad (18)$$

for almost every $y \in \mathbb{R}$.

In view of our construction, (2), we have:

$$\begin{aligned} & \lim_{t \downarrow s} \int_{|x-y| \geq \epsilon}^{\infty} p(y; s; x; s + \epsilon) dx = \\ & \frac{1}{g(y; s)} \lim_{t \downarrow s} \int_{|x-y| \geq \epsilon}^{\infty} dx \int_{-\infty}^{\infty} k(y; s; x; s + \epsilon) k(x; s + \epsilon; z; T) g(z) dz \end{aligned} \quad (19)$$

By changing the order of integrations (allowed by positivity of the involved functions) we get:

$$\begin{aligned} \lim_{t \downarrow s} \int_{|x-y| \geq \epsilon}^{\infty} p(y; s; x; s + \epsilon) dx &= \frac{1}{g(y; s)} \lim_{t \downarrow s} \int_{-\infty}^{\infty} dz g(z) \\ & \left[\int_{|x-y| \geq \epsilon}^{\infty} dx k(y; s; x; s + \epsilon) k(x; s + \epsilon; z; T) \right] \end{aligned} \quad (20)$$

Because the potential is bounded from below, $c \leq M$ for some $M > 0$, we easily arrive at the estimates (use the "scaled" Brownian bridge argument)

$$k(y; s; x; s + \epsilon) \leq (4\pi\epsilon)^{-1/2} \exp\left[-\frac{(x-y)^2}{4\epsilon}\right] \exp(M\epsilon) \quad (21)$$

and

$$k(x; s + 4s; z; T) = [4(T - s - 4s)]^{1/2} \exp\left[-\frac{(z - x)^2}{4(T - s - 4s)}\right] \exp[M(T - s - 4s)] \quad (22)$$

Then we get:

$$\begin{aligned} 0 & \lim_{4s \neq 0} \int_{\mathbb{R}} k(y; s; x; s + 4s) k(x; s + 4s; z; T) dx \\ & [4(T - s)]^{1/2} \exp[M(T - s)] \lim_{4s \neq 0} (4 - 4s)^{1/2} \\ & \int_{\mathbb{R}} dx \exp\left[-\frac{(x - y)^2}{44s}\right] \exp\left[-\frac{(z - x)^2}{4(T - s - 4s)}\right] = 0 \end{aligned} \quad (23)$$

So, by the classic Lebesgue bounded (dominated) convergence theorem, the required limiting property $\lim_{4s \rightarrow 0} \int_{\mathbb{R}} k(y; s; x; s + 4s) dx = 0$ follows and (16) holds true.

As mentioned before, the stochastic continuity of the Markov process is a necessary condition for the process to be continuous in a more pedestrian sense, i. e. to admit continuous sample paths. However, it is insufficient. Hence, additional requirements are necessary to allow for a standard diffusion process realization of solutions of the general Schrodinger problem, (1)–(3).

In the next section we shall prove that our process can be regarded as continuous, by requiring a certain correlation between the kernel $k(y; s; x; t)$ and a function $g(x; t)$, (14).

3. Continuity of the process

It is well known that a solution of a parabolic equation cannot tend to zero arbitrarily fast, when $|x| \rightarrow \infty$, [19]. Roughly speaking, it cannot fall off faster than a fundamental solution (provided it exists). In fact, the solution is known to fall off as fast as the fundamental solution, when the initial boundary data coincide with the Dirac measure. If a support of the initial data is spread (i.e. not point-wise), then the solution fall off is slower than this of the fundamental one.

In our discussion, where $g(x; t)$ is a generalized solution and $k(y; s; x; t)$ is a Feynman-Kac kernel which does not need to be a fundamental solution, we expect a similar behaviour. Mathematically, our demand will be expressed as follows. Let t be small and K be a compact subset in \mathbb{R} . Because $g(x; t)$ is supported on the whole \mathbb{R} , so in the decomposition

$$g(y; s) = \int_K k(y; s; x; t) g(x; t) dx + \int_{\mathbb{R} \setminus K} k(y; s; x; t) g(x; t) dx \quad (24)$$

the second term becomes relevant when $\|y\| \rightarrow 1$. It amounts to (in the denominator there appears $g(y;s)$):

$$\lim_{\|y\| \rightarrow 1} \frac{\int_K k(y;s;x;t)g(x;t)dx}{\int_K k(y;s;x;t)g(x;t)dx} = 0 \quad (25)$$

where χ_K is an indicator function of the set K , which is equal one for $x \in K$ and zero otherwise.

By means of the transition probability density $p(y;s;x;t)$ let us introduce a transformation

$$(T_s^t f)(y) = \int_K p(y;s;x;t)f(x)dx \quad (26)$$

of a function $f(x)$, continuous and vanishing at infinity (we shall use an abbreviation $f \in C_1(\mathbb{R})$ to express this fact). It is clear that $(T_s^t f)(x)$ is a continuous function. For a suitable compact set K we can always guarantee the property $\int_K f(x)dx < \epsilon$ for every $x \in \mathbb{R}^n$. Then, if we exploit the property $\int_{\mathbb{R}^n} p(y;s;x;t)dx = 1$ if $s < t$ and the definition of p in terms of k and g , we arrive at

$$\begin{aligned} |(T_s^t f)(y)| &\leq \int_K p(y;s;x;t)|f(x;t)|dx + \int_{\mathbb{R}^n \setminus K} p(y;s;x;t)|f(x;t)|dx \\ &\leq \left[\int_K p(y;s;x;t)dx \right] \sup_{x \in K} |f(x;t)| + \int_{\mathbb{R}^n \setminus K} p(y;s;x;t)dx \\ &\leq \left[\int_K |f(x;t)|dx \right] \frac{\int_K k(y;s;x;t)g(x;t)dx}{\int_K k(y;s;x;t)g(x;t)dx} + \end{aligned} \quad (27)$$

It implies that for small $t-s$, $\lim_{\|y\| \rightarrow 1} (T_s^t f)(y) = 0$, and so T_s^t forms an inhomogeneous in time semigroup of positive contractions on $C_1(\mathbb{R})$. For arbitrary t and s the result follows by the obvious decomposition property $T_s^t = T_s^{s_1} T_{s_1}^{s_2} \dots T_{s_n}^t$. In the well established terminology, our $p(y;s;x;t)$ is a C_1 -Feller transition function and leads to a regular Markov process, [17]. Moreover, by the stochastic continuity of $p(y;s;x;t)$, T_s^t is strongly continuous.

As yet, we do not know whether the process itself is continuous i.e. has continuous sample paths. To this end, it suffices to check whether the so called "Dynkin condition", [20]

$$\lim_{t \rightarrow s} \frac{1}{t-s} \sup_{y \in K} \left[\int_K p(y;s;x;t)dx \right] = 0 \quad (28)$$

is valid for every $\epsilon > 0$ and every compact set K . We have (remember that $g(x;t)$ is strictly positive, continuous and bounded):

$$\sup_{y \in K} \int_K p(y;s;x;t)dx = \sup_{y \in K} \frac{1}{g(y;s)} \int_K k(y;s;x;t)g(x;t)dx$$

$$\frac{\sup_x g(x;t)}{\inf_{y \in K} g(y;s)} \int_K k(y;s;x;t) dx = C \int_K k_0(x-y;t-s) dx \quad (29)$$

where (compare e.g. the previous estimate (22))

$$C = \frac{\sup_x g(x;t)}{\inf_{y \in K} g(y;s)} \exp[M(t-s)] \quad (30)$$

and $k_0(x-y;t-s)$ is the heat kernel.

Finally, we arrive at:

$$\lim_{t \rightarrow s} \frac{1}{t-s} \sup_{y \in K} \left[\int_K p(y;s;x;t) dx \right] = C \lim_{t \rightarrow s} \frac{1}{t-s} \int_K k_0(z;t-s) dz = 0 \quad (31)$$

So, the stochastic process we are dealing with, is continuous. Interestingly, "a continuous in time parameter stochastic processes which possesses the (strong) Markov property and for which the sample paths $X(t)$ are almost always (i.e. with probability one) continuous functions of t is called a diffusion process", see e.g. chapter 15 of [20].

4. The interpolating stochastic dynamics: compatibility with the temporally adjoint parabolic evolutions

The formulas (14) determine what is called, [12], the generalized solution of a parabolic equation: it admits functions which are not necessarily continuous and if continuous, then not necessarily differentiable. Before, we have established the continuity of the generalized solution $g(x;t)$ under rather mild assumption about the behaviour of $g(x)$ at spatial infinity. In fact, the same assumption works for $f(x;t)$. But nothing has been said about the differentiability of $f(x;t)$ and $g(x;t)$.

Consequently, our reasoning seems to be somewhat divorced from the original partial differential equations (5), for which we can take for granted that certain solutions $u(x;t)$ and $v(x;t)$ exist in the time interval $0 \leq t \leq T$. For this, we must assume that $c(x;t)$ is a continuous function.

Let us consider the solutions of (5) that are bounded functions of their arguments. It is instructive to point out that we do not impose any restrictions on the growth of $c(x;t)$ when $|x| \rightarrow \infty$, and consequently we do not assume that solutions of parabolic equations (5) have bounded derivatives. Then, [12], the solution

$u(x;t)$ of the forward parabolic equation (5) is known to admit the Feynman-Kac representation with the integral kernel (7), (13), where

$$u(x;t) = \int_Z k(y;s;x;t) u(y;s) dy \quad (32)$$

for $0 \leq s < t \leq T$. At this point let us define

$$U(x;t) = v(x;T-t) \quad (33)$$

for all $t \in [0;T]$ and observe that, as a consequence of the time adjoint equation (5) for which $v(x;t)$ is a solution, the newly introduced function $U(x;t)$ solves the forward equation (5):

$$\partial_t U(x;t) = \frac{1}{4} \Delta U(x;t) - c(x;T-t) U(x;t) \quad (34)$$

with a slightly rearranged potential: $c(x;t) \rightarrow c(x;T-t)$. By the assumed boundedness of the solution $v(x;t)$ of (5), we arrive at the Feynman-Kac formula

$$U(x;t) = \int_Z K(y;s;x;t) U(y;s) dy \quad (35)$$

with the corresponding kernel $K(y;s;x;t)$ of the (time ordering implicit) operator $\exp[\int_s^t H(T-t) dt]$, where $H(T-t) = \frac{1}{4} \Delta + c(x;T-t)$. Let us emphasize that in case of the time independent potential, $c(x;t) = c(x)$ for all $0 \leq t \leq T$, the kernel K coincides with k .

The previous Brownian bridge argument (12), (13) retains its validity, and we have:

$$K(y;s;x;t) = \left[\frac{4}{(t-s)} \right]^{1/2} \exp\left[-\frac{(x-y)^2}{4(t-s)} \right] \int_Z d(\gamma) \exp\left[\int_s^t c\left(\frac{t}{t-s}y + \frac{s}{t-s}x + \frac{p}{t-s}\left(\frac{s}{t-s}\right); T-t \right) d\gamma \right] \quad (36)$$

which, after specializing to the case of $s = 0; t = T$ and accounting for the invariance of the Brownian bridge measure with respect to the replacement of sample paths $\gamma(\cdot)$ by sample paths $\gamma(T-\cdot)$, [7, 27], gives rise to:

$$K(y;0;x;T) = \left[\frac{4}{T} \right]^{1/2} \exp\left[-\frac{(x-y)^2}{4T} \right] \int_Z d(\gamma) \exp\left[\int_0^T c\left(\frac{1}{T}y + \left(1 - \frac{1}{T}\right)x + \frac{p}{T}\left(\frac{1}{T}\right); \gamma \right) d\gamma \right] \quad (37)$$

where $\gamma = T - \gamma$.

A comparison of (37) with (13) proves that we have derived an identity:

$$K(y;0;x;T) = k(x;0;y;T) \quad (38)$$

whose immediate consequence is the formula

$$U(x;T) = v(x;0) = \int_0^Z k(x;0;y;T)v(y;T)dy \quad (39)$$

for the backward propagation of $v(y;T)$ into $v(x;0)$.

We shall utilize (39) and (32), under an additional assumption that the previous, hitherto arbitrary, probability density data $\rho_0(x); \rho_T(x)$, actually are determined by the initial and terminal values of the solutions $u(x;t); v(x;t)$ of (5), according to:

$$\begin{aligned} \rho_0(x) &= u(x;0)v(x;0) \\ \rho_T(x) &= u(x;T)v(x;T) \end{aligned} \quad (40)$$

Our present aim is to show that with this assumption, we can identify the (still abstract) functions $f(x;t), g(x;t)$, (14), with $u(x;t)$ and $v(x;t)$ respectively. By (32), (39) there holds:

$$\begin{aligned} \rho_0(x) &= u(x;0) \int_0^Z k(x;0;y;T)v(y;T)dy \\ \rho_T(x) &= v(x;T) \int_0^Z k(y;0;x;T)u(y;0)dy \end{aligned} \quad (41)$$

and, in view of the uniqueness of solution of the Schrodinger system, once the boundary densities and the continuous strictly positive kernel are specified, we realize that the propagation formulas (14) involve solutions of (5) through the respectively initial and terminal data:

$$\begin{aligned} f(x) &= u(x;0) \\ g(x) &= v(x;T) \end{aligned} \quad (42)$$

Moreover, (5), (14) imply that $f(x;t) = u(x;t)$ holds true identically for all $t \in [0;T]$.

What remains to be settled is whether the function $g(x;t)$ can be identified with the solution $v(x;t)$ of (5) for all $t \in [0;T]$.

This property is obvious, when the time independent potential $c(x)$ is investigated instead of the more general $c(x;t)$. As well, the identification is with no doubt in case when $k(y;s;x;t)$ is a fundamental solution of the parabolic equation in variables $x;t$. In this case, $k(y;s;x;t)$ is a unique solution of the system (5), and solves the adjoint equation in variables $y;s$, [21, 22, 23]. Then, because $f(x); g(x)$ are locally integrable, an immediate consequence is, [24], that $f(x;t)$ and $g(x;t)$

are positive solutions of (5). The identification of them with $u(x;t)$ and $v(x;t)$ respectively, follows from the uniqueness of positive solutions, [22].

Let us begin from a minor generalization of (22), and define:

$$U_s(x;t) = v(x;T+s-t) \quad ; \quad t \in [s;T] \quad (43)$$

Clearly, a parabolic equation (34) is satisfied by $U_s(x;t)$, if instead of $c(x;T-t)$, the potential $c(x;T+s-t)$ is introduced. An immediate propagation formula follows

$$U_s(x;t) = \int_s^T K_s(y;s;x;t) U_s(y;s) dy \quad (44)$$

The integral kernel K_s differs from the previous K , (36), in the explicit time dependence of the potential $c(x;T+s-t) \neq c(x;T-t)$. By putting $T = t$ in (44) we get:

$$v(x;s) = \int_s^T K_s(y;s;x;T) v(y;T) dy \quad (45)$$

and by the previous part of our demonstration we know that

$$g(x;s) = \int_s^T k(x;s;y;T) v(y;T) dy \quad (46)$$

At this point, it is enough to prove that the identity (cf. (38))

$$K_s(y;s;x;T) = k(x;s;y;T) \quad (47)$$

takes place for any $s; 0 \leq s \leq T$.

Let us exploit the Brownian bridge scaling (13) again, so that

$$k(x;s;y;T) = [4(T-s)]^{-1/2} \exp\left[-\frac{(x-y)^2}{4(T-s)}\right] \quad (48)$$

$$\int_s^T d\left(\frac{s}{T}\right) \exp\left[\int_s^T c\left(\frac{T}{T-s}x + \frac{s}{T-s}y + \frac{p}{T-s}\left(\frac{s}{T}\right); \frac{s}{T}\right) d\left(\frac{s}{T}\right)\right]$$

and, analogously

$$K_s(y;s;x;T) = [4(T-s)]^{-1/2} \exp\left[-\frac{(x-y)^2}{4(T-s)}\right] \quad (49)$$

$$\int_s^T d\left(\frac{s}{T}\right) \exp\left[\int_s^T c\left(\frac{T}{T-s}y + \frac{s}{T-s}x + \frac{p}{T-s}\left(\frac{s}{T}\right); T+s\right) d\left(\frac{s}{T}\right)\right]$$

By changing:

$$\left(\frac{s}{T}\right) \rightarrow \left(1 - \frac{s}{T}\right) = \left(\frac{T}{T-s}\right) \quad (50)$$

and substituting $\tau = T + s$, where s only is the running variable, we finally recover

$$K_s(y; s; x; T) = [4(T-s)]^{-1/2} \exp\left[-\frac{(x-y)^2}{4(T-s)}\right] \quad (51)$$

$$\int_{\mathbb{R}^d} d(y) \exp\left[\int_{\tau}^T c\left(\frac{s}{T-s}y + \frac{T}{T-s}x + \frac{p}{T-s}\left(\frac{s}{T-s}\right); \right) ds\right] = k(x; s; y; T)$$

Hence,

$$g(x; s) = v(x; s) \quad (52)$$

is valid for all time instants $0 \leq s \leq T$. This implies that $p(y; s; x; t) = k(y; s; x; t) \frac{v(x; t)}{v(y; s)}$ defines a consistent transition probability density of the continuous Markovian interpolation.

We have succeeded to prove that:

(i) If a continuous, strictly positive Feynman-Kac kernel of the forward parabolic equation (5) is employed to solve the Schrodinger boundary data problem (1) for an arbitrary pair of nonzero probability densities $\rho_0(x)$ and $\rho_T(x)$, then we can construct a Markov stochastic process, which is continuous and provides for an interpolation between these boundary data in the time interval $[0; T]$.

(ii) Given the time adjoint parabolic system (5) with bounded solutions $u(x; t); v(x; t)$ in the time interval $[0; T]$. If the boundary densities are defined according to (40), then the Schrodinger problem (1)-(3) provides us with a unique continuous Markov interpolation, that is compatible with the time evolution of $\psi(x; t) = u(x; t)v(x; t); t \in [0; T]$.

5. Whence diffusions?

Our strategy, of deducing a probabilistic solution of the Schrodinger boundary data problem in terms of Markov stochastic processes running in a continuous time, was accomplished in a number of steps accompanied by the gradual strengthening of restrictions imposed on the Feynman-Kac potential, to yield a continuous process (cf. Section II.3), and eventually to get it compatible with a given a priori parabolic evolution (Section II.4). In a broad sense, [20], it can be named a diffusion.

However, this rather broad definition of the diffusion process is significantly narrowed in the physical literature: while demanding the continuity of the process, the additional restrictions are imposed to guarantee that the mean and variance of the infinitesimal displacements of the process have the standard meaning of the drift and diffusion coefficient, respectively, [26].

According to the general wisdom, diffusions arise in conjunction with the parabolic evolution equations, since then only the conditional averages are believed to make sense in the local description of the dynamics. It is not accidental that forward parabolic equations (5) are commonly called the generalized diffusion equations. Also, the fact that the Feynman-Kac formula involves the integration over sample paths of the Wiener process, seems to suggest some diffusive features of the Schrodinger interpolation, even if we are unable to establish this fact in a canonical manner.

Clearly, the conditions valid for any $\epsilon > 0$:

- (a) there holds $\lim_{t \rightarrow s} \frac{1}{t-s} \int_{|y-x| \leq \epsilon} p(y;s;x;t) dx = 0$, (notice that (a) is a direct consequence of the stronger Dynkin condition, (28)),
 - (b) there exists a drift function $b(x;s) = \lim_{t \rightarrow s} \frac{1}{t-s} \int_{|y-x| \leq \epsilon} (y-x)p(x;s;y;t) dy$,
 - (c) there exists a diffusion function $a(x;s) = \lim_{t \rightarrow s} \frac{1}{t-s} \int_{|y-x| \leq \epsilon} (y-x)^2 p(x;s;y;t) dy$,
- are conventionally interpreted to define a diffusion process, [26].

To our knowledge, no rigorous demonstration is available in the Schrodinger problem context, in case when the involved semigroup kernel is not a fundamental solution of the parabolic equation.

Let us impose a restriction on a lower bound of a solution $v(x;t)$ of the backward equation (5). Namely, we assume that there exist constants $c_1 > 0; c_2 > 0$ such that $v(y;s) \geq c_1 \exp(-c_2 y^2)$ for all $s \in [0;t]; t < T$. This property was found to be respected by a large class of parabolic equations, [25], and it automatically ensures that the condition (25) of Section II.3 is satisfied. Indeed:

$$0 \leq \lim_{|y| \rightarrow 1} \frac{1}{v(y;s)} \int_{|x| \leq 1} k(y;s;x;t) v(x;t) dx$$

$$\leq \frac{1}{c_1} [4(t-s)]^{1/2} \exp[M(t-s)]$$

$$[\sup_{x \in K} v(x;t)] \lim_{|y| \rightarrow 1} \exp(c_2 y^2) \int_K \exp\left[-\frac{(x-y)^2}{4(t-s)}\right] dx = 0$$

if $t-s \rightarrow 0$ for sufficiently small $\epsilon > 0$ (like for example $\epsilon = 1/16$).

It is our purpose to complete the previous analysis by demonstrating that, with the above assumption on $v(x;t)$, the continuous Markov process we have constructed actually is the diffusion process.

Our subsequent arguments will rely on the Dynkin treatise [17]. It is well known that the infinitesimal (local) characteristics of a continuous Markov process can be defined in terms of its, so called, characteristic operator. It is closely linked with the standard infinitesimal (Markov) generator of the process, and we shall

take advantage of this link in below. Let us agree, following Dynkin, to call a continuous Markov process a diffusion, if its characteristic operator U is defined on twice differentiable functions (we skip more detailed definition, [17]). In this case $x \mapsto x - x_0$ and $x \mapsto (x - x_0)^2$ allow for the definition of a drift and diffusion function respectively:

$$\begin{aligned} [U(x - x_0)](x_0; s) &= b(x_0; s) \\ [U((x - x_0)^2)](x_0; s) &= a(x_0; s) \end{aligned} \quad (54)$$

By results of Sections II.3 and II.4 we know that our transition probability density $p(y; s; x; t) = k(y; s; x; t) \frac{v(x; t)}{v(y; s)}$, inspired by the Schrodinger boundary data problem, gives rise to a continuous Markov process. To see whether it can be regarded as a diffusion, we must verify the above two defining properties (54).

At first, let us consider the infinitesimal operator A (Markov generator) of the corresponding strongly continuous semigroup $T_s^t : C_1(\mathbb{R}) \rightarrow C_1(\mathbb{R})$, which we have introduced via the formula (26). We are interested in domain properties of A , in view of the fact that the characteristic operator U is a natural extension of A , $A \subset U$, [17].

We denote $C_c^2(\mathbb{R})$ the space of continuous functions with compact support which possess continuous derivatives up to second order. For $h \in C_c^2(\mathbb{R})$ we have

$$\begin{aligned} \lim_{\# \rightarrow 0} \frac{1}{\#} \left[\int_1^{Z+\frac{1}{\#}} p(y; s; x; s + \frac{1}{\#}) h(x) dx - h(y) \right] &= \\ \frac{1}{v(y; s)} \lim_{\# \rightarrow 0} \frac{1}{\#} \left[\int_1^{Z+\frac{1}{\#}} k(y; s; x; s + \frac{1}{\#}) v(x; s + \frac{1}{\#}) h(x) dx - v(y; s) h(y) \right] \end{aligned} \quad (55)$$

Because v is continuously differentiable with respect to time, we have

$$v(x; s + \frac{1}{\#}) = v(x; s) + \frac{1}{\#} \partial_t v(x; s^0) \quad (56)$$

where (cf. the standard Taylor expansion formula) $s^0 = s + \frac{1}{\#}$; $0 \leq \frac{1}{\#} \leq 1$. Hence

$$\begin{aligned} \lim_{\# \rightarrow 0} \frac{1}{\#} \left[\int_1^{Z+\frac{1}{\#}} p(y; s; x; s + \frac{1}{\#}) h(x) dx - h(y) \right] &= \\ \frac{1}{v(y; s)} \lim_{\# \rightarrow 0} \frac{1}{\#} \left[\int_1^{Z+\frac{1}{\#}} dx k(y; s; x; s + \frac{1}{\#}) v(x; s) h(x) - v(y; s) h(x) \right] + \\ \frac{1}{v(y; s)} \lim_{\# \rightarrow 0} \left[\int_1^{Z+\frac{1}{\#}} k(y; s; x; s + \frac{1}{\#}) \partial_t v(x; s^0) h(x) dx \right] \end{aligned} \quad (57)$$

We shall exploit the strongly continuous semigroup evolution associated with the parabolic system (5). Because of the domain property: $C_c^1(\mathbb{R}) \subset D(H)$ the smooth

functions with compact support are acted upon by $H = 4 - c(x; s)$ and H is closed as an operator on $C_1(\mathbb{R})$. But then also $C_c^2(\mathbb{R}) \subset D(H)$ and so the first term in (57) takes the form :

$$\frac{1}{v(y; s)} [4 (vh)(y; s) - c(y; s)v(y; s)h(y)] \quad (58)$$

while the second equals

$$\frac{1}{v(y; s)} [\partial_s v(y; s)]f(y) = \frac{1}{v(y; s)} [4 v(y; s) + c(y; s)v(y; s)]f(y) \quad (59)$$

Thus, (55) is point-wise convergent:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left[\sum_{j=1}^{Z+1} p(y; s; x; s + \frac{j}{n}) h(x) dx - h(y) \right] = \quad (60)$$

$$\frac{1}{v(y; s)} [4 v(y; s)h(y) + 2r v(y; s)r h(y) + v(y; s)4 h(y) - c(y; s)v(y; s)h(y)$$

$$(4 v(y; s))h(y) + c(y; s)v(y; s)h(y)] = 4 h(y) + 2\left(\frac{r v}{v}\right)(y; s)r h(y)$$

Now, we shall establish the boundedness of:

$$\sup_{y \in \mathbb{R}, \rho < \infty} \left[\frac{1}{n} \sum_{j=1}^{Z+1} p(y; s; x; s + \frac{j}{n}) h(x) dx - h(y) \right] \quad (61)$$

for some small ϵ .

Because $C_c^2(\mathbb{R}) \subset D(H)$, so there holds

$$\frac{1}{n} \left[\sum_{j=1}^{Z+1} k(y; s; x; s + \frac{j}{n}) v(x; s) h(x) dx - v(y; s)h(y) \right] \leq [4 - c(y; s)](vf)(y; s) \quad (62)$$

uniformly in y , as $\epsilon > 0$. It implies that for any compact set K there is

$$\sup_{y \in K, \rho < \infty} \left[\frac{1}{n} \sum_{j=1}^{Z+1} p(y; s; x; s + \frac{j}{n}) h(x) - h(y) \right] \quad (63)$$

$$\left[\sup_{y \in K} \frac{1}{v(y; s)} \right] \sup_{y \in K, \rho < \infty} \left[\frac{1}{n} \sum_{j=1}^{Z+1} k(y; s; x; s + \frac{j}{n}) v(x; s) h(x) dx - v(y; s)h(y) \right] + \sum_{j=1}^{Z+1} k(y; s; x; s + \frac{j}{n}) [v(x; s)h(x) dx] < 1$$

We have thus the required boundedness for all $y \in K$ i.e. on compact sets.

For $y \in \mathbb{R}^n_K$ we shall make the following estimations. Because the support of h is compact, we can define $\text{supp } h \subset [-n; n]$ for some natural number n . Let $K = [-3n; 3n]$. Then:

$$\begin{aligned} \sup_{y \in \mathbb{R}^n_K, |y| < \frac{1}{2}} \int_{-1}^1 p(y; s; x; s + \frac{1}{2}) h(x) dx - h(y) &= \\ \sup_{y \in \mathbb{R}^n_K, |y| < \frac{1}{2}} \int_K p(y; s; x; s + \frac{1}{2}) h(x) dx & \quad (64) \\ [\sup_{x \in K} |h(x)|] \sup_{y \in \mathbb{R}^n_K, |y| < \frac{1}{2}} \int_K \frac{1}{v(y; s)} k(y; s; x; s + \frac{1}{2}) v(x; s + \frac{1}{2}) dx \\ [\sup_{x \in K} |h(x)|] [\sup_{x \in K, |s| \leq s_0 + \frac{1}{2}} v(x; s_0)] \sup_{y \in \mathbb{R}^n_K, |y| < \frac{1}{2}} \int_K \frac{1}{v(y; s)} k(y; s; x; s + \frac{1}{2}) dx \end{aligned}$$

In view of our assumption $v(y; s) = c_1 \exp(-c_2 y^2)$, there holds:

$$\begin{aligned} \sup_{y \in \mathbb{R}^n_K, |y| < \frac{1}{2}} \int_K p(y; s; x; s + \frac{1}{2}) h(x) dx & \quad (65) \\ C \sup_{|y| \leq 3n, |y| < \frac{1}{2}} \exp(c_2 y^2) \int_K \exp[-\frac{(x - y)^2}{4}] dx \end{aligned}$$

where

$$C = c_1 (4)^{-1/2} \exp(M) [\sup_{x \in K} |h(x)|] \sup_{|s| \leq s_0 + \frac{1}{2}} v(x; s_0) \quad (66)$$

If we choose $\epsilon = 1/16$, then

$$\exp(c_2 y^2) \int_K \exp[-\frac{(x - y)^2}{4}] dx \leq 4 \exp(-\frac{n^2}{4}) \quad (67)$$

for every $|y| \leq 3n$, and so

$$\sup_{y \in \mathbb{R}^n_K, |y| < \frac{1}{2}} \int_K p(y; s; x; s + \frac{1}{2}) h(x) dx \leq 4C \sup_{|y| < \frac{1}{2}} \exp(-\frac{n^2}{4}) < 1 \quad (68)$$

Consequently, the desired boundedness (62) holds true for all $y \in \mathbb{R}^n$, together with the previously established point-wise convergence (61).

Altogether, it means, [17], that the weak generator of T_s^t is defined at least on $C_c^2(\mathbb{R})$. Moreover, while acting on $h \in C_c^2(\mathbb{R})$ it gives $\frac{1}{2} h'' + (r \ln v) r h$. Because T_s^t is strongly continuous in $C_1(\mathbb{R})$, the Markov generator A coincides with the weak generator, [17], i.e. $A = \frac{1}{2} h'' + (r \ln v) r h$ on $C_c^2(\mathbb{R})$.

Finally, let us choose $h_0 \in C_c^2(\mathbb{R})$ such that $h_0(x) = 1$ in some neighbourhood of the point x_0 . Then, $(x - x_0)h_0(x)$ and $(x - x_0)^2 h_0(x)$ both belong to $C_c^2(\mathbb{R})$ and therefore:

$$A[(x - x_0)h_0](x_0; s) = \frac{1}{2} [(x - x_0)h_0](x_0) + \quad (69)$$

$$2(r \ln v)(x_0; s) r[(x - x_0)h_0](x_0) = 2(r \ln v)(x_0; s)$$

$$A[(x - x_0)^2 h_0](x_0; s) = 2$$

Because $A \in U$ and U is a local operator, [17], we have the following inclusion $C_c^2(\mathbb{R}) \subset D(U)$ and (we can get rid of h_0):

$$[U(x - x_0)](x_0; s) = 2(r \ln v)(x_0; s) \quad (70)$$

$$[U(x - x_0)^2](x_0; s) = 2$$

It means that we indeed obtain a diffusion process with the drift $r \ln v$ and a constant diffusion coefficient, according to the standards of [5, 27, 28].

It is worth emphasizing that since $(x - x_0)h_0(x)$ and $(x - x_0)^2 h_0(x)$ belong to $D(A)$, and since functions from $C_c^2(\mathbb{R})$ can be used to approximate, under an integral, an indicator function of the set $[x_0 - \epsilon; x_0 + \epsilon]; \epsilon > 0$, we can directly evaluate:

$$\lim_{t \downarrow s} \frac{1}{t - s} \int_{x_0 - \epsilon}^{x_0 + \epsilon} p(x_0; s; x; t) (x - x_0) h_0(x) dx = \quad (71)$$

$$\lim_{t \downarrow s} \frac{1}{t - s} \int_{x_0 - \epsilon}^{x_0 + \epsilon} p(x_0; s; x; t) (x - x_0) dx = 2(r \ln v)(x_0; s)$$

and similarly

$$\lim_{t \downarrow s} \frac{1}{t - s} \int_{x_0 - \epsilon}^{x_0 + \epsilon} p(x_0; s; x; t) (x - x_0)^2 dx = 2 \quad (72)$$

Because the Dynkin condition (28) implies that

$$\lim_{t \downarrow s} \frac{1}{t - s} \int_{x_0 - \epsilon}^{x_0 + \epsilon} p(x_0; s; x; t) dx = 0 \quad (73)$$

we arrive at the commonly accepted definition of the diffusion process, summarized in formulas (71)–(73), with the functional expression for the drift, (71), given in the familiar, [5, 27, 8], gradient form.

III. Nonstationary Schrodinger dynamics: from the Feynman-Kac kernel to diffusion process

In our previous paper [9], the major conclusion was that in order to give a definitive probabilistic description of the quantum dynamics as a unique diffusion process solving Schrodinger's interpolation problem, a suitable Feynman-Kac semigroup must be singled out. Let us point out that the measure preserving dynamics, permitted in the presence of conservative force fields, was investigated in [8], see also [29, 12].

The present analysis was performed quite generally and extends to the dynamics affected by time dependent external potentials, with no clear-cut discrimination between the nonequilibrium statistical physics and essentially quantum evolutions. The formalism of Section II encompasses both groups of problems. Presently, we shall restrict our discussion to the free Schrodinger picture quantum dynamics. Following Ref. [9] we shall discuss the rescaled problem so as to eliminate all dimensional constants.

The free Schrodinger evolution $i\partial_t = -\frac{1}{4}$ implies the following propagation of a specific Gaussian wave packet:

$$\langle x;0 \rangle = (2)^{-1/4} \exp\left(-\frac{x^2}{4}\right) \quad (74)$$

$$\langle x;t \rangle = \left(\frac{2}{1+t^2}\right)^{1/4} \exp\left[-\frac{x^2}{4(1+t^2)}\right]$$

So that

$$\rho_0(x) = \int |\langle x;0 \rangle|^2 = (2)^{-1/2} \exp\left[-\frac{x^2}{2}\right] \quad (75)$$

$$\langle x;t \rangle = \int |\langle x;t \rangle|^2 = \left[\frac{2}{1+t^2}\right]^{1/2} \exp\left[-\frac{x^2}{2(1+t^2)}\right]$$

and the Fokker-Planck equation (easily derivable from the standard continuity equation $\partial_t \rho = -\nabla \cdot (v \rho)$; $v(x;t) = xt/(1+t^2)$) holds true:

$$\partial_t \rho = -\frac{1}{4} \nabla \cdot (b \rho) ; \quad b(x;t) = \frac{1-t}{1+t^2} x \quad (76)$$

The Madelung factorization $\psi = \exp(R + iS)$ implies (notice that $v = 2R/S$ and $b = 2R/(R + S)$) that the related real functions $\langle x;t \rangle = \exp[R(x;t) + S(x;t)]$ and $\rho(x;t) = \exp[2R(x;t) - S(x;t)]$ read:

$$\begin{aligned} \langle x;t \rangle &= \left[\frac{2}{1+t^2}\right]^{1/4} \exp\left(-\frac{x^2}{4} \frac{1-t}{1+t^2} - \frac{1}{2} \arctan t\right) \\ \rho(x;t) &= \left[\frac{2}{1+t^2}\right]^{1/2} \exp\left(-\frac{x^2}{4} \frac{1+t}{1+t^2} + \frac{1}{2} \arctan t\right) \end{aligned} \quad (77)$$

They solve a suitable version of the general parabolic equations (5), namely :

$$\partial_t \rho = -\frac{1}{4} \rho + c \quad (78)$$

$$\partial_t \langle x;t \rangle = -\frac{1}{4} \langle x;t \rangle + c$$

with

$$c(x;t) = \frac{x^2}{2(1+t^2)^2} - \frac{1}{1+t^2} = 2 \frac{4-t^2}{(1+t^2)^2} \quad (79)$$

Anticipating further discussion, let us mention that the Feynman-Kac kernel, in this case, is a fundamental solution of the time adjoint system (78). For clarity of exposition, let us recall that a fundamental solution of the forward parabolic equation (5) is a continuous function $k(y;s;x;t)$, defined for all $x,y \in \mathbb{R}$ and all $0 \leq s < t \leq T$, which has the following two properties:

(a) for any fixed $(y;s) \in \mathbb{R} \times (0;T)$, the function $(x;t) \mapsto k(y;s;x;t)$ is a regular (i.e. continuous and continuously differentiable the needed number of times) solution of the forward equation (5) in $\mathbb{R} \times (s;T]$

(b) for all continuous functions $\phi(x)$ with a compact support, there holds $\lim_{(t,x) \rightarrow (s,z)} \int_{\mathbb{R}} k(y;s;x;t) \phi(y) dy = \phi(z)$.

First, we need to verify (this will be done self-explanatorily) that $c(x;t)$, (79), is Hölder continuous of exponent one on every compact subset of $\mathbb{R} \times [0;T]$. It follows from direct estimates:

$$\begin{aligned} |c(x_2;t_2) - c(x_1;t_1)| &\leq \frac{1}{2} \left| \frac{x_2^2}{(1+t_2^2)^2} - \frac{x_1^2}{(1+t_1^2)^2} \right| + \left| \frac{1}{1+t_2^2} - \frac{1}{1+t_1^2} \right| \\ &\leq \frac{1}{2} \left| \frac{x_2}{1+t_2^2} - \frac{x_1}{1+t_1^2} \right| + |t_2 - t_1| \left| \frac{t_1}{(1+t_1^2)(1+t_2^2)} + \frac{t_2}{(1+t_1^2)(1+t_2^2)} \right| \end{aligned} \quad (80)$$

But, in case of $|x_1|, |x_2| \leq K$ and $|t_1|, |t_2| \leq T$ we have

$$|c(x_2;t_2) - c(x_1;t_1)| \leq K \left| \frac{x_2}{1+t_2^2} - \frac{x_1}{1+t_1^2} \right| + 2T |t_2 - t_1| \quad (81)$$

Furthermore:

$$\begin{aligned} \left| \frac{x_2}{1+t_2^2} - \frac{x_1}{1+t_1^2} \right| &\leq \left| \frac{x_2 - x_1}{(1+t_2^2)(1+t_1^2)} \right| + \left| \frac{x_2 t_1^2 - x_1 t_2^2}{(1+t_2^2)(1+t_1^2)} \right| \\ &\leq |x_2 - x_1| + T^2 |x_2 - x_1| + 2KT |t_2 - t_1| \end{aligned} \quad (82)$$

implies (the new constant C majorizes all remaining ones)

$$|c(x_2;t_2) - c(x_1;t_1)| \leq C (|x_2 - x_1| + |t_2 - t_1|) \sqrt{\frac{1}{2} C [(x_2 - x_1)^2 + (t_2 - t_1)^2]}^{1/2} \quad (83)$$

Let us also notice that we can introduce an auxiliary function $h(x;t) = \arctan t$ such that there holds

$$4h - c(x;t)h - \partial_t h = \frac{x^2 h(x;t)}{2(1+t^2)^2} \leq 0 \quad (84)$$

We have thus satisfied the crucial assumptions I and II of Ref. [23]. As a consequence, we have granted the existence of a fundamental solution $k(y;s;x;t) \geq 0$. Moreover, for every bounded and continuous function $\phi(x); |\phi(x)| \leq C$, where $C > 0$ is arbitrary, the function

$$u(x;t) = \int_1^{Z+1} k(y;0;x;t) \phi(y) dy \quad (85)$$

is a solution of the Cauchy problem, i.e. solves (79) under the initial condition $u(x;0) = \phi(x)$, so that $|u(x;t)| \leq C$. All that implies the uniqueness of the fundamental solution $k(y;s;x;t)$, and in view of $c(x;t) \geq 1$ its strict positivity. The function $k(y;s;x;t)$ is also a solution of the adjoint equation with respect to variables $y;s$: $\partial_s k = -\Delta_y k + c(y;s)k$ in $R \times [0;T)$. It is obvious that the Chapman-Kolmogorov composition rule holds true, in view of the validity of the Feynman-Kac representation in the present case.

Basically, we must be satisfied with the Feynman-Kac representation of the fundamental solution, whose existence we have granted so far. In our case, the so called parametrix method, [21], can be used to construct fundamental solutions. In fact, since $c(x;t)$ is locally Lipschitz i.e. Hölder continuous of exponent one and quadratically bounded $|c(x;t)| \leq x^2 + 1$, the infinite series:

$$k(y;s;x;t) = \sum_{n=0}^{\infty} (-1)^n k_n(y;s;x;t) \quad (86)$$

where $k_0(y;s;x;t) = [4(t-s)]^{-1/2} \exp[-(x-y)^2/4(t-s)]$ is the heat kernel and

$$k_n(y;s;x;t) = \int_s^t \int_1^{Z+1} dz c(z;\tau) k_{n-1}(y;s;z;\tau) k_0(z;\tau;x;t) \quad (87)$$

are known to converge for all $x,y \in R, 0 \leq s < t < T$, and $t-s < T_0$ where $T_0 < T$, and define the fundamental solution, [30].

By putting $p(y;s;x;t) = k(y;s;x;t) \frac{c(x;t)}{c(y;s)}$ we arrive at the fundamental solution of the second Kolmogorov (Fokker-Planck) equation

$$\partial_t p(y;s;x;t) = -\Delta_x p(y;s;x;t) - r_x [b(x;t)p(y;s;x;t)] \quad (88)$$

where $b = 2x$ and $r = uv$, and in particular $p = p_j = p_j^2$, are consistently propagated by p . It is the transition probability density of the Nelson diffusion associated with the solution (74) of the Schrödinger equation, and at the same time a solution of the first Kolmogorov (backward diffusion) equation

$$\partial_s p(y;s;x;t) = -\Delta_y p(y;s;x;t) - b(y;s)r_y p(y;s;x;t) \quad (89)$$

Equations (88), (89) prove that the pertinent process is a diffusion: it has the standard local (in infinitesimal) characteristics of the diffusion process, [26].

Obviously, the above definition of p in terms of k induces the validity of the compatibility condition

$$c(x;t) = 2 [\partial_t \ln \quad (x;t) + \frac{1}{2} [\frac{b^2(x;t)}{2} + r b(x;t)]] \quad (90)$$

connecting the drift of the diffusion process with the Feynman-Kac potential governing its local dynamics: cf. Refs. [29, 8] and [31] where the Ehrenfest theorem analogue was formulated for general (non-quantal included) Markovian diffusions.

Let us point out that our quantumly motivated example was chosen not to show up a typical for quantum wave functions property of vanishing somewhere. In fact, because of restricting our considerations to strictly positive Feynman-Kac kernels and emphasizing the uniqueness of solutions, we have left aside an important group of topics pertaining to solution of the Schrodinger boundary data problem when:

- (i) the boundary densities have zeros
- (ii) the interpolation itself is capable of producing zeros of the probability density, even if the boundary ones have none.

Only the case (i) can be (locally) addressed by means of strictly positive semigroup kernels, however the uniqueness of solution is generally lost in space dimension higher than one, [2, 3, 4]. General existence theorems are available [28, 29] and indicate that one deals with diffusion-type processes in this case, see e.g. also [5, 6, 8].

The case (ii) seems not to be ever considered in the literature, see however [32].

Acknowledgement: Both authors receive a financial support from the KBN research grant No 2 P 302 057 07. We would like to thank Professors John Klauder and Gert Roepstorff for correspondence concerning the differentiability of Feynman-Kac kernels and Jean-Claude Zambrini for reference suggestion.

References

- [1] E. Schrodinger, Ann. Inst. Henri Poincare, 2, (1932), 269
- [2] R. Fortet, J. Math. Pures Appl., 9, (1940), 83
- [3] A. Beurling, Ann. Math. 72, (1960), 189
- [4] B. Jamison, Z. Wahrsch. verw. Geb. 30, (1974), 65
- [5] J. C. Zambrini, J. Math. Phys. 27, (1986), 3207

- [6] J.C.Zambrini, Phys.Rev.A 35, (1987), 3631
- [7] M.Nagasawa, Prob.Theory Relat.Fields, 82,(1989), 109
- [8] Ph.Blanchard, P.Garbaczewski, Phys.Rev.E, 49, (1994), 3815
- [9] P.Garbaczewski, R.Olkiewicz, Phys.Rev.A, 51, (1995), 3445
- [10] P.Garbaczewski, J.R.Klauder, R.Olkiewicz, Phys.Rev.E, 51, (1995), 4114
- [11] B.Simon, "Functional Integration and Quantum Physics", Academic Press, New York, 1979
- [12] M.Freidlin, "Functional Integration and Partial Differential Equations", Princeton University Press, Princeton, 1985
- [13] B.Simon, Bull.Amer.Math.Soc. 7, (1982), 447
- [14] P.Garbaczewski, Phys.Lett.A 178, (1993), 7
- [15] J.Glimm, A.Ja e, "Quantum Physics: A Functional Integral Point of View", Springer-Verlag, Berlin, 1981
- [16] G.Roepstor , "Path Integral Approach to Quantum Physics", Springer-Verlag, Berlin, 1994
- [17] E.B.Dynkin, "Theory of Markov Processes", Pergamon Press, Oxford, 1960
- [18] E.Nelson, "Dynamical Theories of Brownian Motion", Princeton University Press, Princeton, 1967
- [19] N.A.Watson, "Parabolic Equations on an Infinite Strip", Marcel Dekker, New York, 1989
- [20] S.Karlin, H.M.Taylor, "A Second Course in Stochastic Processes", Academic Press, New York, 1981
- [21] A.M.Ilin, A.S.Kalashnikov, O.A.Oleinik, Usp.Mat.Nauk (in Russian), 27, (1962), 65
- [22] D.G.Aronson, P.Besala, Colloq.Math. 18, (1967), 125

- [23] P. Besala, *Ann. Polon. Math.* 29, (1975), 403
- [24] N. A. Watson, *J. London Math. Soc.* (2), 8 (1974), 311
- [25] N. A. Watson, *Archiv. Rational Mech. Anal.* 68, (1978), 121
- [26] W. Horsthemke, R. Lefever, "Noise-Induced Transitions", Springer-Verlag, Berlin, 1984
- [27] E. Nelson, "Quantum Fluctuations", Princeton University Press, Princeton, 1985
- [28] E. Carlen, *Commun. Math. Phys.* 94, (1984), 293
- [29] R. Carmona, in *Taniguchi Symposium*, PMMP, Katata 1985, ed. K. Itô, N. Ikeda, Academic Press, Boston, 1987
- [30] M. Krzyżanski, A. Szybiak, *Lincol. Rend. Sc. s. mat. e nat.* 28, (1959), 26
- [31] P. Garbaczewski, *Phys. Lett. A* 185, (1994), 149
- [32] Ph. Blanchard, P. Garbaczewski, R. Olkiewicz, "Nonnegative Feynman-Kac kernels in Schrodinger's interpolation problem", *subm. for publ.*