

Minimum Decision Cost for Quantum Ensembles

Dorje Brody^{*} and Bernhard Mester[†]

^{*} Blackett Laboratory, Imperial College, South Kensington, London SW 7 2BZ, U.K.

[†] Isaac Newton Institute For Mathematical Science, 20 Clarkson Road, Cambridge, CB3 0EH,
U.K.

(March 24, 2022)

For a given ensemble of N independent and identically prepared particles, we calculate the binary decision costs of different strategies for measurement of polarised spin $1/2$ particles. The result proves that, for any given values of the prior probabilities and any number of constituent particles, the cost for a combined measurement is always less than or equal to that for any combination of separate measurements upon sub-ensembles. The Bayes cost, which is that associated with the optimal strategy (i.e., a combined measurement) is obtained in a simple closed form.

PACS Numbers : 02.50.-r, 02.50.Le, 03.65.Bz

In a problem of experimental design, the task of the experimentalist is to find an optimal observational strategy. Ordinarily, one must choose among different strategies before the data can be obtained, and hence one must perform a preposterior analysis. When the experiment involves a decision among different quantum mechanical states, such an analysis is indeed important, since, unlike the classical case, virtual sampling, i.e., repeated samplings of the same system, are not generally permitted.

There are a number of different approaches for finding an optimal strategy. In the information-theoretic approach, one typically determines the strategy that maximises the mutual information (see, e.g., [1]), but this is generally difficult, owing to the nonlinear

Electronic address: d.brody@tp.ph.ic.ac.uk

[†]Electronic address: b.mester@ic.ac.uk

nature of the Shannon information. In the minimum approach [2], one finds the strategy that minimises the maximum cost (or loss) incurred by the decision among different strategies. When certain a priori knowledge concerning the nature of the state is available, then one may seek a strategy that minimises the expected cost, using a Bayes procedure [2], [3].

In the present Letter, we study the Bayesian approach to a binary decision problem for an ensemble of polarised spin 1/2 particles. First, we briefly introduce the Bayesian approach to quantum hypothesis testing. These notions, developed by Helstrom and others ([4], [5], and [6]), are then applied to obtain the optimal strategy for a Bayes decision between two quantum mechanical pure states, for an ensemble of polarised spin 1/2 particles. In this example, we first study the application of quantum Bayes sequential analysis to the ensemble. The result is then compared with a combined measurement of the entire ensemble, treated as a single composite system. Other strategies consisting of combined measurements of sub-ensembles are also considered. The Bayes solution to the problem demonstrates that the Bayes cost for separate sequential measurements of the individual particles is the same as that of a combined measurement. This result differs from that predicted by Peres and Wootters [7]. Any other strategy turns out to entail a higher expected cost. Nevertheless, we conclude, for the reasons given below, that a combined measurement of the entire ensemble is, in general, an optimal one.

First, consider a decision problem requiring a choice among M hypotheses $H_1; \dots, H_M$ concerning a quantum system. Hypothesis H_k asserts that the density operator of the system is $\hat{\rho}_k$, ($k = 1; \dots, M$), and the prior probability of the j -th state is p_j with

$$\sum_{k=1}^M p_k = 1 : \quad (1)$$

From past experience, one knows that the system is in the j -th state with a relative frequency p_j . The self-adjoint operators $\hat{\rho}_k$ act on the vectors of a Hilbert space H , are non-negative definite, and have unit trace.

A quantum decision strategy is characterised by a probability operator measure (pom) on H , i.e., a set of M non-negative definite self-adjoint operators E_j satisfying

$$\sum_{j=1}^M \pi_j = 1 : \quad (2)$$

If this pom is applied to the system when hypothesis H_k is true, then the conditional probability of choosing hypothesis H_j is given by

$$\Pr(X = j | W = k) = \text{Tr}(\pi_k \hat{\pi}_j) : \quad (3)$$

Here, X denotes the random variable that is to be observed, and W , typically being the parameter, is the unknown state of nature.

Now, let C_{ij} be the cost of choosing hypothesis H_i when H_j is true. Then the expected cost of the observational strategy specified by the pom π_j is [4]

$$C = \sum_{i=1}^M \sum_{j=1}^M C_{ij} \text{Tr}(\hat{\pi}_j R_i) = \text{Tr} \left(\sum_{j=1}^M \pi_j R_j \right) ; \quad (4)$$

where the Hermitian risk operators R_i are defined by

$$R_i = \sum_{j=1}^M C_{ij} \hat{\pi}_j : \quad (5)$$

A set π_j of pom that minimises the cost (4), under the constraints (2), is defined as optimal and the cost is Bayes, i.e., $C = C^*$ (the superscript $*$ here corresponds to the optimal strategy). Necessary and sufficient conditions for the optimality of a pom are known to be [5], [6] the self-adjointness of the operator

$$= \sum_{j=1}^M \pi_j R_j - \sum_{j=1}^M \pi_j R_j \quad (6)$$

and the non-negative definiteness of the operator R_j for all $j = 1; \dots; M$. The minimum expected Bayes cost is thus

$$C^*(\pi; \pi_j) = \text{Tr} : \quad (7)$$

In a simple case where $M = 2$, i.e., for binary decisions, one can easily verify [4] that the optimal pom is projection valued, and the Bayes cost becomes

$$C^*(\pi; \pi_j) = \pi_1 C_{11} + \pi_2 C_{12} + \pi_2 (C_{12} - C_{22}) \sum_{i>0}^X \pi_i ; \quad (8)$$

where λ_i are the eigenvalues of the operator $\hat{C}_2 = \hat{C}_1$, with

$$\lambda_i = \frac{1}{2} \frac{(C_{21} \pm C_{11})}{(C_{12} \pm C_{22})} = \frac{1}{2} : \quad (9)$$

Here and in the sequel, we choose a 0-1 cost structure; $C_{ij} = 1 - \delta_{ij}$, i.e., assign cost 1 to an incorrect decision and 0 to a correct decision. Also, the prior probability for state 1 is given by $p_1 = \frac{1}{2}$, and hence $p_2 = \frac{1}{2}$.

Now, we consider an experiment where a physicist must estimate (decide) the direction of polarisation of a given ensemble of N spin $1/2$ particles, using a Stern-Gerlach (s-g) device. The physicist knows that the particles have been filtered through another s-g device with a magnetic field in the $x-y$ plane at a constant angle θ_1 or θ_2 from the x -axis, and in either case the spin up state has been selected. The physicist can select the orientation angle ϕ of the detector relative to the x -axis. When the particle passes through the field of the detector magnet, the physicist observes either the spin up (head) or spin down (tail) state, whereupon he must decide between the alternatives θ_1 (i.e., the polarisation direction $\theta = \theta_1$) and θ_2 . We do not specify the values of the angles θ_k , but the difference between the two angles is given by $\theta_2 - \theta_1 = \theta$.

First, consider the case where the physicist performs sequential observations of each individual spin $1/2$ particle. Suppose, for simplicity, that $N = 1$. The physicist has to decide, either before or after the observation, whether the particle is polarised in the θ_1 or θ_2 direction. If a decision were to be chosen without any observation, then a Bayes decision against the prior distribution $P(W)$ of W (in this case, $W = 1$ or 2) would be optimal. Suppose that X (spin 'up' or 'down') is observed before a decision is chosen. Then, the decision process for the physicist follows the same procedure as the previous case. However, the difference here is that the distribution of W has changed from the prior to the posterior distribution. Hence, a Bayes decision against the posterior distribution of W is now optimal.

The conditional probability for observing the spin up (+1) state, when the state of the system is θ_k , is given by

$$p_k(+1) = \Pr(X = +1 | W = \theta_k) = \cos^2\left(\frac{\theta_k}{2}\right) : \quad (10)$$

If one fixes the angle θ , then the experiment is entirely analogous to a classical coin tossing problem [8], with coins whose bias is given by the above b_k . However, having the freedom to choose the angle θ for each value of the prior p , the physicist must choose an optimal direction given by [9]

$$\theta_{\text{opt}}(\theta) = \tan^{-1} \frac{\sin \theta_1 (1 - \cos \theta_2) \sin \theta_2}{\cos \theta_1 (1 - \cos \theta_2) \cos \theta_2} : \quad (11)$$

Hence, we have a problem of tossing quantum coins whose bias is a function of the prior probability p .

Having chosen the optimal angle θ_{opt} , the Bayes decision rule specifies that σ_1 is to be chosen if the spin up state is observed, and σ_2 otherwise. The Bayes cost against the prior p , when $N = 1$, can easily be obtained by calculating the eigenvalues of $\hat{C}_2 - \hat{C}_1$, with the result [9]

$$C(\theta; 1) = \frac{1}{2} \left[1 - \sqrt{2^2 (2 + \cos 2\theta) + 1} \right] : \quad (12)$$

Now, suppose that $N = 2$, and the result of measurement of the first particle has been obtained. As mentioned above, the physicist must follow the same procedures as in the case $N = 1$, with the posterior distribution $p(\theta)$ instead of the prior p . From Bayes' theorem, the posterior probability that $\sigma = \sigma_1$ is given by

$$p(+|\theta) = \frac{b_1(\theta)}{b_1(\theta) + b_2(\theta) (1 - \cos \theta)} \quad (13)$$

or

$$p(-|\theta) = \frac{(1 - b_1(\theta))}{(1 - b_1(\theta)) + (1 - b_2(\theta)) (1 - \cos \theta)} ; \quad (14)$$

according to the outcome (+ or -) of the first measurement. The optimal orientation angle, before performing the second measurement, is now given by $\theta_{\text{opt}}(+|\theta)$ or $\theta_{\text{opt}}(-|\theta)$, accordingly. The Bayes cost for this case ($N = 2$) is given by the weighted average, i.e.,

$$C = b_1 C(\theta(+); 1) + b_2 (1 - b_1) C(\theta(-); 1) :$$

Next, we consider an arbitrary number N of particles. Again, the procedures are the same as above, except that the prior is now replaced by one of the 2^{N-1} posteriors $[(+ + \dots +)]$, after observations of $N-1$ particles. In a classical Bayes decision procedure [2], [3], it is difficult (or impossible) to obtain the Bayes cost as a closed function of N . The reason is that, first, one must study the tree [10] of the posterior distributions, with branches proliferating as 2^N . To each branch (i.e., posterior) of the tree, one associates the cost $C(\cdot; 1)$, and then calculates the weight (probability) for the sequence of outcomes associated with that branch. After these considerations, one can, in principle, obtain the weighted average of the cost, which involves 2^{N-1} terms. (Note that, for classical coins, the branches of the posterior tree do recombine and hence proliferate as N . However, the weights associated with the branches do not recombine, and therefore one cannot avoid the consideration of 2^{N-1} terms.)

In the case of our "quantum coins", the situation appears even worse, since, after each observation, the physicist must turn the device in accordance with formula (11). This results in changing the bias $b_k(\cdot)$ of the "coins" at each stage, and hence one must also incorporate the bias tree (which proliferates 2^N). However, it turns out that this optimal orientation forces the posterior tree to recombine into two branches, i.e.,

$$(n; \cdot) = \frac{1}{2} \left[1 - \frac{q}{1 - 4(1 - \cdot) \cos^2(n-1)} \right]; \quad (15)$$

where \cdot corresponds to the outcome of the last $(n-1)$ -th trial being spin up (+) or down (-). This result can be proven by induction as follows. First, for $n=1$, it is easily verified that $(1; \cdot) = \cdot$ as given in (13) and (14). Next, assume that the last $(n-1)$ -th outcome of the trial is \cdot , and that the posterior is given by the above $(n; \cdot)$. Then, if the next trial outcome is (+), follows from Bayes' theorem, that the posterior distribution, after $n+1$ observations, is given by

$$(\cdot +) = \frac{b_1(\cdot) (n; \cdot)}{b_1(\cdot) (n; \cdot) + b_2(\cdot) (1 - (n; \cdot))};$$

with $\cdot = \text{opt}(\cdot)$. After some algebra, one can show that the above $(\cdot +) = (n+1; +)$. The other three cases $[(\cdot -), \text{etc.}]$ can also be treated in the same manner.

Although the weights for different branches neither recombine in the quantum case, since $C(n; +; 1) = C(n; -; 1)$, the final average cost is just $C(n; -; 1)$ times the sum of all the different weights (which is just 1), and hence we finally deduce that the Bayes cost for sequential observations is

$$\begin{aligned} C(-; N) &= C(-(N-1; -); 1) \\ &= \frac{1}{2} \left(1 - \frac{1}{2^{N-1}} \right) \cos^{2N} \theta; \end{aligned} \quad (16)$$

for either value of the $N-1$ -th outcome (+ or -).

Next, consider the case where the physicist treats the entire ensemble as a single composite system. The total spin of a system with N particles is just $N/2$, and the density operator for a spin $N/2$ particle polarised in the direction $\mathbf{n} = (\cos \theta; \sin \theta; 0)$ is given by

$$(\hat{\rho})_{mn} = \frac{1}{2^{N/2}} \sum_N C_m C_n e^{i(m-n)\phi}; \quad (17)$$

where $(n; m) = 0; \dots; N$. According to the result in (8), one must find the eigenvalues of the matrix $\hat{\rho}_2 - \hat{\rho}_1$ in order to obtain the Bayes cost. We first show that the matrix $\hat{\rho}_2 - \hat{\rho}_1$ is of rank two, and thus has only two non-zero eigenvalues. Define two vectors $\mathbf{u} = (u_n)$ and $\mathbf{v} = (v_n)$ by

$$u_n = \frac{1}{2^{N/2}} \sum_N C_n e^{in\phi_1}; \quad (18)$$

and

$$v_n = \frac{1}{2^{N/2}} \sum_N C_n e^{in\phi_2}; \quad (19)$$

Then, $(\hat{\rho}_1)_{mn} = u_m u_n$ and $(\hat{\rho}_2)_{mn} = v_m v_n$. Since the inner product $\mathbf{u} \cdot \mathbf{u} = \mathbf{v} \cdot \mathbf{v} = 1$, one obtains

$$\hat{\rho}_1 \mathbf{u} = \sum_n (\hat{\rho}_1)_{mn} u_n = \mathbf{u}$$

and similarly, $\hat{\rho}_2 \mathbf{v} = \mathbf{v}$. Now, let \mathbf{w} and λ be an eigenvector and the corresponding eigenvalue of the matrix $\hat{\rho}_2 - \hat{\rho}_1$, i.e.,

$$(\hat{\lambda}_1 \quad \hat{\lambda}_2)w = w : \quad (20)$$

We may expand the eigenvector w in terms of a basis that contains either u or v , i.e., $w = c_1 u + u_\perp$ or $w = c_2 v + v_\perp$. Here, u_\perp denotes some vector orthogonal to u , and similarly for v_\perp . However, since $\hat{\lambda}_1 u_\perp = \hat{\lambda}_2 v_\perp = 0$, we have

$$w = c_1 u + c_2 v : \quad (21)$$

Therefore, the matrix $\hat{\lambda}_2 \quad \hat{\lambda}_1$ is of rank two, as claimed. On the other hand, if we form the inner product of the two vectors $w = c_1 u + u_\perp$ and u , we obtain

$$w \cdot u = \langle w | u \rangle = \frac{c_1}{2} - c_2 (v \cdot u) ; \quad (22)$$

and similarly,

$$w \cdot v = \langle w | v \rangle = \frac{c_1}{2} (u \cdot v) - c_2 : \quad (23)$$

Without any loss of generality, we may now set $c_1 = 1$, and then by eliminating c_2 from the above equations, we obtain the eigenvalues of the matrix $\hat{\lambda}_2 \quad \hat{\lambda}_1$, i.e.,

$$= \frac{1}{2} (1 \pm \sqrt{(1 \pm \frac{1}{2})^2 - 4 (\frac{1}{2} - 1) }) ; \quad (24)$$

where

$$\begin{aligned} \frac{1}{2} &= (v \cdot u) (u \cdot v) \\ &= 2^{-N} \sum_{m=0}^N C_m e^{2im} = \cos^{2N} \left(\frac{\pi}{4} \right) : \end{aligned} \quad (25)$$

Therefore, the binary Bayes decision cost for a spin $N=2$ particle is

$$C(\cdot; N) = \frac{1}{2} \left(1 \pm \sqrt{1 - 4 (\frac{1}{2} - 1) \cos^{2N} \left(\frac{\pi}{4} \right)} \right) : \quad (26)$$

One immediately observes that the above cost (26) is the same as that obtained from sequential analysis, given by (16). Hence, the Bayes solution to our optimisation problem states that a combined measurement is as advantageous as sequential measurements. These two

strategies, however, are not the only ones, and many other partially combined measurement procedures are possible. However, in the present formalism of sequential analysis, the only effect of any intermediate measurements, either partially combined or not, consists in updating the posterior distributions. Since the Bayes cost is a monotonically decreasing function of the number of updating steps, this implies that any partially combined measurements will increase the cost. Therefore, we may now conclude that the optimal measurement strategy consists in either performing a combined measurement of the entire ensemble or performing sequential measurements of the individual particles. Any other strategies will result in higher costs.

This result is quite different from that expected by Peres and Wootters, who conjectured that sequential measurements can never be as efficient as a combined measurement [7]. However, it is important to note that their conjecture is based upon an information-theoretic approach, and the solution of an optimisation problem using a Bayesian approach can yield a different result. Massar and Popescu [11], on the other hand, have proved the above mentioned conjecture explicitly for the case $N = 2$. The method used therein is effectively similar to a Bayesian approach, without the use of the prior distributions. However, when a prior distribution is available, the Bayes solution is known to be optimal in general [2]. If prior knowledge is not available, one can still apply the Bayesian approach, using a non-informative prior. The analysis of such cases is, however, beyond the scope of the present Letter.

Throughout the present Letter, we have only considered the cost associated with making decisions. In any practical situation, on the other hand, one must take into consideration other costs (e.g., the observational cost, the cost of analysing the results, etc.). In our example of sequential analysis, for example, at each stage before performing an observation, the physicist must analyse the previous results in order to determine the optimal turning angle. One might argue that [2] the analysing cost can be ignored, since, after all, scientists are so underpaid that the cost of their labors is usually negligible! Nonetheless, the observational costs cannot be ignored in general. Assuming the linearity of the utility function (e.g., that

the total cost is just the sum of the decision cost and the observational costs), it is clear that any separate measurements will result in a higher total cost, since the decision cost for optimal sequential measurements (i.e., sequential measurements with optimal angular orientations) can never be lower than that for a combined measurement. Therefore, we conclude, after these considerations, that a combined measurement is optimal in general.

In connection with the decision problem for classical coins which was briefly mentioned above, it is interesting to note that all the quantum results obtained by calculating the eigenvalues of the density operators can, in principle, be recovered from purely classical calculations, even for sequential measurements, if and only if the spins of the particles concerned are $1/2$. That is, provided one does not perform any combined measurements, the results can be obtained from classical calculations. More details of this, as well as a treatment including the observational costs, may be found in [8]. (See, also [9] for a comparison between classical and quantum coin tossings.)

The authors acknowledge their gratitude to J. T. Key, and J. D. Malley for useful discussions of the foregoing topics.

1. Davies, E. B., IEEE Trans. Inform. Theory. IT-24, 596 (1978).
2. Berger, J. O., Statistical Decision Theory and Bayesian Analysis (Springer-Verlag, New York, 1985).
3. DeGroot, M. H., Optimal Statistical Decisions (McGraw-Hill, New York, 1970).
4. Helstrom, C. W., Quantum Detection and Estimation Theory (Academic Press, New York, 1976).
5. Holevo, A. S., J. Multivar. Anal. 3, 337 (1973).
6. Yuen, H. P., Kennedy, R. S., and Lax, M., IEEE Trans. Inform. Theory. IT-21, 125 (1975).
7. Peres, A. and Wootters, W. K., Phys. Rev. Lett. 66, 1119 (1991).

8. Brody, D. and M. Eiser, B., "Bayesian Inference in Quantum Systems", Imperial College Preprint (1995), submitted to Physica A.
9. Alley, J. D. and Hornstein, J., Statist. Sci. 8, 433 (1993).
10. Lindley, D. V., Making Decisions (Wiley, London, 1971).
11. Massar, S. and Popescu, S., Phys. Rev. Lett. 74, 1259 (1995).