

# ALTERNATIVE DERIVATION OF THE HU-PAZ-ZHANG MASTER EQUATION OF QUANTUM BROWNIAN MOTION

J. J. Halliwell and T. Yu<sup>y</sup>

Theoretical Physics Group, Blackett Laboratory, Imperial College, London SW 7 2BZ, U.K.

(June, 1995)

## Abstract

Hu, Paz and Zhang [B.L. Hu, J.P. Paz and Y. Zhang, Phys. Rev. D 45 (1992) 2843] have derived an exact master equation for quantum Brownian motion in a general environment via path integral techniques. Their master equation provides a very useful tool to study the decoherence of a quantum system due to the interaction with its environment. In this paper, we give an alternative and elementary derivation of the Hu-Paz-Zhang master equation, which involves tracing the evolution equation for the Wigner function. We also discuss the master equation in some special cases.

PACS Numbers : 05.40.+j, 42.50.Lc, 03.65.Bz

Typeset using REVTeX

---

E-mail: jhalliwell@ic.ac.uk

<sup>y</sup>E-mail: ting.yu@ic.ac.uk

## I. INTRODUCTION

Quantum Brownian motion (QBM) models provide a paradigm of open quantum systems that has been very useful in quantum measurement theory [1], quantum optics [2] and decoherence [3-5]. One of the advantages of the QBM models is that they are reasonably simple, yet sufficiently complex to manifest many important features of realistic physical processes.

Central to the study of QBM is the master equation for the reduced density operator of the Brownian particle, derived by tracing out the environment in the evolution equation for the combined system plus environment. A variety of such derivations have been given [6-9]. The most general is that of Hu, Paz and Zhang [10,11], who used path integral techniques and in particular, the Feynman-Vernon influence functional.

The purpose of this paper is to provide an alternative and elementary derivation of the Hu-Paz-Zhang master equation for QBM, by tracing the evolution equation for the Wigner function of the whole system.

## II. MASTER EQUATION FOR QUANTUM BROWNIAN MOTION

The system we considered is a harmonic oscillator with mass  $M$  and bare frequency  $\omega_0$ , in interaction with a thermal bath consisting of a set of harmonic oscillators with mass  $m_n$  and natural frequency  $\omega_n$ . The Hamiltonian of the system plus environment is given by

$$H = \frac{p^2}{2M} + \frac{1}{2}M\omega_0^2 q^2 + \sum_n \left[ \frac{p_n^2}{2m_n} + \frac{1}{2}m_n\omega_n^2 q_n^2 + q_n \sum_n C_n \right]; \quad (1)$$

where  $q, p$  and  $q_n, p_n$  are the coordinates and momenta of the Brownian particle and oscillators, respectively, and  $C_n$  are coupling constants.

The state of the combined system (1) is most completely described by a density matrix  $\rho(q, q_1; q^0, q_1^0; t)$  where  $q_1$  denotes  $(q_1; \dots; q_N)$ , and evolves according to

$$\dot{\rho} = \frac{i}{\hbar} [H; \rho]; \quad (2)$$

The state of the Brownian particle is described the reduced density matrix, defined by tracing over the environment,

$$\rho_r(q; q^0; t) = \int \prod_n (dq_n dq_n^0) \rho(q; q_i; q^0; q_i^0; t) : \quad (3)$$

The equation of time evolution for the reduced density matrix is called the master equation. For a general environment, Hu, Paz, and Zhang [10] derived the following master equation by using path integral techniques:

$$\begin{aligned} i\hbar \frac{\partial \rho_r}{\partial t} = & \frac{\hbar^2}{2M} \frac{\partial^2 \rho_r}{\partial q^2} - \frac{\partial^2 \rho_r}{\partial q^2} + \frac{1}{2} M \omega^2 (q^2 - q^0) \rho_r \\ & + \frac{1}{2} M \omega^2(t) (q^2 - q^0) \rho_r \\ & + i\hbar \gamma(t) (q - q^0) \frac{\partial \rho_r}{\partial q} - \frac{\partial \rho_r}{\partial q^0} \\ & + iM \gamma(t) \hbar(t) (q - q^0)^2 \rho_r \\ & + \hbar \gamma(t) f(t) (q - q^0) \frac{\partial \rho_r}{\partial q} + \frac{\partial \rho_r}{\partial q^0} : \end{aligned} \quad (4)$$

The explicit form of the coefficients of the above equation will be given later on. The coefficient  $\omega^2(t)$  is the frequency shift term, the coefficient  $\gamma(t)$  is the "quantum dissipative" term, and the coefficients  $\gamma(t)\hbar(t)$ ;  $\gamma(t)f(t)$  are "quantum diffusion" terms. Generally, these coefficients are time dependent

and of quite complicated behaviour.

We find it convenient to use the Wigner function of the reduced density matrix,

$$\tilde{W}(q; p; t) = \frac{1}{2} \int du e^{iup} \rho_r(q - \frac{u}{2}; q + \frac{u}{2}; t) : \quad (5)$$

Taking the Wigner transform of (4), we obtain<sup>1</sup>

$$\begin{aligned} \frac{\partial \tilde{W}}{\partial t} = & \frac{1}{M} p \frac{\partial \tilde{W}}{\partial q} + M [\omega^2 + \omega^2(t)] q \frac{\partial \tilde{W}}{\partial p} \\ & + 2\gamma(t) \frac{\partial (p\tilde{W})}{\partial p} + \hbar M \gamma(t) \hbar(t) \frac{\partial^2 \tilde{W}}{\partial p^2} \\ & + \hbar \gamma(t) f(t) \frac{\partial^2 \tilde{W}}{\partial q \partial p} : \end{aligned} \quad (6)$$

---

<sup>1</sup>We believe Eq. (2.48) in Ref. [10] contains some incorrect numerical factors.

The inverse transformation of (5) is given by

$$\rho(q; q^0; t) = \int dp e^{ip(q - q^0) - \hbar W \frac{q + q^0}{2}; p; t} \quad (7)$$

Our strategy for deriving the master equation (4) is to derive the Fokker-Planck type equation (6) from the Wigner equation for the total system. The master equation can be obtained from the Wigner equation for the system by using the transformation (7).

We shall make the following two assumptions:

(1) The system and the environment are initially uncorrelated, i.e. the initial Wigner function factors

$$W_0(q; p; q_b; p_b) = W_0^s(q; p) W_0^b(q_b; p_b); \quad (8)$$

where  $W_0^s$  and  $W_0^b$  are the Wigner functions of the system and the bath, respectively, at  $t = 0$ .

(2) The heat bath is initially in a thermal equilibrium state at temperature  $T = (k_B)^{-1}$ . This means that the initial Wigner function of bath is of Gaussian form,

$$\begin{aligned} W_0^b &= \prod_n W_{n0}^b \\ &= \prod_n N_n \exp \left\{ -\frac{2}{\hbar} \tanh \left( \frac{1}{2} \hbar \beta_n \right) H_n g \right\}; \end{aligned} \quad (9)$$

where  $H_n$  is the Hamiltonian of the  $n$ -th oscillator in the bath,

$$H_n = \frac{p_n^2}{2m_n} + \frac{1}{2} m_n \omega_n^2 q_n^2; \quad (10)$$

In addition, one easily sees that the initial moments of the bath are

$$\langle q_n(0) \rangle = \langle p_n(0) \rangle = 0; \quad (11)$$

$$\langle q_n(0) q_m(0) \rangle = 0 \quad (\text{if } m \neq n); \quad (12)$$

$$\langle p_n(0) p_m(0) \rangle = 0 \quad (\text{if } m \neq n); \quad (13)$$

$$\langle q_n(0) p_m(0) + p_m(0) q_n(0) \rangle = 0; \quad (14)$$

and

$$\begin{aligned} \langle q_n^2(0) \rangle &= \frac{\hbar}{2m_n \omega_n} \coth\left(\frac{1}{2}\hbar\omega_n\right); \\ \langle p_n^2(0) \rangle &= \frac{1}{2}\hbar m_n \omega_n \coth\left(\frac{1}{2}\hbar\omega_n\right); \end{aligned} \quad (15)$$

For the QBM problem described by (1) and (2), the Wigner function of the combined system plus environment satisfies

$$\begin{aligned} \frac{\partial W}{\partial t} = & \frac{p}{M} \frac{\partial W}{\partial q} + M^{-2} q \frac{\partial W}{\partial p} \\ & + \sum_n \frac{p_n}{m_n} \frac{\partial W}{\partial q_n} + m_n^{-2} q_n \frac{\partial W}{\partial p_n} \\ & + \sum_n C_n \left( q_n \frac{\partial W}{\partial p} + p_n \frac{\partial W}{\partial q_n} \right); \end{aligned} \quad (16)$$

By integrating over the bath variables on the both sides of the above equation, one obtains

$$\frac{\partial \tilde{W}}{\partial t} = \frac{p}{M} \frac{\partial \tilde{W}}{\partial q} + M^{-2} q \frac{\partial \tilde{W}}{\partial p} + \sum_n C_n \int \prod_i dq_i dp_i q_i \frac{\partial \tilde{W}}{\partial p}; \quad (17)$$

where  $\tilde{W}(q;p)$  is the reduced Wigner function and it follows from Eq. (3) that

$$\tilde{W}(q;p) = \int \prod_i dq_i dp_i W(q;p;q_i;p_i); \quad (18)$$

This definition is equivalent to Eqs. (3) and (5). The first two terms on the right-hand side of the Eq. (17) give rise to the standard evolution equation of the system. The last term contains all the information about the behaviour of the system in the presence of interaction with environment.

In what follows, we shall demonstrate that the quantity

$$G(q;p) = \sum_n C_n \int \prod_i dq_i dp_i q_i W \quad (19)$$

appearing (differentiated with respect to  $p$ ) in (17) can be expressed in terms of  $\tilde{W}$  and its derivatives. To this end, we first perform Fourier transform of  $G(q;p)$

$$\begin{aligned} G(k;k^0) &= \int \prod_i dq_i dp_i e^{ikq + ik^0 p} G(q;p) \\ &= \sum_n C_n \int \prod_i dq_i dp_i q_i e^{ikq + ik^0 p} \tilde{W}(q;p;q_i;p_i); \end{aligned} \quad (20)$$

It is well known that  $q(t); p(t)$  and  $q_h(t); p_h(t)$  are related to the classical evolution of their initial values  $q(0); p(0)$  and  $q_h(0); p_h(0)$  through a canonical transformation:

$$z(t) = U(t)z(0); \quad (21)$$

where

$$z(t) = (q(t); q_1(t); \dots; q_N(t); p(t); p_1(t); \dots; p_N(t)) :$$

Since the Hamiltonian (1) is quadratic, the Eq. (16) has the same form as the classical Liouville equation, so the solution of Eq. (16) is of the form ,

$$W_t(z) = W_0(U^{-1}(t)z) : \quad (22)$$

Changing the integration variables into their initial values by this canonical transformation, we obtain

$$\begin{aligned} G(k; k^0) = & \int \prod_i dq_i(0) dp_i(0) \int \prod_n dq_n(0) dp_n(0) \\ & \exp \{ ik [q(0) + p(0)] + \sum_n (f_n q_n(0) + g_n p_n(0)) \} \\ & \exp \{ ik [M q(0) + p(0)] + \sum_n (a_n q_n(0) + b_n p_n(0)) \} \\ & W_0^s(q(0); p(0)) W_0^b(q_h(0); p_h(0)) : \end{aligned} \quad (23)$$

Here the coefficients  $f; g; f_n; g_n; ; a_n; b_n$  are time dependent. Their explicit values are not required.

Similarly, the Fourier transform of the reduced Wigner function is

$$\begin{aligned} \tilde{W}(k; k^0) = & \int dq dp e^{ikq + ik^0 p} \tilde{W}(q; p) \\ = & \int \prod_i dq_i(0) dp_i(0) \int \prod_n dq_n(0) dp_n(0) \\ & \exp \{ ik [q(0) + p(0)] + \sum_n (a_n q_n(0) + b_n p_n(0)) \} \\ & \exp \{ ik [M q(0) + p(0)] + \sum_n (a_n q_n(0) + b_n p_n(0)) \} \\ & W_0^s(q(0); p(0)) W_0^b(q_h(0); p_h(0)) : \end{aligned} \quad (24)$$

Now compare  $G(k; k^0)$  and  $\tilde{W}(k; k^0)$ . They differ by the terms linear in  $q(0); p(0); q_h(0); p_h(0)$  in the preexponential factor in  $G(k; k^0)$ . Consider the factors  $f_h q_h(0)$  and  $g_h p_h(0)$  in  $G(k; k^0)$ . Since they multiply  $W_0^b(q_h(0); p_h(0))$ , and since  $W_0^b(q_h(0); p_h(0))$  is Gaussian in  $q_h(0); p_h(0)$ , the terms  $f_h q_h(0) W_0^b$  and  $g_h p_h(0) W_0^b$  may be replaced by terms of the form  $\partial W_0^b / \partial q_h(0); \partial W_0^b / \partial p_h(0)$  up to time dependent factors. An integration by parts then may be performed, and these factors are then effectively replaced by multiplicative factors of  $k; k^0$ .

Similarly, the factors  $f q(0); g p(0)$  in the prefactor in  $G(k; k^0)$  may be replaced by  $\partial / \partial k; \partial / \partial k^0$  (plus some more factors of  $k$  and  $k^0$ ). Hence, it is readily seen that  $G(k; k^0)$  is a linear combination of terms of the form  $k; k^0; \partial / \partial k; \partial / \partial k^0$  operating on  $\tilde{W}(k; k^0)$ , with time dependent coefficients.

Inverting the Fourier transform, it follows that

$$G = A(t) q \tilde{W} + B(t) p \tilde{W} + C(t) \frac{\partial \tilde{W}}{\partial q} + D(t) \frac{\partial \tilde{W}}{\partial p} : \quad (25)$$

for some coefficients  $A(t); B(t); C(t); D(t)$  to be determined. This result immediately leads to the general form Wigner equation :

$$\begin{aligned} \frac{\partial \tilde{W}}{\partial t} = & \frac{p}{M} \frac{\partial \tilde{W}}{\partial q} + M^{-2} q \frac{\partial \tilde{W}}{\partial p} + A(t) q \frac{\partial \tilde{W}}{\partial p} \\ & + B(t) \frac{\partial (p \tilde{W})}{\partial p} + C(t) \frac{\partial^2 \tilde{W}}{\partial p \partial q} + D(t) \frac{\partial^2 \tilde{W}}{\partial p^2} : \end{aligned} \quad (26)$$

### III. DETERMINATION OF THE COEFFICIENTS (GENERAL CASE)

Having found the functional form of the Wigner equation (26) of the Brownian particle, the next step is to determine the coefficients in the equation. Undoubtedly, there is more than one way to do this. Here we shall choose a way which is both mathematically simple and physically heuristic. Towards this direction, let us consider the time evolution of the expectation values of the system variables:  $q; p; q^2; p^2$  and  $\frac{1}{2}(pq + qp)$ .

By using Eq. (16), we have

$$\frac{d}{dt}h_{qi} = \frac{1}{M}h_{pi}; \quad (27)$$

$$\frac{d}{dt}h_{pi} = -M^{-2}h_{qi} - \sum_n C_n h_{qn}i; \quad (28)$$

$$\frac{d}{dt}h_{q^2i} = \frac{1}{M}h_{pq} + q_{pi}; \quad (29)$$

$$\frac{d}{dt}h_{p^2i} = -M^{-2}h_{pq} + q_{pi} - 2 \sum_n C_n h_{pn}q_i; \quad (30)$$

$$\frac{d}{dt}h_{pq} + q_{pi} = \frac{2}{M}h_{p^2i} - 2M^{-2}h_{q^2i} - 2 \sum_n C_n h_{qn}q_i; \quad (31)$$

Similarly, using Eq. (26) yields

$$\frac{d}{dt}h_{qi} = \frac{1}{M}h_{pi}; \quad (32)$$

$$\frac{d}{dt}h_{pi} = -(M^{-2} + A)h_{qi} - B h_{pi}; \quad (33)$$

$$\frac{d}{dt}h_{q^2i} = \frac{1}{M}h_{pq} + q_{pi}; \quad (34)$$

$$\frac{d}{dt}h_{p^2i} = -(M^{-2} + A)h_{pq} + q_{pi} - 2B h_{p^2i} + 2D; \quad (35)$$

$$\frac{d}{dt}h_{pq} + q_{pi} = \frac{2}{M}h_{p^2i} - 2(M^{-2} + A)h_{q^2i} - B h_{pq} + q_{pi} + 2C; \quad (36)$$

Since the evolution equations of the expectation values are connected to the system variables, the above two sets of equations must be identical.

Now by comparing (28) with (33) we see that

$$\sum_n C_n h_{qn}i = A h_{qi} + B h_{pi}; \quad (37)$$

Similarly, by comparing (31) with (36), (30) with (35), respectively, we get

$$\sum_n C_n h_{qn}q_i = A h_{q^2i} + \frac{B}{2}h_{pq} + q_{pi} - C; \quad (38)$$

$$\sum_n C_n h_{pn}q_i = \frac{A}{2}h_{pq} + q_{pi} + B h_{p^2i} - D; \quad (39)$$

The coefficients  $A; B; C; D$  may now be determined from (37)–(39) by regarding the expectation values  $h_{qi}; h_{qn}q_i$  etc. as expectation values of Heisenberg picture operators, and by solving the operator equation of motion. For simplicity, we still use ordinary notation to represent an operator without adding a hat on it.



The solution to the equation of motion may be written,

$$q_n(t) = q_n(0) + p_n(0)t + \sum_n^X (a_{nm} q_n(0) + b_{nm} p_n(0)) ; \quad (40)$$

$$q(t) = q(0) + p(0)t + \sum_n^X (a_n q_n(0) + b_n p_n(0)) ; \quad (41)$$

for some time-dependent coefficients  $a_n; b_n; a_{nm}; b_{nm}; ; ; a_n; b_n$ . Note that  $q_n(t)$  has been expressed in terms of the final, not initial values of  $q; p$ . By substituting Eq. (40) into Eq. (37), keeping (11) in mind, and comparing the two sides of the resulting equation, we have

$$A = \sum_n^X C_n a_n ; B = \sum_n^X C_n b_n : \quad (42)$$

Similarly, substituting (40) and (41) into (38) and (39), respectively, we get

$$C = \sum_{m,n}^X C_n (a_{nm} a_m \dot{q}_m^2(0) + b_{nm} b_m \dot{p}_m^2(0)) ; \quad (43)$$

$$D = M \sum_{m,n}^X C_n (a_{nm} a_m \dot{q}_m^2(0) + b_{nm} b_m \dot{p}_m^2(0)) : \quad (44)$$

Here we have made use of  $p = M \dot{q}$ . The coefficients  $A; B; C; D$  are therefore completely determined by solving the equation of motion. We now do this explicitly.

We have

$$q(t) + \ddot{q}(t) = \frac{1}{M} \sum_n^X C_n q_n(t) ; \quad (45)$$

$$q_n(t) + \omega_n^2 q_n(t) = \frac{C_n}{m_n} q(t) : \quad (46)$$

The solution to Eq. (46) is as follows:

$$q_n(t) = q_n(0) \cos(\omega_n t) + \frac{p_n(0)}{m_n \omega_n} \sin(\omega_n t) + \sum_0^t ds \frac{\sin[\omega_n(t-s)]}{\omega_n} \frac{C_n}{m_n} q(s) : \quad (47)$$

Combining (45) and (47) gives

$$q(t) + \ddot{q}(t) + \frac{2}{M} \sum_0^t d(t-s) q(s) = \frac{f(t)}{M} ; \quad (48)$$

where

$$f(t) = \sum_n C_n q_n(0) \cos(\omega_n t) + \frac{p_n(0)}{m_n} \frac{\sin(\omega_n t)}{\omega_n} : \quad (49)$$

The kernel  $\chi(s)$  is defined as

$$\chi(s) = \frac{d}{ds} \psi(s) ; \quad (50)$$

where

$$\psi(s) = \sum_0^{Z+1} d! \frac{I(\omega)}{\omega!} \cos(\omega s) : \quad (51)$$

Here  $I(\omega)$  is the spectral density of the environment:

$$I(\omega) = \sum_n (\omega - \omega_n) \frac{C_n^2}{2m_n \omega_n} : \quad (52)$$

In order to get the expressions (40) and (41), we solve equation (48) with the following two different initial conditions:

$$q(s=0) = q(0) ; \quad \dot{q}(s=0) = \frac{p(0)}{M} : \quad (53)$$

and

$$q(s=t) = q(t) ; \quad \dot{q}(s=t) = \frac{p(t)}{M} : \quad (54)$$

where  $t$  is any given time point. In doing so, we consider the elementary functions  $u_i(s)$  ( $i=1,2$ ) introduced by Hu, Paz and Zhang [10] which satisfy the following homogeneous integro-differential equation

$$\ddot{u}(s) + \omega^2 u(s) + \frac{2}{M} \sum_0^Z d(s - \tau) u(\tau) = 0 \quad (55)$$

with the boundary conditions:

$$u_1(s=0) = 1 ; \quad u_1(s=t) = 0 ; \quad (56)$$

and

$$u_2(s=0) = 0; u_2(s=t) = 1 : \quad (57)$$

The solution to equation (55) with the initial condition (53) is obtained as the linear combination of  $u_1; u_2$ ,

$$w(s) = u_1(s) \frac{u_1(0)}{u_2(0)} u_2(s) q(0) + \frac{u_2(s)}{u_2(0)} \frac{p(0)}{M} : \quad (58)$$

The solution to equation (48) with the homogeneous initial conditions can be formally written as

$$w(s) = \frac{1}{M} \int_0^s ds G_1(s; ) f( ) : \quad (59)$$

Where  $G_1(s_1; s_2)$  is the Green function which can be constructed in terms of  $u_i (i=1;2)$ :

$$G_1(s_1; s_2) = \begin{cases} \frac{u_1(s_2)u_2(s_1) - u_2(s_2)u_1(s_1)}{u_1(s_2)u_2(s_2) - u_1(s_2)u_2(s_2)} & \text{if } s_1 > s_2 \\ 0 & \text{otherwise.} \end{cases}$$

Note that  $G_1(s_1; s_2)$  as the function of  $s_1$  satisfies equation (55) with

$$G_1(s_1 = s_2; s_2) = 0; \frac{d}{ds_1} G_1(s_1 = s_2; s_2) = 1 : \quad (60)$$

Then the solution to the equation (48) with initial condition (53) reads

$$q(s) = w(s) + w(s); \quad (61)$$

explicitly,

$$q(s) = u_1(s) \frac{u_1(0)}{u_2(0)} u_2(s) q(0) + \frac{u_2(s)}{u_2(0)} p(0) + \sum_n \frac{C_n}{M} \int_0^s ds G_1(s; ) \cos(\lambda_n) q_n(0) + \sum_n \frac{C_n}{M} \int_0^s ds G_1(s; ) \frac{\sin(\lambda_n)}{\lambda_n} \frac{p_n(0)}{m_n} : \quad (62)$$

It can be shown that the solution to the homogeneous equation (55) with the initial conditions (54) is

$$u(s) = u_2(s) \frac{u_2(t)}{u_1(t)} u_1(s) q(t) + \frac{u_1(s)}{u_1(t)} \frac{p(t)}{M} ; \quad (63)$$

and

$$\alpha(s) = \frac{1}{M} \int_0^s dG_2(s; \tau) f(\tau); (s \leq t) \quad (64)$$

is the solution to the inhomogeneous equation (48) with the homogeneous initial conditions

$$\alpha(t) = 0; \alpha'(t) = 0; \quad (65)$$

where Green function  $G_2(s_1; s_2)$  is

$$G_2(s_1; s_2) = \begin{cases} \frac{u_1(s_2)u_2(s_1) - u_2(s_2)u_1(s_1)}{u_1(s_2)u_2(s_2) - u_1(s_2)u_2(s_2)} & \text{if } s_2 > s_1 \\ 0 & \text{otherwise.} \end{cases}$$

Hence, we get the solution to Eq. (48) with the initial conditions (54)

$$\begin{aligned} q(s) &= u(s) + \alpha(s) \\ &= u_2(s) \frac{u_2(t)}{u_1(t)} u_1(s) q(t) + \frac{u_1(s)}{u_1(t)} \frac{p(t)}{M} \\ &\quad + \sum_n \frac{C_n}{M} \int_0^t ds G_2(s; \tau) \cos(\omega_n \tau) q_n(0) \\ &\quad + \sum_n \frac{C_n}{M} \int_0^t ds G_2(s; \tau) \frac{\sin(\omega_n \tau) p_n(0)}{\omega_n m_n} : \end{aligned} \quad (66)$$

Substituting (66) into (47), one obtains

$$\begin{aligned} q_n(t) &= \frac{C_n}{m_n \omega_n} \int_0^t ds \sin[\omega_n(t-s)] u_2(s) \frac{u_2(t)}{u_1(t)} u_1(s) q(t) \\ &\quad + \frac{C_n}{m_n \omega_n} \int_0^t ds \sin[\omega_n(t-s)] \frac{u_1(s)}{u_1(t)} \frac{p(t)}{M} \\ &\quad + q_n(0) \cos(\omega_n t) + \frac{p_n(0) \sin(\omega_n t)}{m_n \omega_n} \\ &\quad + \frac{1}{M} \sum_m \frac{C_n C_m}{m_n \omega_n} \int_0^t ds \int_0^t ds' \sin[\omega_n(t-s)] G_2(s; s') \cos(\omega_m s') q_m(0) \\ &\quad + \frac{1}{M} \sum_m \frac{C_n C_m}{m_n \omega_n} \int_0^t ds \int_0^t ds' \sin[\omega_n(t-s)] G_2(s; s') \frac{\sin(\omega_m s')}{m_m \omega_m} p_m(0) : \end{aligned} \quad (67)$$

By using (42) we immediately arrive at

$$A(t) = \sum_n \frac{C_n^2}{m_n \omega_n} \int_0^t ds \sin[\omega_n(t-s)] u_2(s) \frac{u_2(t)}{u_1(t)} u_1(s) ; \quad (68)$$

$$B(t) = \frac{1}{M} \sum_n \frac{C_n^2}{m_n \omega_n} \int_0^t ds \sin[\omega_n(t-s)] \frac{u_1(s)}{u_1(t)} : \quad (69)$$

Furthermore,  $A; B$  can be written as

$$A(t) = 2 \int_0^{Z_t} ds (t-s) u_2(s) - 2 \frac{u_2(t)}{u_1(t)} \int_0^{Z_t} ds (t-s) u_1(s); \quad (70)$$

$$B(t) = \frac{2}{M u_1(t)} \int_0^{Z_t} ds (t-s) u_1(s); \quad (71)$$

From (62), the momentum of the Brownian particle is then

$$\begin{aligned} p(s) &= M \dot{q}(s) \\ &= \frac{u_1(s)}{u_1(0)} \frac{u_1(0)}{u_2(0)} u_2(s) - M \dot{q}(0) + \frac{u_2(s)}{u_2(0)} p(0) \\ &\quad \times \int_0^{Z_s} ds G_1^0(s; \cdot) \cos(\cdot) q_1(0) \\ &\quad \times \int_0^{Z_s} ds G_1^0(s; \cdot) \frac{\sin(\cdot)}{\cdot} \frac{p_1(0)}{m_1}; \end{aligned} \quad (72)$$

Here "prime" stands for derivative with respect to the first variable of  $G_1(s; \cdot)$ .

With these results, It can be easily shown that

$$\begin{aligned} C(t) &= \frac{h}{M} \int_0^{Z_t} ds G_1(t; \cdot) (t-s) \\ &\quad \times \int_0^{Z_s} ds \int_0^{Z_s} ds (t-s) G_1(t; \cdot) G_2(s; \cdot) (\cdot); \end{aligned} \quad (73)$$

and

$$\begin{aligned} D(t) &= h \int_0^{Z_t} ds G_1^0(t; \cdot) (t-s) \\ &\quad \times \int_0^{Z_s} ds \int_0^{Z_s} ds (t-s) G_1^0(t; \cdot) G_2(s; \cdot) (\cdot); \end{aligned} \quad (74)$$

where  $\gamma(s)$  is defined as

$$\gamma(s) = \int_0^{Z+1} d! I(!) \coth\left(\frac{1}{2}h!\right) \cos(!s); \quad (75)$$

It is seen that the coefficients  $A(t); B(t); C(t); D(t)$  are dependent only on the kernels  $\gamma(s)$  and  $\gamma(s)$  and the initial state of the bath, not dependent on the initial state of the system. Once the spectral density of the environment is given, the elementary functions  $u_i (i=1;2)$  can be solved from equation (55), the Green functions  $G_i (i=1;2)$  are then obtained. Thus the coefficients of master equation can be determined.

#### IV. PARTICULAR CASES

In this section, we will consider some special cases. Let us at first treat a special case in which we assume that the interaction between the system and environment is weak, so the  $C_n$  are small. In this case, the coefficients are of simple forms, and the determination of these coefficients is very simple and straightforward. We shall work out these coefficients directly using the method in the last section, rather than the general formulae.

The solution to Eq. (46) may be written as

$$q_n(t) = q_n(0) \cos(\omega_n t) + \frac{p_n(0) \sin(\omega_n t)}{m_n \omega_n} + \frac{C_n}{m_n} \int_0^t dt' \frac{\sin[\omega_n(t-t')]}{\omega_n} \cos[\omega_n(t-t')] q(t') + \frac{C_n}{m_n} \int_0^t dt' \frac{\sin[\omega_n(t-t')]}{\omega_n} \frac{\sin[\omega_n(t-t')] p(t')}{M} + O(C_n^2) : \quad (76)$$

Using Eq. (37) and ignoring terms with higher than the second order of  $C_n$ , we get

$$\begin{aligned} \sum_n C_n h q_n(t) i &= \sum_n \frac{C_n^2}{m_n} \int_0^t dt' \frac{\sin[\omega_n(t-t')]}{\omega_n} \cos[\omega_n(t-t')] g h q(t') i \\ &+ \sum_n \frac{1}{M} \frac{C_n^2}{m_n} \int_0^t dt' \frac{\sin[\omega_n(t-t')]}{\omega_n} \frac{\sin[\omega_n(t-t')] p(t')}{\omega_n} g h p(t') i : \end{aligned} \quad (77)$$

Then we immediately get

$$A(t) = 2 \int_0^t ds \langle s \rangle \cos(s) ; \quad (78)$$

$$B(t) = \frac{2}{M} \int_0^t ds \langle s \rangle \sin(s) : \quad (79)$$

We next evaluate  $\sum_n C_n h q(t) q_n(t) i$  and  $\sum_n C_n h p(t) q_n(t) i$ . After a few manipulations, we arrive at the expressions

$$C(t) = \sum_n C_n f h q(t) q_n(0) i \cos(\omega_n t) + h q(t) p_n(0) i \frac{\sin(\omega_n t)}{m_n \omega_n} g ; \quad (80)$$

and

$$D(t) = \sum_n C_n f h p(t) q_n(0) i \cos(\omega_n t) + h p(t) p_n(0) i \frac{\sin(\omega_n t)}{m_n \omega_n} g : \quad (81)$$

To calculate  $C(t)$  and  $D(t)$ , we need to expand  $q(t)$  up to the second order of  $C_n$ :

$$\begin{aligned}
 q(t) = & q(0) \cos(\omega t) + \frac{p(0) \sin(\omega t)}{M} \\
 & + \sum_n \frac{C_n}{M} \int_0^{\omega t} ds \frac{\sin[\omega(t-s)]}{\omega} \cos(\omega_n s) q_n(0) \\
 & + \sum_n \frac{C_n}{M} \int_0^{\omega t} ds \frac{\sin[\omega(t-s)]}{\omega} \frac{\sin(\omega_n s)}{\omega_n} \frac{p_n(0)}{m_n} \\
 & + O(C_n^2) :
 \end{aligned} \tag{82}$$

The expansion of  $p(t)$  is easily obtained from that of  $q(t)$ ,

$$p(t) = M \dot{q}(t) : \tag{83}$$

With these results it is easy to compute  $C(t)$  and  $D(t)$ :

$$C(t) = \frac{h}{M} \int_0^{\omega t} ds \langle s \rangle \sin(\omega s); \tag{84}$$

$$D(t) = h \int_0^{\omega t} ds \langle s \rangle \cos(\omega s); \tag{85}$$

This simple example exhibits the time dependency of the coefficients of the master equation in a general environment. Eqs. (78), (79), (84), (85) are in agreement with Hu, Paz and Zhang [10].

As another example, we briefly discuss the purely Ohmic case in the Fokker-Planck limit (a particular high temperature limit), which has been extensively discussed in the literature [6,10]. In this case one has

$$\langle s - s^0 \rangle = M^{-1} \langle s - s^0 \rangle \tag{86}$$

$$\langle s - s^0 \rangle = \frac{2M k_B T}{h} \langle s - s^0 \rangle \tag{87}$$

Then the equation (55) reduces to

$$u(s) + \frac{\gamma}{2} u(s) + \frac{1}{2} u(s) = -\frac{1}{2} \langle s \rangle u(0) \tag{88}$$

where  $\frac{\gamma}{2} = \frac{1}{2} \langle s^2 \rangle - \frac{1}{2} \langle s \rangle^2$ . After solving this equation, a few calculations give

$$A(t) = 2M \quad (0); \quad (89)$$

$$B(t) = 2; \quad (90)$$

$$C(t) = 0; \quad (91)$$

$$D(t) = 2M \quad k_B T; \quad (92)$$

then the Wigner equation reads:

$$\frac{\partial}{\partial t} \tilde{W} = \frac{p}{M} \frac{\partial \tilde{W}}{\partial q} + M \quad \overset{2}{\text{ren}} q \frac{\partial \tilde{W}}{\partial p} + 2 \frac{\partial \tilde{W}}{\partial p} + 2M \quad k_B T \frac{\partial^2 \tilde{W}}{\partial p^2} : \quad (93)$$

In this regime, the coefficients of this Wigner equation are constants.

#### ACKNOWLEDGEMENTS

We are grateful to Roland Omnès for suggesting this method of deriving the master equation, and encouraging us to pursue it.



## REFERENCES

- [1] W . Zurek, in Proceedings of the NATO Advanced Study Institute on Non-Equilibrium Statistical Mechanics, Sante Fe, 1984 (Plenum , New York, 1984).
- [2] H . Carmichael, An Open Systems Approach to Quantum Optics (Springer-Verlag, 1993).
- [3] W . G . Unruh and W . Zurek, Phys. Rev. D 40, 1071 (1989).
- [4] M . Gell-Mann and J. Hartle, Phys. Rev. D 47, 3345 (1993).
- [5] H F . Dowker and J.J. Halliwell, Phys. Rev. D 46, 1580 (1992).
- [6] A O . Caldeira and A J. Leggett, Physica 121A , 587 (1983).
- [7] H . Dekker, Phys. Rev. A 16, 2116 (1977).
- [8] F . Haake and R . Reibold, Phys. Rev. A 32, 2462 (1985).
- [9] J.P. Paz, in The Physical Origin of Time Asymmetry, edited by J.J. Halliwell, J. Perez-Mercader, and W . Zurek (Cambridge University Press, Cambridge, 1994).
- [10] B L . Hu, J. Paz, and Y . Zhang, Phys. Rev. D 45, 2843 (1992).
- [11] B L . Hu, J. Paz, and Y . Zhang, Phys. Rev. D 47, 1576 (1993).