

IFUG-95-R-4  
 August 22, 1995  
 quant-ph/9509017

# Penning Trap and Vacuum Noise

H. Rosu<sup>1</sup>

Instituto de Física, Universidad de Guanajuato,  
 Apdo Postal E-143, 37150 León, Gto, México

## Abstract

A number of comments are provided on Rogers's model experiment to measure the circular Unruh vacuum noise by means of a hyperbolic Penning trap inside a microwave cavity. It is suggested that cylindrical Penning traps, being geometrically simpler, and controlled almost at the same level of accuracy as the hyperbolic trap, might be a better choice for such an experiment. Besides, the microwave modes of the trap itself, of known analytical structure, can be directly used in trying to obtain measurable results for such a tiny noise effect.

PACS 03.65.Bz - Foundations, theory of measurement, miscellaneous theories  
 PACS 32.80.Pj - Optical cooling of atoms; trapping

The physics of electromagnetic traps is an extremely vigorous discipline, both theoretically and experimentally speaking [1]. Electrons in the Penning trap (PT), also known as electron geoniums and associated mainly with  $g-2$  experiments, are simple physical systems offering many advantages for measuring fundamental quantum field effects. They have an internal dynamic behaviour which is understood in great detail and they can be prepared and fully controlled by means of modern laser spectroscopic methods. The PTs are characterized by very high detection efficiency and high accuracy and tend to become the favorite devices for studying experimentally the theoretical results on radiative interactions, as well as fundamentals of quantum mechanics [2].

On the other hand, vacuum noise is a concept vehiculated by people in quantum field theory [3]. If a quantum particle is following a classical trajectory endowed with an acceleration parameter then because of the *quantum vacuum* (or *quantum aether*) there could appear a radiative "thermal" noise directly related to the acceleration parameter(s) of the problem. As is well known, such effects have been considered for the first time by Unruh [4], who introduced the concept of *quantum detector* and derived the vacuum temperature,  $T_V = \frac{\hbar}{2\pi c k_B} a$ , for a linear accelerating one, and by Davies [5], who used a mirror model, in the mid-seventies. Hawking effect [6] was shown to be of the same type. They are generally considered amongst the most fundamental things Physics is telling us.

---

<sup>1</sup>e-mail: rosu@ifug.ugto.mx

Traditionally, there are two basic physical mechanisms related to the quantum radiative processes [7]. One is directly related to the *randomness* of the interaction of the particles with the vacuum fluctuations, the other is the so-called electromagnetic self interaction, better known as *radiation reaction*. Here, we shall consider only the *randomness* aspect of the vacuum noise. The general problem to study is the coordinate dependence of the spectrum of vacuum fluctuations as seen by quantum detectors/particles moving along classical trajectories. This noise spectrum is a density of states times the Fourier transform of the autocorrelation function (Wightman two-point function) of a quantum field along the classical trajectory. For a relativistic, linear accelerated detector the scalar vacuum noise has an *exact* thermal representation. This is a remarkable theoretical result because one could identify the temperature parameter directly as a scaled kinematical invariant (the proper acceleration); the rest of physics (quantum mechanics, relativity and statistical mechanics) enters as a scaling factor of the corresponding fundamental constants. One would like to prove this “beautiful truth” by experiment, and indeed a number of proposals to detect such quantum “thermal” effects have been made in the past (for a review, see [8]). One of them belongs to Rogers [9] and refers to the very successful Geonium Physics.

The idea of Rogers is to place a small superconducting PT containing a single electron in a microwave cavity and perform a resonant transfer of the cyclotron vacuum noise at the relativistic-shifted axial frequency to a cavity mode.

Rogers’ trap is somewhat unusual compared to the common geonium traps. The cap electrode separation is  $2z_0 = 2$  mm, the electrode potentials  $\pm U_0 = \pm 10$  kV, and the magnetic field is  $B = 150$  kG, whereas representative values of parameters for geonium traps are electrode separations of tens of mm, electrode potentials of several volts, and magnetic fields of 1-2 T (Dehmelt has used a trap with  $2z_0 = 8$  mm, quadrupole potential of several volts and magnetic field of 5 T). The 15 T of the Rogers trap are attainable with superconducting solenoid magnets. The microwave cavity is of length 1 cm, radius 1.36 cm, and  $Q \approx 10^4$ .

The single electron is constrained to move in a cyclotron orbit around the trap axis by the powerful axial magnetic field. Rogers considered the circular proper acceleration to be  $a = 6 \times 10^{19} g_\oplus$ , i.e., an electron having the velocity  $\beta = 0.6$ . This acceleration, if considered as linear, corresponds to a temperature of the vacuum  $T_V = 2.4K$ . The velocity of the electron is maintained constant by means of a circularly polarized wave with the frequency equal to the cyclotron one, compensating at the same time for the irradiated power (the synchrotron damping width is  $\Gamma_c = \frac{4e^2\omega^2}{3mc^2}$ ). The static quadrupole electric field of the trap creates a quadratic potential well along the trap axis in which the electron oscillates. The frequency of observation is  $\omega = \gamma\omega_z = 10.57$  GHz for the device scale and working conditions chosen by Rogers. At this frequency, the difference in energy densities between cyclotron (circular) noise and the universal Unruh noise are negligible. The spectral power density of the cyclotron noise at the relativistic-shifted axial frequency is  $\partial P/\partial f = 0.47 \cdot 10^{-22}$  W/Hz. This power is resonantly transferred to the  $TM_{010}$  mode of the microwave cavity and a most sensible cryogenic GaAs field-effect transistor amplifier should be used in order to have an acceptable signal-to-noise ratio  $S/N = 0.3$ . According to Rogers, the signal may be distinguished from the amplifier noise in about 12 msec.

The experiment of Rogers requires the top of electronics, cryogenic techniques, and geonium methods.

The great advantage of Rogers’ proposal over that of Bell and Leinaas [10] resides in using a single electron instead of electron bunches. In this way, the very complicated stochasticity of the beam dynamics at a storage ring is avoided and a better control of the dynamics is allowed just because of the small spatial

scale of the Penning device.

The PT problem can be defined as the interaction of the electron with an external field consisting of a uniform magnetic field  $\mathbf{B} = B\hat{\mathbf{e}}_z$  and a superposed electric quadrupole potential :

$$\Phi(x, y, z) = \frac{U_0}{2z_0^2 + r_0^2}(2z^2 - x^2 - y^2) . \quad (1)$$

Details of the practical achievement of this combination of fields are given in [1]. The electrostatic field is  $E = -\nabla\Phi$ .  $U_0$  is the potential difference between a one-sheeted hyperboloid which is the ring electrode and a two-sheeted hyperboloid forming the cap electrodes,  $r_0$  is the inner radius of the ring electrode and  $2z_0$  is the distance between the two end-cap electrodes. In the ideal case (i.e., perfect cylindrical geometry) there exist three bounded motions in the trap: (i) an axial harmonic motion in a parabolic well along the direction of the magnetic field, (ii) the cyclotron motion at a higher frequency, in the perpendicular (x,y) plane, and (iii) a magnetron motion in the same plane at a much lower frequency. The axial and cyclotron motions are usually excited with nearly resonant , radio-frequency drives applied to the trap electrodes. For a single electron in the trap (obtained with an “evaporation” technique due to Dehmelt and collaborators) the classical equations of motion are as follows

$$\ddot{z} + \omega_z^2 z = 0 , \quad (2a)$$

$$\ddot{r} = \frac{1}{2}\omega_z^2 r - i\omega_c \dot{r} , \quad (2b)$$

where  $r = x + iy$ ,  $\omega_z^2 = 4eU_0/m(r_0^2 + 2z_0^2)$ ,  $\omega_c = eB/mc$ . The solutions are

$$z = r_z \cos\omega_z t \quad (3a)$$

and

$$r = r_c e^{-i\omega'_c t} + r_m e^{-i\omega_m t} , \quad (3b)$$

where

$$\left\{ \begin{array}{c} \omega'_c \\ \omega_m \end{array} \right\} = \frac{1}{2}\omega_c \pm [(\omega_c/2)^2 - (\omega_z^2)/2]^{1/2} . \quad (4)$$

The amplitudes  $r_m$ ,  $r_c$ ,  $r_z$ , as well as some possible phases are determined by the initial conditions. The motion is stable only for  $\omega_c^2 > 2\omega_z^2$ , see the rhs of Eq. (4), which is the so-called trapping condition, implying periodic solutions. A typical xy orbit for the case  $\omega_m \ll \omega_c$ ,  $r_m < r_c$  is shown in Fig. 1 of Ref. [11]. Only under such conditions, the magnetron motion can be interpreted as an  $\vec{E} \times \vec{B}$  drift of the center of the cyclotron orbit around the trap axis.

The electron motion in the trap can be analyzed by many procedures in analytical mechanics [12]. However, we are interested in the transition rates of a quantum electron due to the vacuum fluctuations affecting its classical trajectory. Unfortunately, the classical trajectory in the trap is not a simple one. From the point of view of the circular (cyclotron) vacuum noise we are not in the ideal situation; the magnetron drift is also there, though usually considered as negligible, or diminished by pumping at  $\omega_z + \omega_m$  [13]. Besides, one has to take care of all the other, more common sources of noise (for a discussion of the axial Brownian noise see Ref. 1b). Even in the case of pure cyclotron motion, it is well known that the circular vacuum noise is not at all universal thermal ambience [14]. Rogers has used the formula for the spectral energy density of a massless scalar field as given by Kim, Soh and Yee (KSY) [14], i.e.,

$$\left[ \frac{de}{d\omega} \right]_{\omega=\gamma\omega_z} = \frac{\omega^3}{\pi^2} \left( \frac{1}{2} + \frac{1}{2\gamma^2 r} \sum_{n=0}^{\infty} v^{2n} f_n(r) \right) , \quad (5)$$

where  $r = \frac{\omega}{\gamma\omega_0}$ ,  $v$  is the electron velocity,  $\gamma$  is the Lorentz factor, and  $f_n$  is a sum over an index  $k$  of step functions of argument  $n - k - r$ , indicating the excitation of zero-point field modes at frequencies of the type  $p\omega_c \pm \omega_z$ , with  $p$  a positive integer; in other words, the vacuum noise at  $\gamma\omega_z$  is a sum of cyclotron axial-displaced harmonics.

The KSY parametrization of the circular spectrum gives a quasi-continuous spectrum which is similar to the black-body one only at low frequencies and for the first few terms in the sum (Rogers considered the first four terms). The real PT cyclotron-axial-magnetron motion produces a more complicated vacuum noise, that one may call the Penning vacuum noise (PVN). Moreover, Levin, Peleg, and Peres [15] have shown that further Casimir corrections might be important. As a matter of fact, cavity effects have been thoroughly investigated in the literature in view of the decade-old debate on their direct substantial shifting of the spin magnetic moment [16]. Also, according to Becker, Wodkiewicz and Zubairy [17], a tiny squeezing effect of the cavity modes is possible just because of the presence of the electron in the cavity.

I would like now to emphasize that recently constructed cylindrical PTs [20], for which the trap itself is a microwave cavity, should be considered as an important simplification of the experimental setup for detecting the Penning Unruh-like noise. In this case, small slits ( $\approx 0.015$  cm) at  $\pm z_0$  above and below the trap center, incorporating choke flanges, divide the oxygen-free high-conductivity copper cavity walls into the required Penning electrode configuration, i.e., the two end-cap electrodes at the level of the slits and the ring electrode (of radius  $r_0$ ). Furthermore, there are two essential compensation electrodes of height  $\Delta z_c/z_0 = 0.20$  placed near the end-cap ones, on which the applied potential is tuned in order to make the axial motion of the electron as harmonic as possible. It has been shown [20] that an “orthogonalized” geometry of the trap, i.e.,  $r_0 = z_0$ , reduces a lot the nonlinear axial frequency shifts. The three motions of the hyperbolic trap, the cyclotron, axial, and magnetron ones take place at  $\approx 166$  GHz, 63 MHz, and 12 kHz, respectively [20]. Moreover, the driven axial resonance for this configuration has been observed with almost the same signal-to-noise ratio as in hyperbolic PTs. By means of these cylindrical cavity-traps, a direct coupling to the cavity modes may be achieved, especially in the common weak coupling regime, where the cyclotron oscillator and the cavity mode cannot form normal modes, and thus supplementary nonlinear effects are not coming into play. The cylindrical Penning Unruh-like noise will be a tiny form of radiative cooling of the electron oscillator. The cylindrical  $TM_{010}$  mode is essentially a Bessel function of zero order in the radial direction without nodes along the  $z$ -axis. The frequency of  $TM_{010}$  is given by  $\omega_{010} = \frac{c\xi_{01}}{\rho_0}$ , where  $\xi_{01} = 2.405$  is the first zero of the Bessel function of zero order. The price to pay in the case of the cylindrical trap is a loss in the quality of the electrostatic quadrupole potential, despite the compensating electrodes helping partial control of this problem.

I now pass to some comments on other related problems in Penning trap physics. Some time ago Fernández and Nieto [18] used the Weyl-Wigner-Moyal phase space formalism to calculate the energy levels of a spinning charged particle in a PT. Here we wish to point out that the study of the PT phase space is a problem one should place in the more general context of Chaos Science [19]. In phase space the uncertainty principle replaces the continuum of classical states within a volume  $\hbar^N$  (where  $N$  is the number of degrees of freedom of the system) by a single quantum state, and as such, quantum chaos is less ‘powerful’ than the classical one. In a certain sense, to go beyond quantum mechanics, at least in phase space, one might think of more accurate topologies defining the proximity of points (states) [21]. In this way, one will have an intermediate

picture between quantum chaos and the standard geometric (classical) chaos. The geometric chaos is a mathematical one devoided of physical reality. For the physical world it is the ideal-limiting case never satisfied by real measurements. However, one would like to estimate as best as possible how far we are from the ideal case in an experiment. In this sense, the proximity topology might help progress in a better evaluation of experimental precision.

One should also keep in mind that in the case of a particle moving in a plane to which a magnetic field is applied in the normal direction, the momentum operators cease to commute:

$$[p_1, p_2] = ieB\hbar . \quad (6)$$

This introduces a cellular structure in the momentum plane, which becomes divided into Landau cells of area proportional to  $eB\hbar$ . Thus, in the PT case, the Landau cells are to be found in the momentum plane normal to the axial magnetic field.

We also recall that in the semiclassical limit the magnetic field determines not only the classical trajectory of the particle through the Lorentz force (dynamical effect), but also contributes to the phase accumulated along the trajectory through the line integral of the vector potential along it (geometric effect) [22].

In conclusion, as far as the electromagnetic trapping is considered as one of the most precise tools at our disposal, one might think that extremely tiny fundamental effects are best accessible by means of such experimental techniques. However, one should be fully aware of the host of more standard effects which are also involved in the trap laboratory. In my short inspection of Rogers' proposal I put forth a number of suggestions that presumably are worthy to be pursued in future more detailed studies. The use of cylindrical traps may contribute to turn not only Rogers' proposal into a more realistic one, but also other even more exotic proposals [23].

Finally, if one takes the viewpoint that the circular Unruh noise is actually the common electromagnetic radiation of an accelerated electron (this is the standpoint of the present author too) then the discussions of the vacuum noise are still of some worth, this time aimed merely at introducing a new temperature parameter in the PT physics.

## Acknowledgments

This work was partially supported by the CONACyT Project 4868-E9406.

## References

- [1] D.J. Wineland, W.M. Itano, and R.S. van Dyck, "High-resolution spectroscopy of stored ions", *Adv. At. Mol. Phys.* **19**, 135 (1983); L.S. Brown and G. Gabrielse, "Geonium theory: Physics of a single electron or ion in a Penning trap", *Rev. Mod. Phys.* **58**, 233 (1986); H. Dehmelt, "Experiments with an isolated subatomic particle at rest", (Nobel lecture, December 8, 1989), *Rev. Mod. Phys.* **62**, 525 (1990); W. Paul, "Electromagnetic traps for charged and neutral particles", (Nobel lecture, December 8, 1989), *ibid*, **62**, 531 (1990)
- [2] I. Marzoli and P. Tombesi, "Quantum non-demolition measurement of cyclotron excitations and wave function collapse", *Europhysics Lett.* **24**, 515 (1993); R.G. Hulet et al., "Precise test of quantum jump theory", *Phys. Rev. A* **37**, 4544 (1988)

- [3] S. Takagi, “Vacuum noise and stress induced by uniform acceleration”, *Prog. Theor. Phys. Suppl.* **88**, 1 (1986)
- [4] W.G. Unruh, “Notes on black hole evaporation”, *Phys. Rev. D* **14**, 870 (1976)
- [5] P.C.W. Davies, “Scalar particle production in Schwarzschild and Rindler metrics”, *J. Phys. A: Math. and Gen.* **8**, 609 (1975)
- [6] S.W. Hawking, “Black hole explosions?”, *Nature* **248**, 30 (1974)
- [7] J. Dalibard, J. Dupont-Roc and C. Cohen-Tannoudji, “Vacuum fluctuations and radiation reaction: identification of their respective contribution”, *J. Physique* **43**, 1617 (1982)
- [8] H.C. Rosu, “Hawking-like and Unruh-like effects: Towards experiments?”, *gr-qc/9406012* (1994)
- [9] J. Rogers, “Detector for the temperaturelike effects of acceleration”, *Phys. Rev. Lett.* **61**, 2113 (1988)
- [10] J.S. Bell and J.M. Leinaas, “Electrons as accelerated thermometers”, *Nucl. Phys. B* **212**, 131 (1983); For comments on this proposal see H. Rosu, “On the circular vacuum noise in storage rings”, *Nuovo Cimento B* **109**, 423 (1994)
- [11] W.M. Itano and D.J. Wineland, “Laser cooling of ions stored in harmonic and Penning traps”, *Phys. Rev. A* **25**, 35 (1982)
- [12] M. Kretzschmar, “A theory of anharmonic perturbations in a Penning trap”, *Z. Naturforsch.* **45 a**, 965 (1990); “Particle motion in a Penning trap”, *Eur. J. Phys.* **12**, 240 (1991); “Single particle motion in a Penning trap: Description in the classical canonical formalism”, *Physica Scripta* **46**, 544 (1992); “Excitation of particle motions in a Penning trap: Description in the classical canonical formalism”, *Physica Scripta* **46**, 555 (1992)
- [13] D. Wineland, P. Ekstrom, and H. Dehmelt, “Monoelectron oscillator”, *Phys. Rev. Lett.* **31**, 1279 (1973)
- [14] S.K. Kim, K.S. Soh, and J.H. Yee, “Zero-point field in a circular-motion frame”, *Phys. Rev. D* **35**, 557 (1987). These authors actually confirmed the results of Letaw and Pfautsch obtained during 1980-1982. See J.R. Letaw and J.D. Pfautsch, “Quantized scalar field in rotating coordinates”, *Phys. Rev. D* **22**, 1345 (1980); “Quantized scalar field in the stationary coordinate systems of flat spacetime”, *Phys. Rev. D* **24**, 1491 (1981); “The stationary coordinate systems in flat spacetime”, *J. Math. Phys.* **23**, 425 (1982). See also, G. Denardo and R. Percacci, “Quantum field theory for a rotating observer”, *Nuovo Cimento* **48 B**, 81 (1978)
- [15] O. Levin, Y. Peleg and A. Peres, “Unruh effect for circular motion in a cavity”, *J. Phys. A: Math. Gen.* **26**, 3001 (1993)
- [16] L.S. Brown et al., “Cyclotron motion in a microwave cavity: Lifetime and frequency shifts”, *Phys. Rev. A* **32**, 3204 (1985) and references therein.
- [17] W. Becker, K. Wodkiewicz, and M.S. Zubairy, “Squeezing of the cavity vacuum by charged particles”, *Phys. Rev. A* **36**, 2167 (1987)
- [18] D.J. Fernández C. and L.M. Nieto, “Penning trap from the phase space quantum mechanics”, *Phys. Lett. A* **157**, 315 (1991)

- [19] N. Srivastava, C. Kaufman, and G. Müller, “Hamiltonian chaos III”, *Computers in Physics* **6/1**, 84 (1992)
- [20] J. Tan and G. Gabrielse, “One electron in an orthogonalized cylindrical Penning trap”, *Appl. Phys. Lett.* **55**, 2144 (1989); “Parametrically pumped electron oscillators”, *Phys. Rev. A* **48**, 3105 (1993)
- [21] Yu.M. Smirnov, “On proximity spaces”, *Mat. Sbornik, N.S.*, **31/73**, 543 (1952); S. Leader 1959, “On clusters in proximity spaces”, *Fund. Math.* **47**, 205 (1959); “On completion of proximity spaces by local clusters”, *Fund. Math.* **48**, 201 (1960)
- [22] D.J. Fernández and N. Bretón, “Is there a prescribed parameter’s space for the adiabatic geometric phase”, *Europhys. Lett.* **21**, 147 (1993); D.J. Fernández C., “Transformations of a wave packet in a Penning trap”, *Nuovo Cimento B* **107**, 885 (1992)
- [23] M. Karim, “Observable effect of curvature on super-radiant states”, *Int. J. Mod. Phys. A* **5**, 143 (1990); “Free fall of the vacuum”, *Phys. Lett. A* **171**, 7 (1992)