

# The Accelerated Observer and Quantum Effects

Roberto Casadio<sup>y</sup> and Giovanni Venturi<sup>z</sup>

Department of Physics, University of Bologna, and  
Istituto Nazionale di Fisica Nucleare, Sezione di Bologna, Italy

## Abstract

An extended monopole detector at constant acceleration coupled to a massless scalar field is allowed to evolve quantum mechanically. It is found that while in the classical, followed by the point particle, limit the usual result (Unruh effect) is recovered, in the point particle (before the classical) limit the detector decouples from the scalar field and therefore the effect disappears.

## List of keywords:

Accelerated observer  
Classical limit  
Extended monopole detector  
Point particle limit  
Quantum mechanics  
Unruh effect

---

<sup>y</sup> e-mail: Casadio@bologna.infn.it

<sup>z</sup> e-mail: Amitage@bologna.infn.it

## I. Introduction

Hawking's remarkable discovery [1] that black holes evaporate due to quantum particle emission and behave as if they had an effective temperature of  $(8M)^{-1}$  with  $M$  the mass of the black hole also encouraged, because of the equivalence principle, the study of field theory in accelerated systems [2,3,4]. In particular it was found that a detector with uniform acceleration in the usual vacuum state of Minkowski space will be thermally excited to a temperature  $T = a/2\pi$ .

In the derivation of the result [5] a massless, otherwise free, scalar field  $(\phi)$  is linearly coupled to a particle detector. This then serves as a localized probe of the field which is assumed to be in the Minkowski vacuum. The detector is described by the DeWitt monopole moment [6] and evolves on a trajectory of constant acceleration. One then uses time dependent perturbation theory and examines the two point function of the scalar field in terms of the detector's proper time. A question which one can naturally ask is what happens if one considers detectors of finite extent which evolve quantum mechanically [7]. Such a model is considered in the next Section. We shall use units for which  $c$  and the Boltzmann constant are set equal to unity.

## II. An Extended Quantum Mechanical Detector

The Lagrangian density describing the field  $\phi$  and its interaction with a pointlike detector in four dimensional Minkowski space will be given by:

$$L' = \int d^4x \left[ -\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - Q(x) \phi(x) \right] \quad (2.1)$$

where  $x_c(\tau)$  denotes the worldline of the idealized classical pointlike detector,  $Q(\tau)$  represents its monopole moment at time  $\tau$  and  $\int d^4x$  serves to transform inertial time integrals into proper time integrals.

Concerning  $x_c(\tau)$  it is convenient to introduce Rindler coordinates  $(\xi, \eta)$  and  $(\tau, z)$  (associated with the transformation to the accelerated frame) related to the Minkowski coordinates  $x^0$  and  $x^3$  through:

$$\begin{aligned} x^0 &= a^{-1} e^a \sinh a \tau \\ x^3 &= a^{-1} e^a \cosh a \tau \end{aligned} \quad (2.2)$$

where the acceleration is in  $x^3$  direction and has a local value equal to  $a$  on the hypersurface  $\eta = 0$  and  $a e^{-a \xi}$  elsewhere. In what follows we set  $\eta = 0$  without loss of generality, further we denote by  $x_c^0$  and  $x_c^3$  the corresponding Minkowski coordinates associated with this  $\eta = 0$  trajectory. The uniformly accelerating classical observer will then have a path given by  $x_c^0, x_c^3(\tau)$ , with proper time  $\tau$  and  $x_c^1(\tau) = x_c^2(\tau) = 0$ .

In Eq. (2.1) we have the Lagrangian for the scalar field  $\phi$  and its coupling to the detector, to this we should add a Lagrangian for the observer which leads to the desired solution for the classical path  $x_c$ . A convenient Lagrangian yielding an action whose stationarity leads to the classical path  $x_c$  is [8] (it is sufficient to just exhibit  $x^3 = z$  since the acceleration is in that direction, and the corresponding Lagrangians  $L_x, L_y$  along other directions are obtained by setting  $a = 0$  and replacing  $z$  by  $x, y$  respectively):

$$L_z = \frac{m}{2} \left( \frac{dz}{d\tau} \right)^2 + a^2 z^2 = \frac{m}{2} \dot{z}^2 + a^2 z^2 \quad ; \quad (2.3)$$

where  $m$  is the mass of the observer which is subject to a field of force  $m a^2 z$  corresponding to an inverted harmonic oscillator potential and the total Lagrangian will of course be the sum of Eq. (2.3) with  $L_x$  and  $L_y$ . Further the Hamiltonian associated with  $L_z$  is given by:

$$H_z = \frac{m}{2} \dot{z}^2 - a^2 z^2 \quad ; \quad (2.4)$$

which is a constant along the path. At this point we shall further pursue the analogy with the harmonic oscillator and if we write the total wave function for the three-dimensional monopole detector in a factorized form  $\psi(x; y; z)$  where for each we shall later consider a gaussian wavepacket, we then obtain a quantum Lagrangian density:

$$L_z = \frac{1}{2} \dot{x}^2 + \frac{1}{2} \dot{y}^2 - i\hbar \psi^* \left( z; \right) \left( -z; \right) \frac{\hbar^2}{2m} \frac{\partial^2 \psi(z; )}{\partial z^2} + \frac{m}{2} a^2 z^2 \psi(z; )^2 ; \quad (2.5)$$

which in the WKB approximation leads to the Lagrangian Eq. (2.3) [8]. We note the dynamics is associated with the z coordinate (and not  $x^1 = x, x^2 = y$ ) thus again we have just exhibited the pertinent term in the total Lagrangian density. The corresponding terms for  $\psi(x; y)$  are obtained from the above by setting  $a = 0$  and interchanging z and x (y).

The inverted harmonic oscillator has been studied [9] and in particular the associated Green's function  $K(z; z^0; 0)$  can be obtained from the corresponding harmonic oscillator one through the replacement of the frequency of oscillation by  $ia$ . One then has:

$$K(z; z^0; 0) = \frac{m a}{2 i\hbar \sinh a} \exp \frac{im a}{2 \hbar \sinh a} (z^2 + z^2 \cosh a - 2 z z^0) ; \quad (2.6)$$

in which case if one begins with a gaussian wavepacket  $\psi(z^0; 0)$ :

$$\psi(z^0; 0) = \frac{1}{b} \exp \frac{(z^0 - z_0)^2}{2b^2} ; \quad (2.7)$$

one will obtain:

$$\psi(z; ) = \frac{1}{b} \frac{m a}{2 i\hbar \sinh a} \frac{1}{\frac{1}{2b^2} - i \cosh a} \exp \left[ i \left( z^2 \cosh a + \frac{1}{4b^4} \left( \frac{1}{4b^4} + \cosh^2 a \right) z_0^2 \cosh a - 2 z z_0 \right) \right] \exp \left[ \frac{1}{2b^2} \left( \frac{1}{4b^4} + \cosh^2 a \right) z z_0 \cosh a \right] ; \quad (2.8)$$

where  $m a = (2h \sinh a)$  and:

$$j(z; )j^2 = \frac{1}{b^{1=2}} \frac{m a}{2 h \sinh a} \frac{1}{\frac{1}{4b^4} + \cosh^2 a} \frac{1}{1=2} \exp \left[ \frac{2}{b^2} \frac{1}{4b^4} + \cosh^2 a \right] z - z_0 \cosh a \quad (2:9)$$

where  $= b \frac{1}{4b^4} + \cosh^2 a$  and  $z_c = z_0 \cosh a$ . It is clear that in the limit for  $\rightarrow 1$  the R.H.S. of Eq. (2.9) will tend to  $(z - z_c)$  and this suggests we replace  $L$  by:

$$L' = \int_0^Z dx^0 - x_c^0 ( ) Q ( ) j(x; )j^2 j(y; )j^2 j(z; )j^2 (x) \quad (2:10)$$

where  $j(x; )j^2 (j(y; )j^2)$  is given by Eq. (2.8) with  $z_0 = a = 0$  and  $z$  replaced by  $x (y)$  and leads to  $(x) (y)$  for  $(a = 0) \rightarrow 1$ . Thus in the limit for  $\rightarrow 1$ ,  $(a = 0) \rightarrow 1$  Eq. (2.10) leads to Eq. (2.1).

As usual [5,6] we shall suppose the detector has a discrete set of internal eigenstates described by vectors  $jE_i$  where  $E = 0$  corresponds to the ground state and evaluate the probability  $P(E)$  for the detector make a transition from the vacuum to an excited state of energy  $E (h!)$  while the scalar field undergoes a transition from the Minkowski vacuum to any final state. One then has:

$$P(E) = hE jQ(0)j_0 i^2 \int_0^Z dZ_L \int_0^Z dZ_L e^{-i(E-h!)Z} G(t; 0) = hE jQ(0)j_0 i^2 \int_0^Z dT \int_{L+2j}^Z dT e^{-i!t} G(t; T); \quad (2:11)$$

where, for convenience, we have considered a finite time interval  $(0;L)$  for the detector [10], defined  $t = T + t=2$ ,  $t^0 = T - t=2$  and:

$$\begin{aligned}
 G(t;T) &= \frac{1}{3} \int_0^L dx dx^0 dy dy^0 dz dz^0 h_0 j'(\mathbf{x};\mathbf{y};\mathbf{z};\mathbf{x}_c^0)'(\mathbf{x}^0;\mathbf{y}^0;\mathbf{z}^0;\mathbf{x}_c^0) j_0 i \\
 &= \frac{1}{4^5} \int_0^L dx dx^0 dy dy^0 dz dz^0 \exp\left[-\frac{a^2}{4} \left( \frac{z - z_c}{2} \right)^2 - \frac{a^2}{4} \left( \frac{z + z_c}{2} \right)^2 - \frac{a^2}{4} (x^2 + y^2) - \frac{a^2}{4} (x^0^2 + y^0^2) \right] \\
 &\quad \frac{\exp\left[-\frac{a^2}{4} \left( \frac{z - z_c}{2} \right)^2 - \frac{a^2}{4} \left( \frac{z + z_c}{2} \right)^2 - \frac{a^2}{4} (x_c^0 - x_c^0)^2 - \frac{a^2}{4} (i)^2 \right]}{(x - x^0)^2 + (y - y^0)^2 + (z - z^0)^2 + (x_c^0 - x_c^0)^2 + (i)^2} \quad (2:12)
 \end{aligned}$$

where the unprimed (primed) quantities are evaluated at  $(t)$  and  $(t^0)$  ( $a = 0$ ). If we now define  $u = z - z_c - z^0 + z_c^0$ ,  $v = (z - z_c + z^0 - z_c^0)/2$ ,  $p = x - x^0$ ,  $q = (x + x^0)/2$ ,  $r = y - y^0$  and  $s = (y + y^0)/2$  one obtains:

$$\begin{aligned}
 G(t;T) &= \frac{1}{4^5} \int_0^L du dv dp dr dq ds \frac{\exp\left[-\frac{a^2}{4} (v - u/2)^2 - \frac{a^2}{4} (v + u/2)^2 - \frac{a^2}{4} (q - p/2)^2 + \frac{a^2}{4} (s - r/2)^2 - \frac{a^2}{4} (q + p/2)^2 + \frac{a^2}{4} (s + r/2)^2 \right]}{D \left( \frac{u^2}{2} + \frac{p^2}{2} + \frac{r^2}{2} \right)} \\
 &= \frac{1}{4^5} \int_0^L \frac{\exp\left[-\frac{a^2}{2} \left( \frac{u^2}{2} + \frac{p^2 + r^2}{2} \right) \right]}{D \left( \frac{u^2}{2} + \frac{p^2}{2} + \frac{r^2}{2} \right)} du dp dr \quad ; \quad (2:13)
 \end{aligned}$$

where  $D = \frac{4}{a^2} \sinh^2 \frac{a}{2} (t - t^0) i$  and  $z_c = z_c^0$ .

We may now consider the integral over the time variable  $t$  of  $G$  :

$$\begin{aligned}
 \int_{L-2j}^{L+2j} dt e^{-i!t} G(t;T) &= \\
 &= \int_{L-2j}^{L+2j} du dp dr \frac{a^2}{4} \int_0^L dt \frac{\exp\left[-\frac{a^2}{2} \left( \frac{u^2}{2} + \frac{p^2 + r^2}{2} \right) \right]}{\sinh \frac{a t}{2} \left( \frac{u^2}{2} + \frac{p^2}{2} + \frac{r^2}{2} \right)} e^{-i!t} \quad ; \quad (2:14)
 \end{aligned}$$

where we have inverted orders of integration and:

$$u = \frac{a}{2} u \sinh aT \sqrt{u^2 \sinh^2 aT + u^2 + p^2 + r^2}^{1=2}$$

$$= u \frac{a}{2} \sinh aT \sqrt{\cosh^2 aT + \frac{p^2 + r^2}{u^2}}^{1=2}; \quad (2:15)$$

which implies the existence of poles at (we shall close the contour in the lower half plane):

$$t = \frac{2 \ln}{a} + t^n; \quad (2:16)$$

where  $n > 0$  and  $t^n$  is purely real and is given by:

$$t^n = (-1)^n \frac{2}{a} \sinh^{-1} u; \quad (2:17)$$

On performing the integral with respect to  $t$  in Eq. (2.14) using the theory of residues one has:

$$\int_{L+2j\pi}^{L+2j\pi} dt e^{it} G(t;T) = \quad (2:18)$$

$$= \frac{a}{2} \int_1^{z^1} du dp dr \sum_{n=1}^N \frac{(-1)^n e^{\frac{2n}{a}}}{2^4 u_+} \frac{e^{-it} t^n}{(1+u_+^2)^{1=2}} \frac{e^{-it} t^n}{(1+u_+^2)^{1=2}};$$

where we have omitted an integral along the contour which does not contribute for  $L$  large and  $N$  is the largest integer  $(L+2j\pi, L=2j) a=2$ . Further:

$$= \int_0^1 \frac{1}{2 + u^2} \frac{1}{2} \exp \left[ \frac{2 \ln}{2 + u^2} u^2 - \frac{2}{2} (p^2 + r^2) \right]; \quad (2:19)$$

with:

$$= \frac{1}{4 + b^2} + b^2 (-1)^n (1+u^2)^{1=2} \cosh aT + u \sinh aT^2 \quad (2:20)$$

$$= \frac{1}{4 + b^2} + b^2 (-1)^n (1+u^2)^{1=2} \cosh aT - u \sinh aT^2 \quad (2:21)$$

$$= \frac{m a}{2h} (-1)^n (1+u^2)^{1=2} \sinh aT + u \cosh aT \quad (2:22)$$

$$= \frac{m a}{2h} (-1)^n (1+u^2)^{1=2} \sinh aT - u \cosh aT \quad (2:23)$$

$$= 0 = (a = 0); \quad (2:24)$$

Finally, for the transition probability Eq. (2.11) one will obtain:

$$P(E) = hE \int_0^L \int_0^1 dT \int_0^1 du dp dr \frac{1}{4} \sum_{n=1}^{\infty} \frac{e^{-\frac{2n}{a}}}{2u \cosh^2 aT + \frac{p^2+r^2}{u^2}} \frac{\sin^n(\frac{1}{u}t_+)}{(1+u^2)^{1/2}} \frac{\sin^n(\frac{1}{u}t_+)}{(1+u^2)^{1/2}} ; \quad (2.25)$$

and we note the appearance of the familiar Planck distribution factor. Concerning the above expansion we note that the first integral may lead to a divergence as  $L \rightarrow \infty$  which as usual [5] can be eliminated by considering the transition probability per unit time, further we observe that the  $u, p, r$  integrations are finite and we may now consider the limits for  $h \rightarrow 0$  and  $b \rightarrow 0$ .

Let us first consider  $h \rightarrow 0$ , in such a limit one has that  $\lambda, \lambda_0$  tend to infinity and the Compton wavelength of the detector will disappear from Eqs. (2.20), (2.21) and (2.25), which will however remain a complicated expression which nonetheless still exhibits a Planck distribution. The expression however simplifies if we then consider the  $b \rightarrow 0$  limit (point detector), indeed in such a case:

$$\lim_{b \rightarrow 0} \lim_{h \rightarrow 0} P(E) = \lim_{b \rightarrow 0} \int_0^1 \frac{1}{2b \cosh aT} \exp \left[ -\frac{u^2}{2b^2 \cosh^2 aT} - \frac{p^2+r^2}{2b^2} \right] \exp \left[ -\frac{p^2+r^2}{2b^2} \right] du dp dr ; \quad (2.26)$$

where we have omitted terms of higher order in  $(u;p;r)$ . The transition probability  $P(E)$  will then become:

$$\lim_{b \rightarrow 0} \lim_{h \rightarrow 0} P(E) = hE \int_0^L \int_0^1 dT \frac{1}{2} \sum_{n=1}^{\infty} e^{-\frac{2n}{a}} ; \quad (2.27)$$

in agreement with previous results [5].

One may now consider the limit  $b \rightarrow 0$ , one then has:

$$\lim_{b \rightarrow 0} P(E) = \lim_{b \rightarrow 0} 2b \int_0^1 \frac{1}{2 + \frac{p^2+r^2}{b^2}} \exp \left[ -\frac{4b^2}{2 + \frac{p^2+r^2}{b^2}} u^2 \right] du dp dr e^{-\frac{p^2+r^2}{2b^2}} = 0 ; \quad (2.28)$$

where  $\omega_0 = m = \hbar (u^2 + p^2 + r^2)^{1/2}$ . Since the  $u, p, r$  integrals in Eq. (2.25) are finite and furthermore in this case so is the  $T$  integral even in the limit for  $L \rightarrow 1$ , one has:

$$\lim_{\hbar \rightarrow 0} P(E) = 0; \quad (2.29)$$

thus implying that a quantum mechanically evolving point detector at constant acceleration decouples from the scalar field, which is rather surprising.

### III. Conclusion

We have considered a massless scalar field coupled to a monopole detector which from a given time is subjected to a constant acceleration. For the detector we considered a gaussian wavepacket which we allowed to evolve according to an inverted harmonic oscillator potential corresponding to a constant acceleration. On examining the probability per unit time for the detector to excite itself due to the absorption of scalar quanta, we observed that on first considering the classical limit ( $\hbar \rightarrow 0$ ) and then the pointlike limit for the detector we reproduced the usual results. If however one considers the pointlike limit first the detector decouples from the scalar field. This rather surprising result is due to the fact that once quantum mechanical evolution is considered another length enters the theory: the Compton wavelength of the detector (let us remember that we have taken  $b, m, \hbar$  and  $a$  as independent).

Indeed in another, quite different, situation a similar result is obtained [11]. One considers, rather than the quantum electrodynamics of point particles, the point particle limit of a theory of extended particles and takes this limit on the quantum level rather than the classical level. Then, in the nonrelativistic approximation, such a method leads to a vanishing self-energy and to the absence of runaway and preacceleration effects.

One may also ask oneself whether an approach such as the above is possible for a black hole. Since black hole evaporation essentially takes place in  $s$ -waves

one can make a 2-dimensional model for a black hole and near the event horizon the Schwarzschild metric can be put in Rindler form [2,6]. Thus one may hope that since the acceleration becomes infinite near the event horizon it could compensate the zero due to the point particle limit. However for the scalar field Wightman function in two dimension one has a logarithmic singularity rather than a pole; usually through integration by parts one changes it to a pole, but for our wavepacket case such an approach is considerably more involved. We hope to return to this point.

#### References

- [1] S.W. Hawking, *Nature* 248 (1974) 30;  
S.W. Hawking, *Comm. Math. Phys.* 43 (1975) 199.
- [2] W.G. Unruh, *Phys. Rev. D* 14 (1976) 870.
- [3] P.C.W. Davies, *J. Phys. A* 6 (1975) 609.
- [4] D.W. Sciama, P. Chandela and D. Deutsch, *Adv. in Phys.* 30 (1981) 327.
- [5] See for example N.D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, England, 1982).
- [6] B.S. Dewitt in *General Relativity: an Einstein Centenary Survey*, eds. by S.W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1979).
- [7] For another discussion of particle detectors see P.G. Grove and Q.C. Ottewill, *J. Phys. A* 16 (1983) 3905.
- [8] R. Brout and Ph. Spindel, *Nucl. Phys. B* 348 (1991) 405;  
R. Brout, *Z. Phys. B* 68 (1987) 339.
- [9] G. Barton, *Annals of Physics* 166 (1986) 322.
- [10] In this context see *Response of finite time particle detectors in non inertial frames and curved backgrounds* by L. Sriram Kumar and T. Padmanabhan.
- [11] See H. Grotch, E. Kazes, F. Rohrlich and D.H. Sharp, *Acta Phys. Austr.* 54 (1982) 31 and references therein.