

Chaos in the Quantum Measurement Record

M. A. Nielsen^y

Department of Physics, University of Queensland 4072, Australia^z
(March 24, 2022)

We investigate measures of chaos in the measurement record of a quantum system which is being observed. Such measures are attractive because they can be directly connected to experiment. Two measures of chaos in the measurement record are defined and investigated numerically for the case of a quantum kicked top. A smooth transition between chaotic and regular behavior is found.

PACS Nos. 03.65.-w, 03.65.Bz, 05.45.+b

Understanding how chaotic behavior arises in quantum systems has been the subject of much research in recent years [1,2]. The purpose of this Letter is to investigate measures of chaos which are determined wholly by the measurement record of a quantum system which is undergoing observation. Such measures are attractive because they can be directly connected to experiment. It is well known that the dynamics of a quantum system depend on how that system is observed. This is in contrast to classical systems, for which it is usually assumed that measurements can be performed without disturbing the system. Thus we are led to a view of chaos in which the initial state, the free (Hamiltonian) dynamics, and the observation scheme are all regarded as parameters which can be varied in order to gain a better understanding of chaos in specific systems.

The Letter is organized as follows. We begin by reviewing some ideas from algorithmic information theory, which are then used to define a measure of chaos in the measurement record. A second measure of chaos in the measurement record is defined using the Shannon entropy. An inequality between these two measures is proved. A numerical investigation of these measures is then performed for the specific example of the quantum kicked top. We conclude with some general observations and comments on how the present approach to quantum chaos may be extended.

Suppose a series of measurements is made on a system, either classical or quantum. The result is a sequence of results, $R_1; R_2; \dots$, which we call the measurement

record. An intuitively appealing measure of chaos is the rate of growth of the information needed to describe the measurement record. If this occurs very quickly, then the measurement record is difficult to describe, and we would say that the system is chaotic. If the rate of information growth is slow, then the measurement record is easy to describe, and we would say that the system is regular.

To make this idea precise, we need to know how to quantify the "information needed to describe the measurement record". The solution to this problem is known as algorithmic information theory, which was created independently by Solomonov, Kolmogorov and Chaitin [3-5]. They defined the algorithmic information content $I(s)$ of a string s to be the length of the shortest possible description of that string. More rigorously, $I(s)$ is the length of the shortest algorithm which generates the string s . An algorithm, as defined by Turing [6], can be represented as a program for a universal computer, U : A data string s is used as input to such a program p , which then performs the algorithm on the data string. If the program ever halts then it prints an output string, $U(p;s)$.

For a particular universal computer U we define the algorithmic information content $I_U(s)$ of a string s to be the minimum length $|p|$ taken over all programs p for U such that $U(p;) = s$, that is the program p prints out s when nothing is input. The conditional algorithmic information content $I_U(s;t)$ of s given another string t is defined to be the minimum length $|p|$ taken over all programs p such that $U(p;t) = s$. Note that here and elsewhere it is assumed for convenience that p and s are represented as binary strings. In chapter six of [7] it is proved that

$$I_U(s;t) \leq I_U(s) + o(1); \quad (1)$$

where $o(1)$ is an order one constant which does not depend on s or t . This intuitively reasonable inequality is important in our later reasoning.

We now turn briefly to classical systems, using algorithmic information theory to define a measure of chaos, and seeing how this measure relates to other measures of chaos for classical systems. Suppose a classical system is initially at a point x_0 in some phase space. At times $t_1; t_2; \dots$ a measurement is carried out on the system. We suppose that this measurement can be modelled as follows: There is a partition of cells covering phase space, labelled $1; \dots; n$ and the measurement at time t_i determines which of these cells, R_i , the system is in at that

Submitted to Physical Review Letters.

^yElectronic address: m.nielsen@tangelo.phys.unm.edu

^zNew address: Center for Advanced Studies, Department of Physics and Astronomy, University of New Mexico, Albuquerque NM 87131-1156

time. Define the rate of information production of the classical system to be

$$R(x_0) = \limsup_{n \rightarrow \infty} \frac{I(R_1; \dots; R_n)}{n} \quad (2)$$

A minor technical point is that this definition is independent of which universal computer U is used to define algorithmic information, and thus no subscript U is needed on I or R . This follows from the fact [7] that for any two universal computers U_1 and U_2 , there is a constant C such that for any string s ,

$$|I_{U_1}(s) - I_{U_2}(s)| \leq C \quad (3)$$

It can be proved (see [8] for a review and references) that

$$R(x_0) \leq h; \quad (4)$$

where h is the topological entropy of the system. Positivity of the topological entropy is widely regarded as one of the best indicators of chaos in a classical system, and thus we can see that if $R(x_0)$ is ever positive then the topological entropy is positive and thus the system is chaotic.

Consider now the case of a quantum mechanical system which is observed at times $t_1; t_2; \dots$, producing a measurement record $R_1; R_2; \dots$. Again, the rate of information production may be defined as

$$R = \limsup_{n \rightarrow \infty} \frac{I(R_1; \dots; R_n)}{n} \quad (5)$$

In general the sequence $R_1; R_2; \dots$ is a random sequence because of the inherently probabilistic nature of quantum measurements. In order to estimate the rate of information production we use the result (see [9] for recent results and references) that the average algorithmic information of a random sequence $R_1; \dots; R_n$ given background information $B = \{p_{r_1; \dots; r_n}\}$ consisting of the probabilities of each possible measurement record $r_1; \dots; r_n$ satisfies the inequality

$$H_n - \bar{I}(R_1; \dots; R_n | B) \leq H_n + o(1); \quad (6)$$

where H_n is a form of the Shannon entropy H , which is defined by

$$H_n = \sum_X H(R_1; \dots; R_n) \\ = - \sum P(r_1; \dots; r_n) \log_2 P(r_1; \dots; r_n);$$

and the sum is over all the possible values $r_1; \dots; r_n$ that $R_1; \dots; R_n$ can take, and by convention $0 \log_2 0 = \lim_{x \rightarrow 0} x \log_2 x = 0$.

Applying the inequality (1) and the approximation result (6) we find that the average rate of information production satisfies the inequality

$$\overline{R} = \limsup_{n \rightarrow \infty} \frac{H(R_1; \dots; R_n)}{n} =: R_S; \quad (7)$$

where R_S is the asymptotic rate at which Shannon information is produced. R_S is another measure of chaos for the system, and can be interpreted in the following way. Suppose an experimentalist repeats the experiment a large number of times, obtaining on each run a measurement record $R_1; \dots; R_n$. Then the Shannon noiseless coding theorem [10] tells us that $H(R_1; \dots; R_n)$ is the smallest average codeword length that can be used to encode the measurement record. Thus R_S represents the asymptotic rate of growth of the average codeword length needed to encode the measurement record. See [9] for a precise account of the distinction between Shannon entropy and algorithmic information.

In practice, R_S was found to be considerably easier to compute than R . Furthermore, the inequality (7) allows us to obtain bounds on R by computing R_S . We will now numerically estimate R_S for the quantum kicked top.

The kicked top (see [11] and references therein) is a simple system whose classical analogue is known to exhibit chaos. In units where $\hbar = 1$, the evolution of the quantum kicked top from kick to kick is given by the unitary operator

$$U = \exp \left(i \frac{3}{2} J_z^2 \right) \exp \left(i \frac{1}{2} J_y \right); \quad (8)$$

where j is the angular momentum quantum number for the system. In this Letter we have used $j = 18$.

We suppose a projective measurement is performed on the system immediately after each kick. The projectors used are

$$P_+ = \sum_{m=0}^X |j; m\rangle \langle j; m| \quad (9)$$

$$P_- = \sum_{m<0} |j; m\rangle \langle j; m| \quad (10)$$

where $J_z |j; m\rangle = m |j; m\rangle$. This measurement determines whether the J_z component of angular momentum is non-negative or negative. If the state just before the measurement is $|j; i\rangle$, then the respective probabilities for the outcomes are given by the projection postulate as $\langle P_+ | j; i \rangle$ and $\langle P_- | j; i \rangle$.

In the numerical examples studied in this Letter the initial states of the system were chosen to be spin coherent states, which are defined by [11]

$$|j; i\rangle; \quad i = \exp(i(\phi \cos \theta - J_y \sin \theta)) |j; j\rangle; \quad (11)$$

In particular, we consider the state

$$|j; i\rangle = |j; j\rangle = 18; \quad \theta = 2.25; \quad \phi = 0.63i; \quad (12)$$

for which the means $\langle J_x \rangle; \langle J_y \rangle$ and $\langle J_z \rangle$ are located in the region of the classical configuration space for which

the classical kicked top is found to be regular, and the state

$$|j\rangle = |18\rangle; \quad |i\rangle = |1.64\rangle; \quad |i\rangle = |1.50\rangle; \quad (13)$$

for which the means $\langle h_{J_x} \rangle; \langle h_{J_y} \rangle$ and $\langle h_{J_z} \rangle$ are located in the region of the classical configuration space for which the classical kicked top is found to be chaotic.

In order to numerically estimate the rate of growth R_S of the Shannon entropy for an initial state $|j\rangle$ the following procedure was used. Suppose N kicks are performed on the system, and N measurements, one immediately after each kick. This will result in a sequence of measurement results $Z_1; \dots; Z_N$. The probability of this measurement history is given by the projection postulate,

$$P(Z_1; \dots; Z_N) = \langle j | U^{\dagger} P_{Z_1} U P_{Z_2} U^{\dagger} \dots U^{\dagger} P_{Z_N} U P_{Z_{N-1}} U \dots P_{Z_1} U | j \rangle; \quad (14)$$

Using these probabilities the Shannon information of the measurement record,

$$H_N = -H(Z_1; \dots; Z_N); \quad (15)$$

was computed for the range $1 \leq N \leq N$.

The results of this procedure when $N = 15$ are shown in figure 1. The lower line plots H_N for an initial state $|j\rangle$, and the upper line plots H_N for an initial state $|i\rangle$. There are two important things to note about this graph. First, for both initial states the growth in I_N settles down to be roughly linear in n very quickly. This has also been found to be the case with other initial states. Second, the system started in the state $|j\rangle$ corresponding to the classically regular region show a much lower rate of information production than the state $|i\rangle$, which corresponds to the classically chaotic region.

The nature of chaos in this system is strikingly illustrated by choosing 500 points at random from the $X > 0; Y > 0; Z < 0$ octant of the unit sphere, and calculating

$$R_S = \frac{H_{15}}{15}; \quad (16)$$

which, assuming H_N settles down quickly to a linear rate of growth, as found above, can be regarded as an approximation to R_S . This quantity was computed for each of the initial spin coherent states whose means correspond to the random points on the sphere. Figure 2 plots R_S as a function of the angle on the unit sphere between the point on the unit sphere corresponding to the initial coherent state, and the point at $\theta = 2.25; \phi = 0.63$, which is an elliptic fixed point for the classical map, and located deep in the regular region. The graph shows that the rate of information production generally increases as this distance increases. That is, the quantum system becomes more chaotic as the means are moved away from the elliptic fixed point of the classical map. This is broadly

similar to the classical situation, where the elliptic fixed point is surrounded by a regular region, which is surrounded by a sea of chaos. What is striking is that in the quantum case there is a much smoother transition between the two regimes. Instead of a sharp chaotic and a sharp regular region there is a continuous transition between chaotic and regular behavior.

Furthermore, although R has not been computed, using the inequality (7) we can place an approximate lower bound on R , corresponding to the numerically computed value of R_S . We see that R is always positive for the initial states plotted in figure 2.

Two measures of quantum chaos have been developed in this Letter, and numerical evidence for the existence of a smooth transition between chaotic and regular behaviour in the quantum kicked top presented. The chief advantage of using measures determined wholly by the measurement record is that they can be directly related to the data available in an experiment. Technically, the approach to quantum chaos sketched in this Letter may appear similar to that of Schack, Caves and co-workers [12,16], in the sense that algorithmic information theory is used in both approaches. However, physically the two approaches are quite different. The present approach uses measures determined by the measurement record, whereas Schack, Caves and co-workers focus on measures determined by the state vector or density operator of the quantum system.

The reader may object that in the present approach to quantum chaos an appropriate choice of observation scheme may induce or suppress chaos in the measurement record of systems whose classical analogues are, respectively, chaotic or not chaotic. For example, the quantum Zeno effect (see [17] for an overview and references) predicts the complete suppression of quantum dynamics when appropriate measurements are performed on a quantum system. Such a system will never show chaos in the measurement record, according to our definitions, even if it is classically chaotic. Other examples may also be found [18]. A brief answer to such objections is that the degree of chaos depends on all three of the initial state, the free dynamics, and the observation scheme. To obtain classical results we must assume the system is being observed in an appropriate "classical" fashion. This is certainly not the case for the quantum Zeno effect. However, this should not be taken to imply that observation is the sole cause of chaos in the measurement record. As we have seen, chaos in the measurement record depends also on the system's initial condition, and it can be shown [18] that it depends on the free dynamics as well. These issues will be dealt with more completely in [18].

The approach to quantum chaos sketched in this Letter may be extended in a number of ways. More general measurement schemes may be considered, using formalisms such as quantum trajectories [19,22]. In partic-

ular, it is necessary to consider in perfect measurements, which introduce extra noise into the measurement record. Closely related is the idea of studying the effect of interactions with the environment other than the coupling to a measuring device. Finally, it is interesting to examine more thoroughly how varying the initial state, the system Hamiltonian and the observation scheme affect the degree of chaos seen in the measurement record. These and other issues will be dealt with in a more extensive paper [18].

ACKNOWLEDGMENTS

I would like to thank J.K. Breslin, J. Twamley and G. J. Milburn for many useful discussions about this work. Thanks also to C.M. Caves and Matthew R. Semak who read the manuscript and provided helpful comments. This work was supported by an Australian Postgraduate Award and a Fulbright Scholarship.

- [18] M. A. Nielsen, in preparation. See also the web page <http://tangelo.phys.unm.edu/~mnielsen/qc.html>.
- [19] C.W. Gardiner, A.S. Parkins, and P. Zoller, Phys. Rev. A 46, 4363 (1992).
- [20] K. Molmer, Y. Castin, and J. Dalibard, J. Opt. Soc. Am. B 10, 524 (1993).
- [21] H. J. Carmichael, An Open Systems Approach to Quantum Optics (Springer-Verlag, Berlin, 1993).
- [22] M. A. Nielsen, accepted to Quantum and Semiclassical Optics (1996).
- [23] M. D. Davis, The Undecidable (Raven Press, New York, 1965).

FIG 1. Shannon information as a function of time for the kicked top started in the states $|j_i\rangle$ (bottom) and $|\bar{j}_i\rangle$ (top).

FIG 2. Rate of production of Shannon information as a function of distance from the elliptic fixed point in the $X > 0; Y > 0; Z < 0$ octant of the unit sphere.

-
- [1] M. C. Gutzwiller, Chaos in Classical and Quantum Mechanics (Springer-Verlag, New York, 1990).
 - [2] L. E. Reichl, The Transition to Chaos in Conservative Classical Systems (Springer-Verlag, New York, 1992).
 - [3] R. J. Solomono, Inform. Control 7, 1 (1964).
 - [4] A. N. Kolmogorov, Probl. Inf. Trans. 1, 1 (1965).
 - [5] G. J. Chaitin, Information-Theoretic Incompleteness (World Scientific, Singapore, 1992).
 - [6] A. M. Turing, Proc. Lond. Math. Soc. ser. 2 42, 230 (1937), reprinted on p 115 of [23].
 - [7] G. J. Chaitin, Algorithmic Information Theory (Cambridge University Press, Cambridge, 1987).
 - [8] V. M. Alekseev and M. V. Yakobson, Phys. Rep. 75, 287 (1981).
 - [9] R. Schack, submitted to Phys. Rev. E. LANL e-print xxx/hep-th/9409022 (1995).
 - [10] R. B. Ash, Information Theory (Dover, New York, 1965).
 - [11] F. Haake, Quantum Signatures of Chaos (Springer-Verlag, Berlin Heidelberg, 1991).
 - [12] C. M. Caves, in Physical Origins of Time Asymmetry, edited by J. J. Halliwell, J. Perez-Mercader, and W. H. Zurek (Cambridge University Press, Cambridge, 1994).
 - [13] R. Schack and C. M. Caves, Phys. Rev. Lett. 71, 525 (1993).
 - [14] R. Schack, G. M. D'Ariano, and C. M. Caves, Phys. Rev. E 50, 972 (1994).
 - [15] R. Schack and C. M. Caves, submitted to Phys. Rev. E. LANL e-print xyz/chao-dyn/9506002 (1995).
 - [16] R. Schack and C. M. Caves, submitted to Phys. Rev. E. LANL e-print xyz/chao-dyn/9506008 (1995).
 - [17] A. Peres, Quantum Theory: Concepts and Methods (Kluwer Academic, Dordrecht, 1993).



