

Time-frequency transfer with quantum fields

Marc-Thierry Jaekel^a and Serge Reynaud^b
(a) *Laboratoire de Physique Théorique, CNRS, ENS, UPS,*
24 rue Lhomond, F75231 Paris Cedex 05 France
(b) *Laboratoire Kastler Brossel, UPMC, ENS, CNRS,*
4 place Jussieu, F75252 Paris Cedex 05 France
(LPTENS 96/02)

Clock synchronisation relies on time-frequency transfer procedures which involve quantum fields. We use the conformal symmetry of such fields to define as quantum operators the time and frequency exchanged in transfer procedures and to describe their transformation under transformations to inertial or accelerated frames. We show that the classical laws of relativity are changed when brought in the framework of quantum theory.

PACS: 03.70 04.60 06.30

As known since the advent of relativity theory [1], time is not an absolute notion. Positions in time of events occurring at different locations in space have to be compared through clock synchronisation procedures. From a practical point of view, electromagnetic signals are used to transfer a time or frequency reference, allowing to compare respectively the time delivered by remote clocks or their clock rates [2]. The quality of these procedures is optimized by using a field pulse having respectively a short duration or a good spectral purity. At some high level of precision, synchronisation procedures have to reach a limit associated with the quantum nature of the signals used in the transfer [3]. Time and frequency of a field pulse cannot be defined simultaneously with an arbitrary precision, so that clock synchronisation meets the problem of localisability of quantum particles [4]. This leads to specific difficulties for evaluating the spacetime dependent frequency shifts arising from the effect of gravitational fields or non inertial frames. At the quantum level, consistency of time and frequency transfers with frame transformations is not ensured a priori by classical covariance. The aim of the present letter is to build up such consistency properties by using the underlying symmetries of quantum fields.

In 1905, Einstein proposed the hypothesis of light quanta [5] and, a few months later, introduced the principle of relativity [1]. In the latter paper, he incidentally noticed that energy and frequency of an electromagnetic field change in the same manner in a transformation from an inertial frame to another, thus implicitly referring to the consistency of the hypothesis of light quanta with the principle of relativity. Two years later, Einstein discovered that clock rates and frequency shifts arising in transformations to accelerated frames depend on position [6]. This allowed him to lay down the equivalence of gravity and acceleration and to predict the existence

of gravitational redshifts. He again showed that energy and frequency change in the same manner in the frame transformation.

In modern quantum theory, the similarity of energy and frequency changes has to be interpreted as an invariance property for particle number. This property is well known for Lorentz transformations, but it is usually not admitted for transformations to accelerated frames. In particular, the commonly considered hyperbolic parametrizations of accelerated frames transform vacuum into a thermal bath [7], spoiling the attempts to accommodate the notions of particle number or vacuum to accelerated frames as well as the efforts to understand the principle of equivalence in the quantum domain [8].

For Lorentz transformations, it is also usually considered that the frequency changes, i.e. the Doppler shifts, are consistent with the spacetime transformations described by the Poincaré group. Although similar properties still hold for accelerated frames in general relativity, they have not to our knowledge been brought in the framework of quantum theory. In the absence of a satisfactory quantum description, one is left in the quite uncomfortable position of having only the application of classical covariance rules at one's disposal.

To be more explicit, let us consider a field used either as a frequency reference or as a time reference to be shared by two remote observers. This reference is in any case a quantity preserved by field propagation. In classical physics, this quantity is the field frequency ω or the light-cone variable $t - \frac{x}{c}$, where t is the time coordinate, x the space coordinate along which the transfer is performed and c the velocity of light. In order to build quantum variables associated with the classical ones and to study how they change in frame transformations, we will make use of symmetries of quantum field theory.

Electromagnetism in four-dimensional spacetime is invariant not only under Lorentz transformations, which fit inertial motions, but also under the larger group of conformal transformations which fit uniformly accelerated motions [9]. It follows that particle number, and hence vacuum, have their definitions invariant under conformal transformations to accelerated frames. In the present letter, we will use these invariance properties to obtain a consistent quantum description of time and frequency transfers and to describe how they are affected by transformations to inertial or accelerated frames. We will

present the discussion for the simple case of a scalar massless field in two-dimensional spacetime. With minor reservations to be indicated in the following, the results can be translated to electromagnetic fields in four-dimensional spacetime, at the price of more technical algebraic manipulations, but with their physical significance preserved.

A free massless scalar field $\phi(t, x)$ in two-dimensional spacetime is the sum of two counterpropagating components:

$$\phi(t, x) = \varphi^+(t - x) + \varphi^-(t + x) \quad (1)$$

From now on, we use natural spacetime units ($c = 1$). In the following, we study only one of the two counterpropagating components, that we simply denote $\varphi(u)$:

$$\begin{aligned} \varphi(u) &= \int_0^\infty \frac{d\omega}{2\pi} \sqrt{\frac{\hbar}{2\omega}} (a_\omega e^{-i\omega u} + a_\omega^\dagger e^{i\omega u}) \\ [a_\omega, a_{\omega'}^\dagger] &= 2\pi\delta(\omega - \omega') \end{aligned} \quad (2)$$

u is the light-cone variable to be shared in a time transfer procedure; in the simple field theory considered here, ω represents the frequency as well as the wavevector; a_ω and a_ω^\dagger are the standard annihilation and creation operators and δ is the Dirac distribution.

The infinitesimal transformations of the light-cone variable u are characterized by relations between two coordinate systems \bar{u} and u or, equivalently, by transformations of the field:

$$\begin{aligned} \bar{u} &= u + \varepsilon_m u^m & \varphi(u) &= \bar{\varphi}(\bar{u}) \\ \delta\varphi(u) &\equiv \bar{\varphi}(u) - \varphi(u) & &= -\varepsilon_m u^m \partial_u \varphi(u) \end{aligned} \quad (3)$$

ε_m is an infinitesimal quantity. All the conformal transformations corresponding to any value of m preserve the propagation equation for a massless field theory in two-dimensional spacetime. Here, we will consider only the transformations corresponding to $m = 0, 1, 2$ which respectively describe translations, dilatations and transformations to accelerated frames. These transformations formally correspond to conformal transformations in four-dimensional spacetime and they are known to preserve vacuum fluctuations [10]. Other two-dimensional conformal transformations, not corresponding to those in four-dimensional spacetime, change vacuum fluctuations, which leads to the emission of radiation from mirrors moving in vacuum with a non-uniform acceleration [11]. Vacuum is invariant not only under the infinitesimal transformations (3), but also under the finite transformations obtained by exponentiation [10], since such transformations form a group. This latter property, which is essential for extending the symmetry associated with the Poincaré group, is not obeyed by hyperbolic parametrizations of accelerated frames. Vacuum is indeed invariant under infinitesimal hyperbolic transformations, but transformed into a thermal state under finite ones [7].

In consistency with the canonical commutation relations (2), the field transformation (3) may be described by commutators with conformal generators defined as moments of the stress tensor [12]:

$$\begin{aligned} \delta\varphi &= \frac{\varepsilon_m}{i\hbar} [T_m, \varphi] \\ T_m &= \int u^m e(u) du \quad e(u) =: (\partial_u \varphi(u))^2 : \end{aligned} \quad (4)$$

The symbol $: \quad :$ prescribes a normal ordering of products of operators, and means that the generators vanish in vacuum. Vacuum invariance under the action of T_0 , T_1 and T_2 ensures the consistency of this definition. The conformal generators are also conserved quantities: the translation generator T_0 is the energy-momentum operator associated with the light-cone variable u ; T_1 corresponds to dilatations and T_2 to transformations to accelerated frames. The commutation relations between the conformal generators [13] are recovered, either by inspecting the composition law for coordinate transformations, or by evaluating quantum commutators:

$$[T_m, T_n] = i\hbar(n - m)T_{m+n-1} \quad m, n = 0, 1, 2 \quad (5)$$

In order to discuss the invariance of the photon number, we introduce spectral decompositions for the conformal generators T_0 and T_1 :

$$\begin{aligned} T_m &= \int_0^\infty \frac{d\omega}{2\pi} \rho_m[\omega] \\ \rho_0[\omega] &= \hbar\omega n_\omega & n_\omega &= a_\omega^\dagger a_\omega \\ \rho_1[\omega] &= \hbar\omega\sqrt{n_\omega}(\partial_\omega\delta_\omega)\sqrt{n_\omega} & a_\omega &= e^{i\delta_\omega}\sqrt{n_\omega} \end{aligned} \quad (6)$$

The density ρ_0 is related to the particle number density n_ω while the density ρ_1 is also related to the phase operators δ_ω . Although the definition of these operators is ambiguous [14], the density ρ_1 is properly defined and it vanishes in vacuum. Notice that the operators $\partial_\omega\delta_\omega$ are hermitian even for non-hermitian definitions of the phases [15], and that they are obtained by differentiating phases δ_ω versus frequency ω in complete analogy with the semiclassical definition of scattering phase-delays [16]. The transformation of the energy-momentum density ρ_0 is easily obtained from the field transformation (3):

$$\begin{aligned} [T_0, \rho_0[\omega]] &= 0 \\ [T_1, \rho_0[\omega]] &= i\hbar\omega\partial_\omega\rho_0[\omega] \\ [T_2, \rho_0[\omega]] &= 2i\hbar\omega\partial_\omega\rho_1[\omega] \end{aligned} \quad (7)$$

ρ_0 is unchanged under T_0 , while it is changed under T_1 through a mapping in the frequency domain equivalent to the Doppler shift. Its transformation under T_2 is determined by the density ρ_1 , thus appearing as a spectral decomposition of the commutator $[T_2, T_0]$ of equation (5). Equations (7) imply that the photon number N is preserved under the three generators:

$$N = \int_0^\infty \frac{d\omega}{2\pi} n_\omega \quad [T_m, N] = 0 \quad (8)$$

Hence, the action of T_2 , like that of T_1 , amounts to a redistribution of particles in the frequency domain without any change of the total particle number. In particular, vacuum, the $N = 0$ state, is preserved under T_2 . The transformations (7) have a simple interpretation when the spacetime distribution of the field may be considered as dispersionless. In this case, the operator $\partial_\omega \delta_\omega$ appearing in eqs. (6) may be semiclassically approximated as the classical light-cone variable u . The Doppler shift under T_1 is then proportional to ω , while the frequency shift under T_2 is proportional to $2\omega u$, in consistency with the classical predictions [6].

We will now write down expressions generalizing the classical transformation laws to the quantum domain. To this aim, we first define operators U and Ω associated with the light-cone variable u and field frequency ω . To emphasize the physical content of the equations, we will denote E the energy, which is also the translation generator, D the dilatation generator and C the generator of conformal transformations to accelerated frames:

$$E \equiv T_0 \quad D \equiv T_1 \quad C \equiv T_2 \quad (9)$$

In the case of a scalar field considered here, or equivalently of spin-0 particles [4], the operator U is simply defined as the center of inertia of field energy [17], that is precisely for any state orthogonal to vacuum ($E \neq 0$) as the ratio of D and E (compare with eqs. (4)):

$$U = \frac{1}{2} \{D, E^{-1}\} \quad (10)$$

We have taken care of the non-commutativity of the generators by symmetrizing the expression; $\{ , \}$ denotes an anticommutator. The operator Ω is then defined as the ratio of the energy E to the particle number N :

$$\Omega = \frac{E}{\hbar N} \quad (11)$$

For a single particle state ($N = 1$), Ω plays exactly the same role as $E = \hbar\Omega$. For a more general state however, the quantum fluctuations of Ω and E have independent meanings, since N also possesses its proper quantum fluctuations. Ω is thus a new quantum concept which represents the mean frequency of the field quanta. Note that N always commutes with U as well as with Ω :

$$[N, U] = [N, \Omega] = 0 \quad (12)$$

Having given definitions for the operators U and Ω , we discuss in the following how these definitions are affected by frame transformations. To avoid any ambiguity, we may first recall that the light-cone variable u and the frequency ω are preserved by field propagation in a classical analysis. This property is still true in the present

quantum analysis, where the operators U and Ω are conserved quantities, which may thus be used to transfer time or frequency information between two remote observers. However these operators are not invariant under frame transformations. According to the discussion in the introduction, their transformations are expected to reveal the basic relativistic properties of time and frequency, within the framework of quantum theory.

The commutator $[D, E]$ (see eqs (5,9)):

$$[D, E] = -i\hbar E \quad (13)$$

implies that the operator U transforms under E and D as the classical variable u in the corresponding frame transformations [4]:

$$[E, U] = i\hbar \quad [D, U] = i\hbar U \quad (14)$$

This means in particular that U is canonically conjugated to E . The commutator $[C, E]$ then reads as:

$$[C, E] = -2i\hbar D = -i\hbar \{E, U\} \quad (15)$$

The frequency shifts are finally given by equations (13,15) combined with the invariance (8) of N :

$$[D, \Omega] = -i\hbar \Omega \quad [C, \Omega] = -i\hbar \{\Omega, U\} \quad (16)$$

These laws reproduce the Doppler shifts associated with Lorentz transformations as well as the position dependent frequency shifts arising in transformations to accelerated frames. They fit the form of the classical transformation laws [6], while holding in any quantum state orthogonal to vacuum. As already alluded to in the introduction, the consistency between energy change and frequency change in frame transformations reflects the invariance of the particle number.

These results bring the derivation of frequency shifts in the framework of quantum theory. Precisely, frequency shifts may be evaluated from the quantum transformation laws (16), which are identical to the classical laws, but do not merely rely upon a classical covariance rule. A fact of great interest for the physical analysis of time-frequency transfer is that these expressions are available in the same theoretical framework where quantum fluctuations of the various physical quantities may be analyzed. They may thus be considered as setting the quantum limits in time-frequency transfer. The canonical commutator $[E, U]$ may indeed be read:

$$[\Omega, U] = \frac{i}{N} \quad (17)$$

In the limiting case of a large number of particles, this commutator goes to 0. This allows to build field pulses with nearly dispersionless distributions of Ω and U and to perform time-frequency transfer in a semiclassical regime.

We have found that the transformations (14) of the position operator U under the generators E and D , as well

as the transformations (16) of the frequency operator Ω under all generators have the simple form required by 'classical relativity' [6]. We show now that the transformation of U under C does not conform to these classical covariance rules. To this aim, we introduce the following quadratic form Δ^2 of the generators, which is a Casimir invariant of the conformal algebra (5):

$$\begin{aligned} \Delta^2 &= \frac{1}{2} \{C, E\} - D^2 \\ [E, \Delta^2] &= [D, \Delta^2] = [C, \Delta^2] = 0 \end{aligned} \quad (18)$$

Definitions (10,18) and commutation relations (5) of the conformal algebra allow to rewrite the generator C and its action on the operator U as:

$$\begin{aligned} C &= UEU + \frac{\Delta^2}{E} + \frac{\hbar^2}{4E} \\ [C, U] &= i\hbar \left(U^2 - \frac{\Delta^2}{E^2} - \frac{\hbar^2}{4E^2} \right) \end{aligned} \quad (19)$$

The first term in each expression corresponds to the symmetric ordering of their classical analogs. The other terms are corrections associated with the pulse duration, as it follows from the relation:

$$\frac{1}{2} \int du \{ (U - u)^2, e(u) \} = \frac{1}{E} \left(\Delta^2 + \frac{\hbar^2}{4} \right) \quad (20)$$

It can be shown that Δ^2 has a non negative mean value in any field state, and that it vanishes in any 1-particle state. It may in principle be made close to 0 either by using a very short pulse or by using 1-particle pulse. In contrast, the terms proportional to \hbar^2 appear as purely quantum corrections to the classical terms. They only become negligible at the semiclassical limit where a large number of particles is used ($N \gg 1$).

The transformation (19) for the operator U under the acceleration generator C differs from its classical covariant analog. Corrections however involve operators which commute with E , and are therefore unchanged if the pulse used for the transfer is delayed. It may be stated equivalently that time transfer procedures are found to be invariant under time translation. This statement is the expression of the consistency between time and frequency transfers in the quantum domain. It does however not imply that the corrections still disappear when successive transfer operations are performed. In this case, a sequence of field pulses has indeed to be used and the corrections may vary from one pulse to the next one.

The problem raised here is not a practical limitation in present time-frequency metrology, even at the state-of-the-art level. As a matter of principle however, we emphasize once more that clock synchronisation has to involve quantum fields. As a consequence, the consistency of synchronisation operations, which follows in the

quantum domain from conformal symmetry, entails a departure from the classical covariant laws for frame transformations.

The common conception of spacetime associated with the theory of general relativity is known to remain ambiguous [18]. The result of the present letter extends the connection between symmetries of quantum fields and relativistic properties of spacetime from inertial frames to accelerated frames. It thus advocates a novel conception of spacetime which would be free from its difficulties inherited from classical physics [19].

-
- [1] A. Einstein, *Annalen der Physik* **17** 891 (1905).
 - [2] See for example the Special Issue on Time and Frequency of the *Proceedings of IEEE* **79** 891-1079 (1991).
 - [3] H. Salecker and E.P. Wigner, *Physics Review* **109** 571 (1958).
 - [4] T.D. Newton and E. Wigner, *Review of Modern Physics* **21** 400 (1949).
 - [5] A. Einstein, *Annalen der Physik* **17** 132 (1905).
 - [6] A. Einstein, *Jahrb. Radioakt. Elektron.* **4** 411 (1907).
 - [7] W.G. Unruh, *Physical Review* **D14** 870 (1976); N.D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space* (Cambridge, 1982) and references therein.
 - [8] V.L. Ginzburg and V.P. Frolov, *Sov. Physics Uspekhi* **30** 1073 (1987) [*Uspekhi Fiz. Nauk* **153** 633 (1987)].
 - [9] H. Bateman, *Proceedings of the London Mathematical Society* **8** 223 (1909); E. Cunningham, *Proceedings of the London Mathematical Society* **8** 77 (1909); T. Fulton, F. Rohrlich and L. Witten, *Nuovo Cimento* **26** 653 (1962) and references therein.
 - [10] M.T. Jaekel and S. Reynaud, *Quantum and Semiclassical Optics* **7** 499 (1995).
 - [11] S.A. Fulling and P.C.W. Davies, *Proceedings of the Royal Society* **A348** 393 (1976).
 - [12] C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw Hill, 1985).
 - [13] C. Itzykson and J.M. Drouffe, *Statistical Field Theory* (Cambridge University Press, 1989).
 - [14] S.M. Barnett and D.T. Pegg, *Journal of Physics* **A19** 3849 (1986) and references therein.
 - [15] M.T. Jaekel and S. Reynaud, *Brazilian Journal of Physics* **25** (December 1995).
 - [16] E.P. Wigner, *Physical Review* **98** 145 (1955).
 - [17] A. Einstein, *Annalen der Physik* **20** 627 (1906).
 - [18] J.D. Norton, *Reports Progress Physics* **56** 791 (1993).
 - [19] C. Rovelli *Classical Quantum Gravity* **8** 297, 317 (1991).