arXiv:quant-ph/9605004v1 3 May 1996

ACTION AND PASSION AT A DISTANCE An Essay in Honor of Professor Abner Shim ony

Sandu Popescu

Department of Physics, Boston University, Boston, MA 02215, U.S.A.

Daniel Rohrlich

School of P hysics and A stronom y, Tel-A viv U niversity, Ram at-A viv 69978 Tel-A viv, Israel (M arch 26, 2022)

Abstract

Quantum mechanics permits nonlocality | both nonlocal correlations and nonlocal equations of motion | while respecting relativistic causality. Is quantum mechanics the unique theory that reconciles nonlocality and causality? We consider two models, going beyond quantum mechanics, of nonlocality | \superquantum " correlations, and nonlocal \jamming" of correlations | and derive new results for the jamming model. In one space dimension, jamming allows reversal of the sequence of cause and elect; in higher dimensions, however, elect never precedes cause.

To appear in Quantum Potentiality, Entanglement, and Passion-at-a-Distance: Essays for Abner Shimony, R.S.Cohen, M.A.Home and J.Stachel, eds. (Dordrecht, Netherlands: Kluwer A cademic Publishers), in press.

W hy is quantum mechanics what it is? M any a student has asked this question. Some physicists have continued to ask it. Few have done so with the passion of A bner Shim ony. \W hy is quantum mechanics what it is?" we, too, ask ourselves, and of course we haven't got an answer. But we are working on an answer, and we are honored to dedicate this work to you, A bner, on your birthday.

W hat is the problem ? Q uantum mechanics has an axiom atic structure, exposed by von Neum ann, D irac and others. The axiom s of quantum mechanics tell us that every state of a system corresponds to a vector in a complex H ilbert space, every physical observable corresponds to a linear herm it ian operator acting on that H ilbert space, etc. W e see the problem in comparison with the special theory of relativity. Special relativity can be deduced in its entirety from two axiom s: the equivalence of inertial reference frames, and the constancy of the speed of light. B oth axiom s have clear physical meaning. By contrast, the num erous axiom s of quantum mechanics have no clear physical meaning. D expite m any attempts, starting with von Neumann, to derive the H ilbert space structure of quantum mechanics from a \quantum logic", the new axiom s are hardly more natural than the old.

Abner Shim ony o ershope, and a di erent approach. H is point of departure is a rem arkable property of quantum m echanics: nonlocality. Quantum correlations display a subtle nonlocality. On the one hand, as Bell [1] showed, quantum correlations could not arise in any theory in which all variables obey relativistic causality [2]. On the other hand, quantum correlations them selves obey relativistic causality | we cannot exploit quantum correlations to transm it signals at superlum inal speeds [3] (or at any speed). That quantum m echanics com bines nonlocality and causality is wondrous. Nonlocality and causality seem prim a facie incom patible. E instein's causality contradicts Newton's action at a distance. Yet quantum correlations do not perm it action at a distance, and Shim ony [4] has aptly called the nonlocality m anifest in quantum correlations \passion at a distance". Shim ony has raised the question whether nonlocality and causality can peacefully coexist in any other theory

besides quantum mechanics [4,5].

Quantum mechanics also implies nonlocal equations of motion, as Yakir A haronov [6,7] has pointed out. In one version of the A haronov-Bohm e ect [8], a solenoid carrying an isolated magnetic ux, inserted between two slits, shifts the interference pattern of electrons passing through the slits. The electrons therefore obey a nonlocal equation of motion: they never pass through the ux yet the ux a ects their positions when they reach the screen [9]. A haronov has shown that the solenoid and the electrons exchange a physical quantity, the modular momentum, nonlocally. In general, modular momentum is measurable and obeys a nonlocal equation of motion. But when the ux is constrained to lie between the slits, its modular momentum is completely uncertain, and this uncertainty is just su cient to keep us from seeing a violation of causality. Nonlocal equations of motion at a distance, but quantum mechanics manages to respect relativistic causality. Still, nonlocal equations of motion seem so contrary to relativistic causality that A haronov [7] has asked whether quantum mechanics is the unique theory combining them.

The parallel questions raised by Shimony and Aharonov lead us to consider models for theories, going beyond quantum mechanics, that reconcile nonlocality and causality. Is quantum mechanics the only such theory? If so, nonlocality and relativistic causality together in ply quantum theory, just as the special theory of relativity can be deduced in its entirety from two axioms [7]. In this paper, we will discuss model theories [10{12] manifesting nonlocality while respecting causality. The mst model manifests nonlocality in the sense of Shimony: nonlocal correlations. The second model manifests nonlocality in the sense of Aharonov: nonlocal dynamics. We might that quantum mechanics is not the only theory that reconciles nonlocality and relativistic causality. These models raise new theoretical and experimental possibilities. They imply that quantum mechanics is only one of a class of theories combining nonlocality and causality; in some sense, it is not even the most nonlocal of such theories. Our models raise a question: W hat is the minimal set of physical principles | \nonlocality plus no signalling plus something else simple and fundamental" as Shimony put it [13] from which we may derive quantum mechanics?

The C lauser, H ome, Shim ony, and H olt [14] form of Bell's inequality holds in any classical theory (that is, any theory of local hidden variables). It states that a certain combination of correlations lies between -2 and 2:

2
$$E(A;B) + E(A;B^{0}) + E(A^{0};B) = (A^{0};B^{0}) 2 :$$
 (1)

Besides 2, two other numbers, 2^{p} and 4, are important bounds on the CHSH sum of correlations. If the four correlations in Eq. (1) were independent, the absolute value of the sum could be as much as 4. For quantum correlations, however, the CHSH sum of correlations is bounded [15] in absolute value by 2^{p} . Where does this bound come from? Rather than asking why quantum correlations violate the CHSH inequality, we might ask why they do not violate it more. Suppose that quantum nonlocality in plies that quantum correlations violate the CHSH inequality at least som etim es. W e m ight then guess that relativistic causality is the reason that quantum correlations do not violate it maxim ally. C ould relativistic causality restrict the violation to 2^{2} instead of 4? If so, then nonlocality and causality would together determ ine the quantum violation of the CHSH inequality, and we would be closer to a proof that they determ ine all of quantum mechanics. If not, then quantum mechanics cannot be the unique theory combining nonlocality and causality. To answer the question, we ask what restrictions relativistic causality imposes on pint probabilities. Relativistic causality forbids sending messages faster than light. Thus, if one observer measures the observable A, the probabilities for the outcom es A = 1 and 1 must be independent of whether the other observer chooses to measure B or B^{0} . A =However, it can be shown [10,16] that this constraint does not lim it the CHSH sum of quantum correlations to 2^{p} . For example, in agine a \superguantum " correlation function E for spin measurements along given axes. A ssume E depends only on the relative angle

between axes. For any pair of axes, the outcom es j""i and j##i are equally likely, and sim ilarly for j"#i and j#"i. These four probabilities sum to 1, so the probabilities for j"#i and j##i sum to 1=2. In any direction, the probability of j#i or j#i is 1=2 irrespective of a measurement on the other particle. Measurements on one particle yield no information about measurements on the other, so relativistic causality holds. The correlation function then satis es E () = E (). Now let E () have the form

(i) E () = 1 for 0 = 4;

(ii) E () decreases m onotonically and sm oothly from 1 to -1 as increases from =4 to 3 = 4;

(iii) E() = 1 for 3 = 4

Consider four m easurements along axes de ned by unit vectors \hat{a} , \hat{b} , \hat{a} , and \hat{b}^0 separated by successive angles of =4 and lying in a plane. If we now apply the CHSH inequality Eq. (1) to these directions, we not that the sum of correlations

$$E(\hat{a};\hat{b}) + E(\hat{a}^{0};\hat{b}) + E(\hat{a};\hat{b}^{0}) = (\hat{a}^{0};\hat{b}^{0}) = 3E(=4) = 4$$
 (2)

violates the CHSH inequality with the maximal value 4. Thus, a correlation function could satisfy relativistic causality and still violate the CHSH inequality with the maximal value 4.

III. NONLOCALITY II: NONLOCAL EQUATIONS OF MOTION

A libough quantum mechanics is not the unique theory combining causality and nonlocal correlations, could it be the unique theory combining causality and nonlocal equations of motion? Perhaps the nonlocality in quantum dynamics has deeper physical sign cance. Here we consider a model that in a sense combines the two form sofnonlocality: nonlocal equations of motion where one of the physical variables is a nonlocal correlation. Jamming, discussed by Grunhaus, Popescu and Rohrlich [11] is such a model. The jamming paradigm involves three experimenters. Two experimenters, call them A lice and Bob, make measurements on systems that have locally interacted in the past. A lice's measurements are spacelike separated from Bob's. A third experimenter, Jim (the jammer), presses a button on a black box. This event is spacelike separated from A lice's measurements and from Bob's. The

black box acts at a distance on the correlations between the two sets of systems. For the sake of de niteness, let us assume that the systems are pairs of spin-1/2 particles entangled in a singlet state, and that the measurements of A lice and B ob yield violations of the C H SH inequality, in the absence of jamming; but when there is jamming, their measurements yield classical correlations (no violations of the C H SH inequality).

Indeed, Shim ony [4] considered such a paradigm in the context of the experiment of A spect, D alibard, and R oger [17]. To probe the implications of certain hidden-variable theories [18], hew rote, \Suppose that in the interval after the commutators of that experiment have been actuated, but before the polarization analysis of the photons has been completed, a strong burst of laser light is propagated transverse to but intersecting the paths of the propagating photons.... Because of the nonlinearity of the fundamental material medium which has been postulated [in these models], this burst would be expected to generate excitations, which could conceivably interfere with the nonlocal propagation that is responsible for polarization correlations." Thus, Shim ony asked whether certain hidden-variable theories would predict classical correlations after such a burst. (Q uantum mechanics, of course, does not.)

Here, our concern is not with hidden-variable theories or with a mechanism for jamming; rather, we ask whether such a nonlocal equation of motion (or one, say, allowing the third experimenter nonlocally to create, rather than jam, nonlocal correlations) could respect causality. The jamming model [11] addresses this question. In general, jamming would allow Jim to send superluminal signals. But remarkably, some forms of jamming would not; Jim could tamper with nonlocal correlations without violating causality. Jamming preserves causality if it satis es two constraints, the unary condition and the binary condition. The unary condition states that Jim cannot use jamming to send a superluminal signal that A lice (or B ob), by examining her (or his) results alone, could read. To satisfy this condition, let us assume that A lice and B ob each measure zero average spin along any axis, with or without jamming. In order to preserve causality, jamming must a ect correlations only, not average measured values for one spin component. The binary condition states that Jim cannot use

jam ming to send a signal that A lice and B ob together could read by comparing their results, if they could do so in less time than would be required for a light signal to reach the place where they meet and compare results. This condition restricts spacetime congurations for jamming. Let a, b and j denote the three events generated by A lice, B ob, and Jim, respectively: a denotes A lice's measurements, b denotes B ob's, and j denotes Jim's pressing of the button. To satisfy the binary condition, the overlap of the forward light cones of a and b must lie entirely within the forward light cone of j. The reason is that A lice and B ob can compare their results only in the overlap of their forward light cones. If this overlap is entirely contained in the forward light cone of j, then a light signal from j can reach any point in spacetime where A lice and B ob can compare their results. This restriction on jam ming congurations also rules out another violation of the unary condition. If Jim could obtain the results of A lice's measurements prior to deciding whether to press the button, he could send a superlum inal signal to B ob by selectively jamming [11].

IV.AN EFFECT CAN PRECEDE ITS CAUSE!

If jam ming satis es the unary and binary conditions, it preserves causality. These conditions restrict but do not preclude jam ming. There are congurations with spacelike separated a, b and j that satisfy the unary and binary conditions. We conclude that quantum mechanics is not the only theory combining nonlocal equations of motion with causality. In this section we consider another remarkable aspect of jam ming, which concerns the time sequence of the events a, b and j de ned above. The unary and binary conditions are manifestly Lorentz invariant, but the time sequence of the events a, b and j is not. A time sequence a, j, b in one Lorentz fram e may transform into b, j, a in another Lorentz fram e. Furtherm ore, the jam ming model presents us with reversals of the sequence of cause and e ect: while j may precede both a and b in one Lorentz fram e, in another fram e both a and b may precede j.

To see how jamming can reverse the sequence of cause and e ect, we specialize to the

case of one space dimension. Since a and b are spacelike separated, there is a Lorentz frame in which they are simultaneous. Choosing this frame and the pair (x;t) as coordinates for space and time, respectively, we assign a to the point (-1,0) and b to the point (1,0). W hat are possible points at which j can cause jamming? The answer is given by the binary condition. It is particularly easy to apply the binary condition in 1+1 dimensions, since in 1+1 dimensions the overlap of two light cones is itself a light cone. The overlap of the two forward light cones of a and b is the forward light cone issuing from (0,1), so the jammer, Jim, may act as late as t = 1 after A lice and Bob have completed their measurements and still jam their results. More generally, the binary condition allows us to place j anywhere in the backward light cone of (0,1) that is also in the forward light cone of (0,-1), but not on the boundaries of this region, since we assume that a, b and j are mutually spacelike separated. (In particular, j cannot be at (0,1) itself.)

Such reversals m ay boggle the m ind, but they do not lead to any inconsistency as long as they do not generate self-contradictory causal loops [19,20]. Consistency and causality are intim ately related. We have used the term relativistic causality for the constraint that others call no signalling. W hat is causal about this constraint? Suppose that an event (a \cause") could in uence another event (an \e ect") at a spacelike separation. In one Lorentz fram e the cause precedes the e ect, but in some other Lorentz frame the e ect precedes the cause; and if an e ect can precede its cause, the e ect could react back on the cause, at a still earlier time, in such a way as to prevent it. A self-contradictory causal loop could arise. A man could kill his parents before they met. Relativistic causality prevents such causal contradictions [19]. Jam m ing allows an event to precede its cause, but does not allow selfcontradictory causal bops. It is not hard to show [11] that if jamming satis es the unary and binary conditions, it does not lead to self-contradictory causal loops, regardless of the number of jammers. Thus, the reversal of the sequence of cause and e ect in jamming is consistent. It is, how ever, su ciently remarkable to warrant further comment below, and we also show that the sequence of cause and e ect in jamming depends on the space dimension in a surprising way.

The unary and binary conditions restrict the possible jam ming congurations; however, they do not require that jam ming be allowed for all congurations satisfying the two conditions. Nevertheless, we have made the natural assumption that jam ming is allowed for all such congurations. This assumption is manifestly Lorentz invariant. It allows a and b to both precede j. In a sense, it means that Jim acts along the backward light cone of j; whenever a and b are outside the backward light cone of j and full 11 the unary and binary conditions, jam ming occurs.

V.AN EFFECT CAN PRECEDE ITS CAUSE??

That Jim may act after A lice and B ob have completed their measurements (in the given Lorentz frame) is what may boggle the mind. How can Jim change his own past? We may also put the question in a dierent way. Once A lice and Bob have completed their measurements, there can after all be no doubt about whether or not their correlations have been jammed; A lice and Bob cannot compare their results and nd out until after Jim has already acted, but whether or not jamming has taken place is already an immutable fact. This fact apparently contradicts the assumption that Jim is a free agent, i.e. that he can freely choose whether or not to jam. If A lice and B ob have completed their measurements, Jim is not a free agent: he must push the button, or not push it, in accordance with the results of A lice and B ob's measurements.

W e m ay be uncom fortable even if Jim acts before A lice and B ob have both com pleted their m easurements, because the time sequence of the events a, b and j is not Lorentz invariant; a, j, b in one Lorentz fram e m ay transform to b, j, a in another. The reversal in the time sequences does not lead to a contradiction because the e ect cannot be isolated to a single spacetime event: there is no observable e ect at either a or b, only correlations between a and b are changed. All the same, if we assume that Jim acts on either A lice or B ob | whoever m easures later | we conclude he could not have acted on either of them, because both come earlier in some Lorentz fram e.

W hat, then, do we make of cause and e ect in the jam ming model? We o er two points of view on this question. One point of view is that we don't have to worny; jam ming does not lead to any causal paradoxes, and that is all that matters. O fcourse, experience teaches that causes precede their e ects. Yet experience also teaches that causes and e ects are locally related. In jam ming, causes and e ects are nonlocally related. So we cannot assum e that causes must precede their e ects; it is contrary to the spirit of special relativity to im pose such a dem and. Indeed, it is contrary to the spirit of general relativity to assign absolute meaning to any sequence of three mutually spacelike separated events, even when such a sequence has a Lorentz-invariant meaning in special relativity [20]. We only dem and that no sequence of causes and e ects close upon itself, for a closed causal loop | a timetravel paradox | would be self-contradictory. If an e ect can precede its cause and both are spacetime events, then a closed causal loop can arise. But in jam ming, the cause is a spacetime event and the e ect involves two spacelike separated events; no closed causal loop can arise [11].

This point of view interprets cause and e ect in jam m ing as Lorentz invariant; observers in all Lorentz fram es agree that jam m ing is the e ect and Jim's action is the cause. A second point of view asks whether the jam m ing m odel could have any other interpretation. In a world with jam m ing, m ight observers in di erent Lorentz fram es give di erent accounts of jam m ing? Could a sequence a, j, b have a covariant interpretation, with two observers com ing to di erent conclusions about which m easurem ents were a ected by Jim? (No experim ent could ever prove one of them w rong and the other right [21].) Likew ise, perhaps observers in a Lorentz fram e where both a and b precede j would interpret jam m ing as a form of telesthesia: Jim knows whether the correlations m easured by A lice and B ob are nonlocal before he could have received both sets of results. W e m ust assum e, how ever, that observers in such a world would notice that jam m ing always turns out to bene t Jim; they would not interpret jam m ing as m ere telesthesia, so the jam m ing m odel could not have this covariant interpretation.

Finally, we note that a question of interpreting cause and e ect arises in quantum me-

chanics, as well. Consider the measurements of A lice and Bob in the absence of jamming. Their measured results do not indicate any relation of cause and e ect between A lice and Bob; A lice can do nothing to a ect Bob's results, and vice versa. A coording to the conventional interpretation of quantum mechanics, however, the rst measurement on a pair of particles entangled in a singlet state causes collapse of the state. The question whether A lice or B ob caused the collapse of the singlet state has no Lorentz-invariant answer [11,22].

VI. JAMM ING IN MORE THAN ONE SPACE DIMENSION

A fler arguing that jam ming is consistent even if it allows reversals of the sequence of cause and e ect, we open this section with a surprise: such reversals arise only in one space dimension! In higher dimensions, the binary condition itself eliminates such con qurations; jam ming is not possible if both a and b precede j. To prove this result, we rst consider the case of 2+1 dimensions. We choose coordinates (x;y;t) and, as before, place a and b on the x-axis, at (-1,0,0) and (1,0,0), respectively. Let A, B and J denote the forward light cones of a, b and j, respectively. The surfaces of A and B intersect in a hyperbola in the yt-plane. To satisfy the binary condition, the intersection of A and B must lie entirely within J. Suppose that this condition is ful lled, and now we move j so that the intersection of A and B ceases to lie within J. The intersection of A and B ceases to lie within J when its surface touches the surface of J. Either a point on the hyperbola, or a point on the surface of either A of B alone, may touch the surface of J. However, the surfaces of A and J can touch only along a null line (and likewise for B and J); that is, only if j is not spacelike separated from either a or b, contrary to our assumption. Therefore the only new constraint on j is that the hyperbola form ed by the intersection of the surfaces of A and B not touch the surface of J. If we place j on the t-axis, at (0,0,t), the latest time t for which this condition is ful led is when the asymptotes of the hyperbola lie along the surface of J. They lie along the surface of J when j is the point (0,0,0). If j is the point (0,0,0), moving j in either the x-or y-direction will cause the hyperbola to intersect the surface of J. W e conclude that

there is no point j, consistent with the binary condition, with t-coordinate greater than 0. Thus, j cannot succeed both a and b in any Lorentz fram e (although it could succeed one of them).

For n > 2 space dimensions, the proof is similar. The only constraint on j arises from the intersection of the surfaces of A and B. At a given time t, the surfaces of A and B are $(n \ 1)$ -spheres of radius t centered, respectively, at x = 1 and x = 1 on the x-axis; these $(n \ 1)$ -spheres intersect in an $(n \ 2)$ -sphere of radius $(t^2 \ 1)^{1=2}$ centered at the origin. This $(n \ 2)$ sphere lies entirely within an $(n \ 1)$ -sphere of radius t centered at the origin, and approaches it asymptotically for $t \ 1$. The $(n \ 1)$ -spheres centered at the origin are sections of the forward light cone of the origin. Thus, j cannot occur later than a and b.

We nd this result both am using and odd. We argued above that allowing j to succeed both a and b does not entail any inconsistency and that it is contrary to the spirit of the general theory of relativity to exclude such con gurations for jam ming. Nonetheless, we nd that they are autom atically excluded for n 2.

VII.CONCLUSIONS

Two related questions of Shim ony [4,5] and Aharonov [7] inspire this essay. Nonbcality and relativistic causality seem almost irreconcilable. The emphasis is on almost, because quantum mechanics does reconcile them, and does so in two dierent ways. But is quantum mechanics the unique theory that does so? Our answer is that it is not: model theories going beyond quantum mechanics, but respecting causality, allow nonlocality both ways. We qualify our answer by noting that nonlocality is not completely de ned. Relativistic causality is well de ned, but nonlocality in quantum mechanics includes both nonlocal correlations and nonlocal equations of motion, and we do not know exactly what kind of nonlocality we are sæking. A Itematively, we may ask what additional physical principles can we in pose that will single out quantum mechanics as the unique theory. Our \superquantum " and \jmm ing" models open new experimental and theoretical possibilities. The superquantum

m odel predicts violations of the CHSH inequality exceeding quantum violations, consistent with causality. The jamming model predicts new elects on quantum correlations from some mechanism such as the burst of laser light suggested by Shimony [4]. Most interesting are the theoretical possibilities. They oller hope that we may rediscover quantum mechanics as the unique theory satisfying a small number of fundamental principles: causality plus nonlocality \plus som ething else simple and fundamental" [13].

ACKNOW LEDGMENTS

D.R.acknow ledges support from the State of Israel, M inistry of Imm igrant Absorption, Center for Absorption in Science.

REFERENCES

- [1] J.S.Bell, Physics 1, 195 (1964).
- [2] The term relativistic causality denotes the constraint that inform ation cannot be transferred at speeds exceeding the speed of light. This constraint is also called no signalling.
- [3] G.C.Ghirardi, A.Rimini and T.Weber, Lett. Nuovo Cim. 27 (1980) 263.
- [4] A. Shimony, in Foundations of Quantum Mechanics in Light of the New Technology, S. Kamefuchiet al., eds. (Tokyo, Japan Physical Society, 1983), p. 225.
- [5] A. Shimony, in Quantum Concepts in Space and Time, R. Penrose and C. Isham, eds.(Oxford, Claredon Press, 1986), p. 182.
- [6] Y.Aharonov, H.Pendleton, and A.Petersen, Int. J. Theo. Phys. 2 (1969) 213; 3 (1970) 443; Y.Aharonov, in Proc. Int. Symp. Foundations of Quantum Mechanics, Tokyo, 1983, p. 10.
- [7] Y. A haronov, unpublished lecture notes.
- [8] Y. Aharonov and D. Bohm, Phys. Rev. 115 (1959) 485, reprinted in F.W ilczek (ed.) Fractional Statistics and Anyon Superconductivity, Singapore: W orld-Scienti c, 1990;
- [9] It is true that the electron interacts locally with a vector potential. However, the vector potential is not a physical quantity; all physical quantities are gauge invariant.
- [10] S. Popescu and D. Rohrlich, Found. Phys. 24, 379 (1994).
- [11] J.G runhaus, S.Popescu and D.Rohrlich, TelAviv University preprint TAUP-2263-95 (1995), to appear in Phys. Rev. A.
- [12] D. Rohrlich and S. Popescu, to appear in the Proceedings of 60 Years of E. P. R. (Workshop on the Foundations of Quantum Mechanics, in honor of Nathan Rosen), Technion, Israel, 1995.

[13] A. Shim ony, private communication.

- [14] J.F.C lauser, M.A. Home, A. Shim ony and R.A. Holt, Phys. Rev. Lett. 23, 880 (1969).
- [15] B. S. Tsirelson (Cirel'son), Lett. M ath. Phys. 4 (1980) 93; L. J. Landau, Phys. Lett. A 120 (1987) 52.
- [16] For the maxim alviolation of the CHSH inequality consistent with relativity see also L. K hal n and B.T sirelson, in Sym posium on the Foundations of Modern Physics '85, P. Lahti et al., eds. (W orld-Scienti c, Singapore, 1985), p. 441; P.R astall, Found. Phys. 15, 963 (1985); S. Sum m ers and R.W erner, J.M ath. Phys. 28, 2440 (1987); G.K renn and K. Svozil, preprint (1994) quant-ph/9503010.
- [17] A. Aspect, J. Dalibard and G. Roger, Phys. Rev. Lett. 49, 1804 (1982).
- [18] D. Bohm, W holeness and the Implicate Order (Routledge and Kegan Paul, London, 1980); D. Bohm and B. Hiley, Found. Phys. 5, 93 (1975); J.P. Vigier, Astr. Nachr. 303, 55 (1982); N. Cufaro-Petroni and J.P. Vigier, Phys. Lett. A 81, 12 (1981); P. Droz-Vincent, Phys. Rev. D 19, 702 (1979); A. Ganuccio, V. A. Rapisarda and J.P. Vigier, Lett. Nuovo Cim. 32, 451 (1981).
- [19] See e.g. D. Bohm, The Special Theory of Relativity, W. A. Benjam in Inc., New York (1965) 156-158.
- [20] We thank Y. A haronov for a discussion on this point.
- [21] They need not be incompatible. An event in one Lorentz fram e offen is another event in another fram e. For example, absorption of a virtual photon in one Lorentz fram e corresponds to emission of a virtual photon in another. In jamming, Jim might not only send instructions but also receive information, in both cases unconsciously. (Jim is conscious only of whether or not he jam s.) Suppose that the time reverse of \sending instructions" corresponds to \receiving information". Then each observer interprets the sequence of events correctly for his Lorentz fram e.

[22] Y.Aharonov and D.Albert, Phys. Rev. D 24, 359 (1981).