

Comment on “Sonoluminescence as Quantum Vacuum Radiation”

We contest the recent claim that sonoluminescence may be explained in terms of quantum vacuum radiation [1]. Due to fundamental physical limitations on bubble surface velocity, the predicted number of photons per flash is indeed much smaller than unity. Therefore, quantum vacuum radiation cannot be considered as an explanation of the observed sonoluminescence phenomenon.

As the starting point of our critical comment we will use the theoretical evaluation of radiation emitted into vacuum by a moving spherical interface between two dielectric media [1,2]. In addition to the results presented in these papers we will calculate the number of photons emitted in each sonoluminescence cycle by interpreting it in terms of parametric emission. Each elementary radiation process corresponds to the emission of a pair of photons (ω, ω') carrying an energy $\hbar(\omega + \omega') = \hbar\Omega$, where Ω is the frequency of the mechanical motion. Equation (10) of [1] or equation (4.8) of [2], which give the energy radiated per pulsation of period T , are thus translated into the following expression of the number of radiated photons

$$\mathcal{N} = \alpha \int_0^\infty d\Omega \Omega^5 \left| \int_0^T d\tau \frac{R^2(\tau)}{c^2} e^{i\Omega\tau} \right|^2 \quad (1)$$

α is a numerical factor ($\alpha \simeq 10^{-4}$ for a water-air interface), and $R(\tau)$ is the time-dependent radius.

An estimate of \mathcal{N} may be derived from a simple lorentzian model (eq. (4.9) of [2] with the same notations for R_0 , R_{\min} and γ)

$$\begin{aligned} \mathcal{N} &= \frac{15\pi^2}{16} \alpha \beta^4 \\ \beta^2 &= \frac{R_0^2 - R_{\min}^2}{c^2 \gamma^2} \end{aligned} \quad (2)$$

The number of photons thus depends on a single parameter β which is the typical velocity of bubble surface measured with respect to the speed of light. The time γ characteristic of bubble collapse, which is assumed to be much shorter than T , determines the frequency spectrum of radiation but does not appear in the integrated photon number. A study of the law of variation of velocity then shows that the photon number (2) may not exceed a value determined by the maximal velocity $\left(\frac{dR}{dt}\right)_{\max}$ of the bubble surface

$$\mathcal{N} \leq .1 \left(\frac{dR}{cdt} \right)_{\max}^4 \quad (3)$$

For any maximal velocity smaller than the speed of light, the photon number (2) therefore remains smaller than unity. In contrast, observed sonoluminescence corresponds to more than 10^5 photons per flash (see for instance Fig.3 of [3]).

Moreover, the characteristic velocity of the bubble surface corresponds to the sound velocity in air or water (see Fig.4 of [4]) rather than to the speed of light. For a maximal velocity equal to sound velocity in water, \mathcal{N} comes out as a very small number of the order of 10^{-23} photons. We may emphasize that, with a radius in the μm range (see Fig.3 of [4]), a velocity of the order of 1km/s leads to a typical time in the ns range, which fits experimental reports (see again Fig.3 of [4]). A much shorter time scale is used in [1] in order to explain the short duration of the flashes and the large width of the thermal-like spectrum. This very short time scale not only differs considerably from the measured values, but it also corresponds to supraluminal velocities.

The precise values of the numerical factors which play a role in the preceding discussion have been obtained from a specific model for the variation of the bubble radius. However, it can hardly be expected that a different variation law of bubble radius may fill the gap between 10^{-23} and 10^5 photons. The predicted number of photons per flash remains much smaller than unity, which unavoidably leads to contest the claim that “the theory of vacuum radiation seems to agree remarkably well with the experimental results on sonoluminescence” [1].

Astrid Lambrecht
Max-Planck-Institut für Quantenoptik
D-85748 Garching, Germany

Marc-Thierry Jaekel
Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, F-75231 Paris Cedex 05, France

Serge Reynaud
Laboratoire Kastler Brossel, Université Pierre et Marie Curie, F-75252 Paris Cedex 05, France

LPTENS 96/40
PACS: 78.60.Mq, 03.70+k, 42.50.Lc

-
- [1] C. Eberlein, *Physical Review Letters* **76** 3842 (1996)
 - [2] C. Eberlein, *Physical Review* **A53** 2772 (1996)
 - [3] R. Hiller, S.J. Putterman and B.P. Barber, *Physical Review Letters* **69** 1182 (1992)
 - [4] R. Löfstedt, B.P. Barber and S.J. Putterman, *Physics of Fluids* **A5** 2911 (1993)