QUANTUM NONLOCALITY AND INSEPARABILITY

A sher Peres

Department of Physics Technion | Israel Institute of Technology 32000 Haifa, Israel

A quantum system consisting of two subsystems is separable if its density matrix can be written as $= {}^{P} w_{K} {}^{0}_{K} {}^{0}_{K}$, where ${}^{0}_{K}$ and ${}^{0}_{K}$ are density matrices for the two subsystems, and the positive weights w_{K} satisfy ${}^{P} w_{K} = 1$. A necessary condition for separability is derived and is shown to be more sensitive than Bell's inequality for detecting quantum inseparability. Moreover, collective tests of Bell's inequality (namely, tests that involve several composite systems simultaneously) may sometimes lead to a violation of Bell's inequality, even if the latter is satisfied when each composite system is tested separately.

1. IN TRODUCTION

From the early days of quantum mechanics, the question has offen been raised whether an underlying \subquantum " theory, that would be determ inistic or even stochastic, was viable. Such a theory would presum ably involve additional \hidden" variables, and the statistical predictions of quantum theory would be reproduced by perform ing suitable averages over these hidden variables.

A fundam ental theorem was proved by Bell [1], who showed that if the constraint of locality was in posed on the hidden variables (nam ely, if the hidden variables of two distant quantum systems were them selves be separable into two distinct subsets), then there was an upper bound to the correlations of results of measurem ents that could be perform ed on the two distant systems. That upper bound, mathematically expressed by Bell's inequality [1], is violated by some states in quantum mechanics, for example the singlet state of two spin $\frac{1}{2}$ particles.

A variant of Bell's inequality, more general and more useful for experim ental tests, was later derived by C lauser, Home, Shimony, and Holt (CHSH) [2]. It can be written

$$j_{A}Bi + h_{A}B'i + h_{A}B'i h_{A}B'ij 2:$$
 (1)

On the left hand side, A and A⁰ are two operators that can be measured by an observer, conventionally called A lice. These operators do not commute (so that A lice has to choose whether to measure A or A⁰) and each one is normalized to unit norm (the norm of an operator is dened as the largest absolute value of any of its eigenvalues). Likewise, B and B⁰ are two normalized noncommuting operators, any one of which can be measured by another, distant observer (Bob). Note that each one of the expectation values in Eq. (1) can be calculated by means of quantum theory, if the quantum state is known, and is also experimentally observable, by repeating the measurements su ciently many times, starting each time with identically prepared pairs of quantum states of quantum states in Eq. (1) can be in the easurement of the expectation of the measurements su ciently many times, starting each time with identically prepared pairs of quantum states of quantum states of quantum states in Eq. (1) can be in the easurement of the expectation of the measurement of the easurement of the expectation states is known, and is also experimentally observable, by repeating the measurement of the easurement of the easurement of the expectation of the easurement of the easurement is also experimentally observable, by repeating the measurement of the easurement of the easurem

(LHV) m odel for the results of these m easurem ents. It is not in general a su cient condition, as will be shown below.

Note that, in order to test Bell's inequality, the two distant observers independently measure subsytems of a composite quantum system, and then report their results to a common site where that inform ation is analyzed [3]. A related, but essentially dimerent, issue is whether a composite quantum system can be prepared in a prescribed state by two distant observers who receive instructions from a common source. For this to be possible, the density matrix has to be separable into a sum of direct products,

$$= \bigvee_{K}^{X} w_{K} \bigvee_{K}^{0} \bigvee_{K}^{0} \bigvee_{K}^{0}; \qquad (2)$$

where the positive weights w_K satisfy $^P w_K = 1$, and where 0_K and 0_K are density matrices for the two subsystems. A separable system always satis es Bell's inequality, but the converse is not necessarily true [4{7]. I shall derive below a simple algebraic test, which is a necessary condition for the existence of the decomposition (2). I shall then give some examples showing that this criterion is more restrictive than Bell's inequality, or than the -entropy inequality [8].

2. SEPARABILITY OF DENSITY MATRICES

The derivation of the separability condition is easiest when the density matrix elements are written explicitly, with all their indices β]. For example, Eq. (2) becomes

$$m_{m} = \sum_{K}^{X} w_{K} \left({}_{K}^{0} \right)_{mn} \left({}_{K}^{0} \right) : \qquad (3)$$

Latin indices refer to the subsystem, G reek indices to the second one (the subsystem s m ay have di erent dimensions). Note that this

equation can always be satis ed if we replace the quantum density matrices by classical Liouville functions (and the discrete indices are replaced by canonical variables, p and q). The reason is that the only constraint that a Liouville function has to satisfy is being non-negative. On the other hand, we want quantum density matrices to have nonnegative eigenvalues, rather than non-negative elements, and the latter condition is more di cult to satisfy.

Let us now de ne a new matrix,

The Latin indices of have been transposed, but not the G reek ones. This is not a unitary transform ation but, nevertheless, the matrix is Herm itian. W hen Eq. (2) is valid, we have

$$= \int_{A}^{X} w_{A} \left(\int_{A}^{0} \right)^{T} \qquad \int_{A}^{0} :$$
 (5)

Since the transposed m atrices $\begin{pmatrix} 0\\A \end{pmatrix}^T$ $\begin{pmatrix} 0\\A \end{pmatrix}^T$ $\begin{pmatrix} 0\\A \end{pmatrix}$ are non-negative m atrices with unit trace, they can also be legitimate density matrices. It follows that none of the eigenvalues of is negative. This is a necessary condition for Eq. (2) to hold [9].

Note that the eigenvalues of are invariant under separate unitary transform ations, U 0 and U 0 , of the bases used by the two observers. In such a case, transform s as

$$! (U^{0} U^{0}) (U^{0} U^{0})^{\vee};$$
(6)

and we then have

$$! (U^{OT} U^{OT}) (U^{OT} U^{OT})^{\gamma};$$

$$(7)$$

which also is a unitary transformation, leaving the eigenvalues of invariant.

As an example, consider a pair of $spin-\frac{1}{2}$ particles in an impure singlet state, consisting of a singlet fraction x and a random fraction (1 x) [10]. Note that the \random fraction" (1 x) also includes singlets, m ixed in equal proportions with the three triplet components. We have

$$m_{in} = x S_{m_{in}} + (1 x)_{m_{in}} = 4;$$
 (8)

where the density matrix for a pure singlet is given by

$$S_{01;01} = S_{10;10} = S_{01;10} = S_{10;01} = \frac{1}{2};$$
 (9)

and all the other components of S vanish. (The indices 0 and 1 refer to any two orthogonal states, such as \up" and \down.") A straightforward calculation shows that has three eigenvalues equal to (1 + x)=4, and the fourth eigenvalue is $(1 \quad 3x)=4$. This lowest eigenvalue is positive if $x < \frac{1}{3}$, and the separability criterion is then fulled. This result may be compared with other criteria: Bell's inequality holds for $x < 1=\frac{p}{2}$, and the -entropic inequality [8] for $x < 1=\frac{p}{3}$. These are therefore much weaker tests for detecting inseparability than the condition that was derived here.

In this particular case, it happens that this necessary condition is also a su cient one. It is indeed known that if $x < \frac{1}{3}$ it is possible to write as a mixture of unentangled product states [11]. This suggests that the necessary condition derived above (has no negative eigenvalue) might also be su cient for any . A proof of this conjecture was indeed recently obtained [12] for composite systems having dimensions 2 2 and 2 3. However, for higher dimensions, the present necessary condition was shown not to be a su cient one.

As a second example, consider a m ixed state consisting of a fraction x of the pure state a j(1i + b)(0i) (with $j_a f + b f = 1$), and fractions $(1 \ x)=2$ of the pure states j(0i) and j(1i). We have

$$_{00;00} = _{11;11} = (1 x)=2;$$
 (10)

$$_{01;01} = x j_{a} j^{2};$$
 (11)

$$_{10;10} = x p_{j}^{2};$$
 (12)

$$_{01;10} = _{10;01} =$$
 xab ; (13)

and the other elements of vanish. It is easily seen that the matrix has a negative determ inant, and thus a negative eigenvalue, when

$$x > (1 + 2jabj)^{-1}$$
: (14)

This is a lower limit than the one for a violation of Bell's inequality, which requires [7]

$$x > [1 + 2jabj(2 1)]^{1}$$
: (15)

An even more striking example is the mixture of a singlet and a maximally polarized pair:

$$m_{jn} = x S_{m_{jn}} + (1 \ x)_{m0 \ n0 \ 0}$$
 (16)

For any positive x, however small, this state is inseparable, because has a negative eigenvalue (x=2). On the other hand, the Horodecki criterion [13] gives a very generous domain to the validity of Bell's inequality: x 0.8.

3. COLLECTIVE TESTS FOR NONLOCALITY

The weakness of Bell's inequality as a test for inseparability is due to the fact that the only use made of the density matrix is for computing the probabilities of the various outcomes of tests that may be performed on the subsystems of a single composite system. On the other hand, an experimental veries cation of that inequality necessitates the use of many composite systems, all prepared in the same way. However, if many such systems are actually available, we may also test them collectively, for example two by two, or three by three, etc., rather than one by one. If we do that, we must use, instead of (the density matrix of a single system), a new density matrix, which is , or

, in a higher dimensional space. It will now be shown that there are some density matrices that satisfy Bell's inequality, but for which , or , etc., violate that inequality [14].

The example that will be discussed is that of the W emer states [4] de ned by Eq. (8). Let us consider n W emer pairs. Each one of the two observers has n particles (one from each pair). They proceed as follows. First, they subject their n-particle systems to suitably chosen local unitary transform ations, U, for A lice, and V, for B ob. Then, they test whether each one of the particles labelled 2, 3, ..., n, has spin up (for simplicity, it is assumed that all the particles are distinguishable, and can be labelled unambiguously). Note that any other test that they can perform is unitarily equivalent to the one for spins up, as this involves only a rede nition of the matrices U and V. If any one of the 2 (n 1) particles tested by A lice and B ob shows spin down, the experiment is considered to have failed, and the two observers must start again with n new W emer pairs.

A similar elimination of \bad" samples is also inherent to any experimental procedure where a failure of one of the detectors to re is handled by discarding the results registered by all the other detectors: only when all the detectors re are their results included in the statistics. This obviously requires an exchange of classical information between the observers. (There is a controversy on whether a violation of Bell's inequality with postselected data [15] is a valid test for non-

7

locality [16]. I shall not discuss this issue here; I only exam ine whether or not Bell's inequality is violated by the postselected data.)

The calculations shown below will refer to the case n = 3, for de niteness. The generalization to any other value of n is straightforward. Spinor indices, for a single spin $\frac{1}{2}$ particle, will take the values 0 (for the \up" component of spin) and 1 (for the \down" component). The 16 components of the density matrix of a W emer pair, consisting of a singlet fraction x and a random fraction (1 x), are, in the standard direct product basis:

$$m_{n;st} = x S_{m_{n;st}} + (1 x) m_{s nt} = 4;$$
 (17)

where I am now using only Latin indices, contrary to what I did in Eq.(8); this is because G reek indices will be needed for another purpose, as will be seen soon. Thus, now, the indices m and s refer to A lice's particle, and n and t to Bob's particle.

W hen there are three W erner pairs, their combined density m atrix is a direct product 0 0, or explicitly, m_n;st m_0n_0;s_0t_0 m_0n_0;s_0t_0. The result of the unitary transform ations U and V is

0
 0 ! (U V) (0 0) (U y V y): (18)

Explicitly, with all its indices, the U matrix satis es the unitarity relation

$$U \circ \omega_{\mathfrak{m}\mathfrak{m}} \circ \omega U \circ \omega_{\mathfrak{m}\mathfrak{m}} \circ \omega U \circ \omega_{\mathfrak{m}\mathfrak{m}} \circ \omega = \circ \circ \omega \circ \omega \circ (19)$$

In order to avoid any possible ambiguity, G reek indices (whose values are also 0 and 1) are now used to label spinor components after the unitary transformations. Note that the indices without primes refer to the two particles of the set W emer pair (the only ones that are not tested for spin up) and the primed indices refer to all the other particles (that are tested for spin up). The V $\circ \alpha_{mn} \gamma_{n} \infty$ matrix elements

of Bob's unitary transform ation satisfy a relationship similar to (19). The generalization to a larger number of W erner pairs is obvious.

A first the execution of the unitary transformation (18), A lice and Bob have to test that all the particles, except those labelled by the

rst (unprimed) indices, have their spin up. They discard any set of n W emer pairs where that test fails, even once. The density matrix for the remaining \successful " cases is thus obtained by retaining, on the right hand side of Eq. (18), only the term s whose primed components are zeros, and then renormalizing the resulting matrix to unit trace. This means that only two of the 2ⁿ rows of the U matrix, namely those with indices 000... and 100..., are relevant (and likewise for the V matrix). The elimination of all the other rows greatly simpli es the problem of optimizing these matrices. W e shall thus write, for brevity,

$$U_{00,mm} \circ_{m} \infty ! U_{mm} \circ_{m} \infty; \qquad (20)$$

where = 0;1. Then, on the left hand side of Eq. (19), we e ectively have two unknown row vectors, U_0 and U_1 , each one with 2^n components (labelled by Latin indices m m⁰m⁰). These vectors have unit norm and are mutually orthogonal. Likewise, Bob has two vectors, V_0 and V_1 . The problem is to optimize these four vectors so as to make the expectation value of the Bell operator [17],

$$C := AB + AB^{0} + A^{0}B A^{0}B^{0}; \qquad (21)$$

as large as possible.

The optim ization proceeds as follows. The new density matrix, for the pairs of spin $\frac{1}{2}$ particles that were not tested by A lice and B ob for spin up (that is, for the st pair in each set of n pairs), is

(new); =

 $N U_{,m m} \circ_{m} \circ V_{,nn} \circ_{n} \circ \circ m_{n}; st m} \circ_{n} \circ_{s} \circ_{t} \circ m \circ_{n} \circ_{s} \circ_{t} \circ u = m \circ_{s} \circ_{s} \circ_{s} \circ V_{,tt} \circ_{t} \circ_{t} \circ (22)$

where N is a norm alization constant, needed to obtain unit trace (N 1 is the probability that all the \spin up" tests were successful). We then have [13], for xed _{new} and all possible choices of C,

$$\max[Tr(C_{new})] = 2^{p} M;$$
 (23)

where M is the sum of the two largest eigenvalues of the real symmetric matrix $T^{y}T$, dened by

$$T_{pq} := Tr[(p_{q})_{new}]: \qquad (24)$$

(In the last equation, $_p$ and $_q$ are the Pauli spin matrices.) Our problem is to nd the vectors U and V that maxim ize M .

At this point, some simplifying assumptions are helpful. Since all matrix elements $m_{n,st}$ are real, we can restrict the search to vectors U and V that have only real components. Furthermore, the situations seen by A lice and B ob are completely symmetric, except for the opposite signs in the standard expression for the singlet state:

These signs can be made to become the same by redening the basis, for example by representing the $\ wn''$ state of B ob's particle by the symbol ${}^{0}_{1}$, without changing the basis used for A lice's particle. This unilateral change of basis is equivalent a substitution

$$V_{\mu n} v_{n} \omega !$$
 (1) $^{+n+n^{0}+n^{\omega}} V_{\mu n} v_{n} \omega;$ (26)

on Bob's side. The m inus signs in Eq. (9) also disappear, and there is complete symmetry for the two observers. It is then plausible that, with the new basis, the optimal U and V are the same. Therefore, when we return to the original basis and notations, they satisfy

$$V_{\mu n n^{0} n^{\infty}} = (1)^{+n+n^{0}+n^{\infty}} U_{\mu n^{0} n^{\infty}}$$
 (27)

W e shall henceforth restrict our search to pairs of vectors that satisfy this relation.

A fter all the above simplications, the problem that has to be solved is the following: nd two mutually orthogonal unit vectors, U_0 and U_1 , each one with 2^n real components, that maximize the value of M (U) dened by Eqs. (23) and (24). This is a standard optimization problem which can be solved numerically. Since the function M (U) is bounded, it has at least one maximum. It may, however, have more than one: there may be several distinct local maxima with diment values. A numerical search leads to one of these maxima, but not necessarily to the largest one. The outcome may depend on the initial point of the search. It is therefore in perative to start from numerous random ly chosen points in order to ascertain, with reasonable con dence, that the largest maximum has indeed been found.

4. NUMERICAL RESULTS

In all the cases that were examined, M (U) turned out to have a localm aximum for the following simple choice:

$$U_{0;00:...} = U_{1;11:...} = 1;$$
 (28)

and all the other components of U_0 and U_1 vanish. Recall that the \vectors" U_0 and U_1 actually are two rows, $U_{000....}$ and $U_{100....}$, of a unitary matrix of order 2^n (the other rows are irrelevant because of the elimination of all the experiments in which a particle failed the spin-up test). In the case n = 2, one of the unitary matrices having the property (28) is a simple permutation matrix that can be implemented by a \controlled-not" quantum gate [18]. The corresponding Boolean operation is known as xor (exclusive or). For larger values of n, matrices that satisfy Eq. (28) will also be called xor-transform ations.

It was found, by num erical calculations, that xor transform ations always are the optim al ones for n = 2. They are also optim al for n = 3 when the singlet fraction x is less than 0.57, and for n = 4 when x < 0.52. For larger values of x, m ore complicated form s of U_0 and U_1 give better results. The existence of two dimensions of maxim a may be seen in Fig. 1: there are discontinuities in the slopes of the graphs for n = 3 and 4, that occur at the values of x where the largest value of hC i jumps from one local maximum to another one.

Forn = 5, a complete determ ination of U_0 and U_1 requires the optim ization of 64 parameters subject to 3 constraints, more than my workstation could handle. I therefore considered only xor-transform ations, which are likely to be optimal for x < 0.5. In particular, for x = 0.5 (the value that was used in W emer's original work [4]), the result is hC i = 2.0087, and the CHSH inequality is violated. This violation occurs in spite of the existence of an explicit LHV model that gives correct results if the W emer pairs are tested one by one.

These results prompt a new question: can we get stronger inseparability criteria by considering , or higher tensor products? It is easily seen that no further progress can be achieved in this way. If is separable as in Eq. (2), so is . Moreover, the partly transposed matrix corresponding to simply is , so that if no eigenvalue of is negative, then too has no negative eigenvalue.

ACKNOW LEDGMENT

This work was supported by the G erard Swope Fund, and the Fund for Encouragement of Research.

References

- [1] J.S.Bell, Physics 1, 195 (1964).
- [2] J.F.Clauser, M.A.Home, A.Shimony, and R.A.Holt, Phys. Rev. Lett. 23, 880 (1969).
- [3] A. Peres, Quantum Theory: Concepts and Methods (Kluwer, Dordrecht, 1993) Chapters 5 and 6.
- [4] R.F.W emer, Phys. Rev. A 40, 4277 (1989).
- [5] S. Popescu, Phys. Rev. Lett. 72, 797 (1994); 74, 2619 (1995).
- [6] N.D.Mermin, in Quantum Mechanics without Observer, edited by R.K.Clifton (Kluwer, Dordrecht, 1996) pp. 57{71.
- [7] N.Gisin, Phys. Lett. A 210, 151 (1996).
- [8] R. Horodecki, P. Horodecki, and M. Horodecki, Phys. Lett. A 210, 377 (1996).
- [9] A. Peres, Phys. Rev. Lett. 77, 1413 (1996).
- [10] J. B lank and P. Exner, Acta Univ. Carolinae, M ath. Phys. 18, 3 (1977).
- [11] C.H.Bennett, G.Brassard, S.Popescu, B.Schum acher, J.Smolin, and W.K.W ootters, Phys. Rev. Lett. 76, 722 (1996).
- [12] M. Horodecki, P. Horodecki, and R. Horodecki, \Separability of mixed quantum states: necessary and su cient conditions" (eprint archive: quant-ph/9605038).
- [13] R.Horodecki, P.Horodecki, and M.Horodecki, Phys. Lett. A 200, 340 (1995).

[14] A.Peres, Phys. Rev. A 54, ... (1996).

- [15] A. Peres, in Quantum Potentiality, Entanglement, and Passionat-a-D istance: Essays for Abner Shimony, ed. by R. S. Cohen, M. Home, and J. Stachel (K luwer, Dordrecht, 1996).
- [16] E. Santos, Phys. Rev. Lett. 66, 1388 (1991).
- [17] S.L.Braunstein, A.Mann, and M.Revzen, Phys. Rev. Lett. 68, 3259 (1992).
- [18] A. Barenco, D. Deutsch, A. Ekert, and R. Jozsa, Phys. Rev. Lett. 74, 4083 (1995).

Caption of gure

FIG.1. Maximal expectation value of the Bell operator, versus the singlet fraction in the W erner state, for collective tests performed on several W erner pairs (from bottom to top of the gure, 1, 2, 3, and 4 pairs, respectively). The CHSH inequality is violated when hC i > 2.