A GravitationalExplanation forQuantum Mechanics

M ark J Hadley Dept. of Physics, Univ. of W arwick COVENTRY CV4 7AL, UK

14 September 1996

A bstract

It is shown that certain structures in classical G eneral R elativity can give rise to non-classical logic, norm ally associated with Q uantum M echanics. A 4-geon m odel of an elementary particle is proposed which is asymptotically at, particle-like and has a non-trivial causal structure. The usual C auchy data are no longer su cient to determ ine a unique evolution. The measurement apparatus itself can in pose non-redundant boundary conditions. M easurements of such an object would fail to satisfy the distributive law of classical physics. Thism odel reconciles G eneral R elativity and Q uantum M echanics without the need for Q uantum G ravity. The equations of Q uantum M echanics are unm odi ed but it is not universal; classical particles and waves could exist and there is no graviton.

1 Comment

This submission reproduces the talk I gave at the 5th UK C onference on C onceptual and Philosophical problems in Physics held in O xford on 10th -14th Septem ber 1996. The content follows the talk very closely but is hopefully m ore coherent - what I m eant to say replaces what I did say. In a similar vein the replies to questions are what I should have said rather than what I actually said; in both cases it is clarity rather than the facts or the arguments which has changed (exceptions to this rule are given as footnotes). Full references are also included.

2 Introduction

I am going to give a gravitational explanation of Q uantum M echanics. By gravitation I m ean E instein's theory of G eneral R elativity – the unm odi ed classical theory. By Q uantum M echanics I m ean the Q uantum M echanics that we all know and love. As far as I am aware nobody has given an explanation for the origin of Q uantum M echanics before, and certainly not in terms of an established classical theory. W hat is m ore I w ill do this in 20 m inutes!!

em ail: m .jhadley@ warw ick.ac.uk

3 The Route from General Relativity to Quantum Mechanics

This diagram shows the route from GeneralRelativity to Schrodinger's equation etc.Quantum logic has a crucial place in the path.




Figure 1: The route from GenialRelativity to Schrodinger's equation via quantum logic.

It is well known that Schrodinger's equation, the D irac equation, P lanck's constant the uncertainty relations etc., etc. can be derived from the H ilbert space structure of Q uantum M echanics, the sym m etries of space and time, and the internal sym m etries of the object. A good reference to the non-relativistic case is given by Ballentine [1], while W einberg gives a useful treatment of the relativistic case [2].

W hat is less well known is that the H ilbert space structure of Q uantum M echanics is a natural representation of quantum logic. In fact it looks increasingly as if T he fam iliar H ilbert space structure is unique as a vectorial representation of quantum logic². Q uantum logic is introduced in the books by Jauch [3] and B eltram etti and C assinelli[4], the latter also describes how the H ilbert space structure is constructed from the logic.

For this talk I will show how quantum logic can arise from the propositions (statem ents) about certain structures in General Relativity. The rest is then already done for me.

4 Quantum Logic

Quantum logic is a non-distributive or non-Boolean logic, which means the failure of the familiar distributive law:

$$a^{(b_c)} \in (a^{(b_c)}) = (a^{(c)})$$
 (1)

^z at least for dim ensions greater than 2

where ^ is the AND operation and _ is the OR operation. a, b and c are the propositions or statements about the system or state. For this talk I will use a special case of equation 1 - taking c to be NOT b, denoted : b, and introducing the trivial operator I, which is TRUE for any state. We then have:

$$a = a^{I} = a^{(b_{:})} \in (a^{b})_{(a^{:})}$$
 (2)

)
$$a \in (a^b)_{(a^:b)}$$
 (3)

In fact quantum logic requires the distributive law to be replaced by a weaker orthom odular condition and for a complete orthom odular orthocom plem ented atom ic lattice with the covering property needs to be constructed. This can be done. It is the subject of a paper submitted to Foundations of Physics and of m y PhD thesis. For this talk I will only show the failure of the distributive law in the form of equation 3, because this marks the departure from a classical system and is by itself a remarkable achievement.

5 General Relativity

For this work the signi cant features of General Relativity are:

The equation, G = 8 T, which relates the curvature of spacetime, of which G is a measure, to the energy momentum tensor T.

It is a non-linear equation for the metric, containing st and second derivatives and both linear and quadratic terms in the metric.

The equations describe distorted, curved spacetime.

The equations are local, they do not prescribe the topology, although they may set constraints on the topology.

The theory allows closed timelike curves, CTCs, (just a respectable way to say time travel). This is one of the great mysteries of General Relativity - if CTCs are possible then how can we make them and use them, and if not, then what forbids them. The mathematical structure of General Relativity allows CTCs and exact solutions are known with CTCs.

6 CTCs

CTCs are crucial for the results which follow, because when interactions are allowed in spacetimes with CTCs the norm alboundary conditions are no longer adequate to uniquely determ ine the evolution.

C onsider a billiard ball in a plane, given an initial position and velocity then the subsequent trajectory is determined, see the dashed line in gure 2; even if there are walls, or hills, or in this example a worm hole.

If the worm hole is replaced with a time machine, so that a particle which enters one mouth exits at a corresponding point from the other mouth, but at an earlier time. The original trajectory is still a possible consistent solution, but now alternatives exist. For example the ball could be hit into the mouth of the worm hole, reappear from the other mouth at an earlier time in such a direction that it causes the original collision (see the solid lines in gure 2). It




Figure 2: The ball travelling from the left m ay be hit by itself into one m outh of the worm hole, to emerge at an earlier time to cause the impact.

must be stressed that these are both consistent evolutions of the system even though the initial data would norm ally (in the absence of CTCs) give a unique trajectory.

The multiplicity of possible solutions is not con ned to this example. It is considered to be a generic feature when self-interacting objects or elds are in a spacetime with CTCs (see for example papers by Friedman et al [5] and Thoma [6]).

7 4-G eon

The strange features of CTCs are exploited in a model of an elementary particle which I call a 4-geon. The idea that an elementary particle is a solution of the eld equations (of General Relativity or any uniedeld theory) dates from the earliest days of General Relativity. Einstein attempted to nd such solutions in all his theories. In the 60's M isner and W heeler[7] continued with the work and used the term geon to describe a topologically non-trivial spacetime estructure held together by its own gravitational attraction. However most of the earlier work used a topologically non-trivial three-manifold evolving with time, and assumed that a global time coordinate existed. By contrast a 4-geon has a non-trivial causal structure. A 4-geon is assumed to have the follow ing properties:

It is a solution of the eld equations of General Relativity.

It has a non-trivial causal structure.

Interactions are taking place around CTCs.

The metric is asymptotically at.

Particle-like: the region of non-trivial topology will be found in one and only one place - otherw ise it would not be recognisable as a particle.

W ith this model of an elementary particle the norm alboundary conditions can no longer be expected to be adequate to determ ine a unique evolution.

8 Boundary Conditions and M easurem ents

The idea that the state preparation sets boundary conditions is obvious. In our real or imagined experiments we look for outcomes consistent with the preparation conditions; this may comprise a source, collimators, shutters liters etc.:




Figure 3: The boundary conditions in posed by state-preparation

W ith the 4-geon model of a particle the state preparation conditions will no longer be adequate to uniquely determ ine the subsequent evolution. The m easurem ent apparatus itself can set further boundary conditions which are not redundant. An x-spin m easurem ent is an example:




Figure 4: The boundary conditions in posed by state-preparation and an x-spin m easurem ent

Note that the m easurem ent apparatus is physically very sim ilar to the state preparation. W ithin the structure of the 4-geon there can be a causal link between the m easurem ent apparatus, state preparation and the evolution, which gives a physical explanation for m easurem ent-dependent e ects.

A lternatively we could measure the y-spin with a very similar apparatus.

However, as is well known, a y-oriented Stern-Gerlach Iter and an x-oriented one are physically incompatible. They set con icting boundary conditions.




Figure 5: The boundary conditions in posed by state-preparation and a y-spin measurem ent

9 Sets of M an ifolds and P ropositions

To see the e ect of these boundary conditions we will consider the possible sets of 4-geon manifolds. Let M denote the set of 4-manifolds consistent with the state preparation conditions. W hile X denotes those manifolds consistent with both the state-preparation and an x-spin measurement. X is partitioned into the two disjoint subsets X⁺ and X .




Figure 6: Sets of 4-m anifolds consistent with both state preparation and the boundary conditions im posed by di erent m easurem ent conditions.

This simple diagram is immediately non-classical, because classically the measurement must simply partition those solutions M consistent with the state preparation; it cannot de ne a proper subset of M .

The y-m easurement de nes a dierent subset of M , denoted $Y\,,\,w\,hich$ is disjoint from X .

The propositions are statements about the state preparation; they are not in one to one correspondence with the measurements because some measurements give the same information about the state (they are indistinguishable by any state preparation). For this system, X corresponds to there is a manifold in M consistent with an x-spin measurement and Y to there is a manifold in M consistent with a y-spin measurement these are both the trivial proposition which is always true and which we denote by I.

By contrast $X^+ \setminus Y^+$ corresponds to there is a manifold in M consistent with a positive x-spin m easurement and also with a positive y-spin m easurement, but X^+ and Y^+ are disjoint and so the intersection corresponds to the trivial proposition which is always false $X^+ \setminus Y^+ = ;$. So we have:

$$X \in (X^+ \setminus Y^+) [(X^+ \setminus Y)]$$
(4)

which is the failure of the distributive law for the propositions.

10 Summary

The conjectured 4-geon description of particles is speculative. I cannot produce a solution of the eld equations with the required properties. I have not tried to. The advice I have received is not to try and nd a solution because it is so di cult so solve E instein's equations, especially if solutions are highly nonlinear, lacking in symmetry and topologically non-trivial.

However, In other respects this theory is extremely conservative – it keeps General Relativity in its unmodied form and it retains 3+1 dimensions for space and time.

The unifying nature of the theory justi es the speculation. Field and particle descriptions of N ature are uni ed as E instein had always hoped and expected. For the st time the origin of Q uantum M echanics is explained in terms of existing theories. In doing so, G eneral Relativity and Q uantum M echanics are reconciled, not with a quantum theory of gravitation as was expected, but with a gravitational explanation for Q uantum M echanics. There is no simpler or m ore conservative theory which reconciles Q uantum M echanics and G eneral Relativity.

11 Predictions

D espite giving standard Q uantum M echanics with the same equations and structure the theory does m ake some new predictions:

There is no quantum theory of gravity.

C lassical objects are possible. The peculiar 4-geon structures give rise to quantum e ects; if these are absent then classical determ inistic evolution would occur.

There is no graviton. This follows from either of the statements above. G ravitational waves are topologically simple solutions of E instein's equations w ithout CTCs. Therefore they cannot exhibit quantum phenomena such as wave particle duality. G ravitational waves are not quantised.

12 Questions

The following questions were asked in open discussions or afterwards. They were most helpful to m e and I thank all those who joined in. A pologies for not giving nam es and for any errors, but I did not make notes at the time.

 ${\tt Q}$. You have shown the failure of the distributive law, but for quantum logic you must show much more – orthom odularity, atom icity etc. C an you show this too?

A. Yes, the failure of the distributive law is the most remarkable feature because it marks the divergence of classical and non-classical systems. Orthom odularity can be shown [8], in fact it follows easily since this construction relies upon the measurement apparatus and so the arguments of Mackey (see [4][page 147]) apply. A tom icity is a mathematical idealisation which cannot be derived, but this sort of idealisation is common to all of mathematical physics; eg. the use of real numbers to represent momentum in classical physics.

Q. How can spin-half arise in a gravitational theory?

A. There is an enorm ous richness in the choice of topology. Certain manifolds can be shown to have the transform ation properties of a spinor provided a xed asymptotically at background metric is assumed. See the fascinating paper by Friedman and Sorkin [9] or the discussion in my thesis).

 ${\tt Q}$. How does the superposition of the wavefunction arise in this model?

A. The wavefunction just gives inform ation about the probability m easurement outcomes. If the logic were Boolean a real number between 0 and 1 would su ce and there would be only trivial superpositions. However to represent probabilities for a non-Boolean logic, complex-valued wavefunctions are required.

Q. How can you get Quantum Mechanics which is formulated on a at spacetime when you are considering manifolds with a nontrivial topology?

A. The manifolds are asymptotically at. Quantum Mechanics can be regarded as a way of mapping information about these knots of spacetime onto the at spacetime which we are familiar with. It is in the asymptotically at region that we set boundary conditions etc.

Q . Do solutions of E instein's equations exist with CTCs which can be traversed in a nite time?

A.Yes.

Q. There are alternative ways of assigning probabilities to an orthom odular lattice which cannot be represented by a Hilbert space. Consider for example the model by M ielnik [10] which can be found in [4] [page 205].

A. I am not aware of that example. However the very existence of nonclassical logic in systems described by a classical theory is by itself most remarkable, to get quantum logic as well is amazing. The familiar Hilbert space structure is certainly compatible with this logic even if it is not unique.

 ${\tt Q}$. A re you aware of other work in which an orthom odular lattice is constructed geometrically from subsets of at M inkowski space?

A . I have seen geom etric constructions of orthom odular lattices eg. W atanabe [11][page 303]. I think that such m odels rely on an innovative de nition of com plem entation, they are interesting but not particularly rem arkable. M y work shows that the orthocom plem ented lattice arises with the de nitions of com plem entation associated with real experim ents. In this respect the construction is unique.

Q.W hat is the energy tensor responsible for the spacetime knots?

A. I have deliberately not m ade assumptions about the energy -m om entum tensor. The most appealing case would be for it be zero i.e. a vacuum solution. Spacetime can be knotted without any source. Indeed the worm holes (not traversable ones) can be solutions of the source-free eld equations.

13 A cknow ledgem ents

I would like to thank Harvey Brown, Mauricio Suarez and Katherine Brading from the Department of Philosophy, Oxford University, for organising the conference. This work has been supported by the University of Warwick.

References

- [1] Leslie E Ballentine. Quantum Mechanics. Prentice Hall, 1989.
- [2] Steven W einberg. The quantum theory of elds. Cambridge University Press, 1995.
- [3] Joseph M Jauch. Foundations of Quantum Mechanics. Addison-Wesley Publishing Company, 1968.
- [4] Enrico G Beltram etti and Gianna Cassinelli. The Logic of Quantum Mechanics, volum e 15 of Encyclopedia of M athematics and its Applications. Addison-W esley Publishing Company, 1981.
- [5] John Friedman, Michael S Morris, Igor D Novikov, and Ulvi Yurtsever. Cauchy problem in spacetimes with closed timelike curves. Physical Review D, 42 (6):1915{1930,1990.
- [6] K ip S T home. C losed tim elike curves. In 13th International C onference on G eneral Relativity and G ravitation 1992. Institute of P hysics P ublishing.
- [7] C W M isner and J A W heeler. C lassical physics as geometry. Annals of Physics, 2:525, 1957.
- [8] Mark J Hadley. The logic of quantum mechanics derived from classical general relativity. Foundations of Physics, Submitted June 1996.
- [9] John L Friedman and Rafael D Sorkin. Half-integral spin from quantum gravity. General Relativity and Gravitation, 14:615{620, 1982.
- [10] Bogden M ielnik. G eom etry of quantum states. C om m unications in M athem atical Physics, 9:55{80, 1968.
- [11] SatosiW atanabe. Knowing and Guessing: A Quantitative Study of Inference and Information. W iley and Sons, 1969.