

Experimental Status of Quaternionic Quantum Mechanics

S. P. Brumby and G. C. Joshi.

School of Physics, University of Melbourne, Research Centre for High Energy Physics,
Parkville, Victoria 3052, Australia.

Abstract

A analysis of the logical foundations of quantum mechanics indicates the possibility of constructing a theory using quaternionic Hilbert spaces. Whether this mathematical structure reflects reality is a matter for experiment to decide. We review the only direct search for quaternionic quantum mechanics yet carried out and outline a recent proposal by the present authors to look for quaternionic effects in correlated multi-particle systems. We set out how such experiments might distinguish between the several quaternionic models proposed in the literature.

Report Number: UM-TP-94/120 ; RCHEP-94/34

PACS: 3.65Bz

arXiv:quant-ph/9610009v1 8 Oct 1996

Typeset using REVTeX

I. QUATERNIONIC THEORIES.

The established formalism of quantum mechanics identifies physical observables with Hermitian operators on a Hilbert space of states. The existence of quantum states consisting of superpositions of pure states is permitted by the Hilbert space structure. Operators corresponding to different observables need not commute, but the outcome of a measurement on a system will be an eigenvalue of the corresponding Hermitian operator and hence real. It is this last fact which led Birkhoff and von Neumann [1] to conclude that it is possible to consider the states to form a vector space over the real, complex or quaternionic algebras. Birkhoff and von Neumann state that these theories cannot be differentiated by known formalistic criteria.

Stueckelberg [2] has shown that the description of polarisation states in a real number formulation of quantum mechanics (RQM) requires the introduction of a superselection operator with all of the algebraic properties of the imaginary unit i . Hence, known phenomena have demanded the enlargement of RQM to be essentially equivalent to complex quantum mechanics (CQM).

The remaining possibility involves considering Hilbert spaces over the division algebra of real quaternions H . This is generated over the real numbers R by a basis of abstract elements $\{i_0 = 1; i_1; i_2; i_3\}$ with the multiplication rule, $i_1^2 = i_2^2 = i_3^2 = -1$ and $i_1 i_2 = i_3$, $i_2 i_1 = -i_3$,

$$i_1 i_2 = i_3, \quad i_2 i_1 = -i_3; \quad i_1^2 = i_2^2 = i_3^2 = -1 \quad \text{and} \quad i_1 i_3 = -i_2, \quad i_3 i_1 = i_2 \quad (1)$$

That is, any quaternion a can be written in the form

$$a = a_0 + a_1 i_1 + a_2 i_2 + a_3 i_3; \quad a_i \in R \quad (2)$$

The algebra has the anti-involution $1 = 1, i_r = -i_r, (ab) = b a$, and a norm $\|a\|^2 = a a^* = a^* a \in R$.

If we introduce an additional complex unit $i_4 = \sqrt{-1}$, defined to commute with each of i_0, i_1, i_2 and i_3 , then the algebra ceases to be a division algebra. This complexification of H is called the biquaternionic algebra, and has been used to treat special relativity and electromagnetism [3]. The Pauli matrices are complex 2×2 matrix representations of the Hermitian biquaternionic units $i_4 i_r$. The Lie algebra of $SU(2)$ is H with the commutator product $[a; b] = ab - ba$. Hence, theories of spin-1/2 fermions, Salam {Weinberg unified electro-weak theory, and Yang-Mills isospin can be written in a quaternionic form [4]. For a review of these and other uses of the quaternion algebra in standard quantum mechanics and quantum field theory, see [5] and references therein.

Quaternionic quantum mechanics (QQM) differs from these other (mainstream) applications of the quaternions, because it considers the Hilbert space of quantum states to be a vector space over H . Hence, eigenvalues of operators on the quaternionic Hilbert space H_H need not commute, though we retain the identification of physical observables with Hermitian operators which, as is well known, necessarily have real eigenvalues only. The relation between kets of H_H and quaternionic wavefunctions follows the usual pattern (where the $\psi \in R$).

$$\langle x | j \rangle = \psi_0 + \psi_1 i_1 + \psi_2 i_2 + \psi_3 i_3 \quad (3)$$

Alternatively, we can express ψ as an ordered pair of i_1 -complex numbers, in its so-called symplectic representation,

$$(\psi) = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} + i_2 \begin{pmatrix} \psi_3 \\ \psi_4 \end{pmatrix}; \quad (4)$$

where $\psi_1 = \psi_0 + i_1 \psi_1$ and $\psi_2 = \psi_2 + i_1 \psi_3$.

If we define $\tilde{j} = \left(\prod_{r=1}^3 \frac{2}{r} \right)^{1/2} \in \mathbb{R}$, then there exists a pure imaginary quaternion of norm unity \tilde{j} , such that

$$(\psi) = \psi_0 (\psi) + \tilde{j} (\psi) \tilde{j} (\psi); \quad (5)$$

From this we see that any two wavefunctions of QQM do not commute unless their imaginary parts are parallel. This property has confounded the construction of a completely satisfactory tensor product, i.e., one which would be quaternion linear in each factor, allow the definition of a tensor product of operators on each factor, and admitting a positive scalar product for the purpose of second quantisation of the theory (see [6,7] for comprehensive discussions).

The obvious and prodigious success of CQM in describing physical systems demands that QQM must reproduce its results wherever CQM has been successful. This led Finkelstein et al. [8] to propose a geometric generalisation of CQM, producing what is in effect an "almost complex quantum mechanics" [9]. Thus QQM is supposed to be complex in any neighbourhood of an observation, modelled by replacing the imaginary unit of CQM with a field of pure imaginary unit quaternions i_x over spacetime, and the language of fibre-bundles is used throughout their work. Over larger distances, topological properties of the spacetime manifold are permitted to manifest themselves in quantum mechanical expectation values. Finkelstein et al. identify an intermediary limit between full geometric QQM and standard CQM, which they call the "electromagnetic limit". In this limit the Q{covariant derivative of the imaginary unit vanishes and the Q{curvature term in the Lagrangian looks like that of EM. The result of this situation is no extra quaternionic degrees of freedom in the wavefunction, and the theory is equivalent to a (complex) Abelian gauge field theory.

Adler [10] and Horowitz [11] have investigated the case where it is possible, by use of a quaternionic gauge transformation, to choose all of the energy eigenkets of a system to be in the same complex subset of H_H . Then QQM effects are dependent upon the existence of new, hyper-complex components of the fundamental forces.

The search for these hypothetical extra components motivated the experimental test of QQM undertaken by Kaiser, George and Werner [12] in 1984.

II. INTERFEROMETRY TEST OF QQM.

The experiment of Kaiser et al. is based on a proposal of Peres [13], who suggested that if a pair of potential barriers were to possess additional hyper-complex components, then a particle (in Peres's proposal, a neutron) traversing the pair will experience a shift in the phase of its wavefunction which will depend upon the order in which the barriers are traversed.

Note that even though relativistic formulations of QQM have been proposed [8,10], this experiment limits itself to the special case of classical, external fields acting on nonrelativistic

matter-waves, as this is the situation amenable to investigation by particle interferometry. The problem of how to quantise quaternionic potentials remains open, while quantisation of the field requires the overcoming of obstacles similar to those attending the construction of a quantised gravity.

Peres's original suggestion contained the idea that the one should use potential barriers with a large absorptive cross-section, which in the standard framework of QM are modelled by potentials with large, non-vanishing imaginary components. The idea behind this was that the imaginary number appearing in the potential might depend upon the material, so potentials corresponding to different barriers would not commute, while having a large imaginary component might make it more likely that the hyper-complex contribution to each potential is significant. The total phase shift experienced by a neutron of wavelength λ , traversing a plate of thickness D and refractive index n is

$$\Delta\phi = \frac{2\pi}{\lambda} (n - 1)D ; \quad (6)$$

where for the thermal neutrons used in the experiment, n is related to the coherent nuclear scattering length b and the atom density N of the barrier material, by

$$n = 1 - \frac{2\pi N b}{\lambda^2} ; \quad (7)$$

Kaiser et al. decided that to achieve very high sensitivity to differences in the total phase shift caused by traversal of the apparatus in different directions, the thickness of the slabs should be such that

$$\frac{2\pi N b D}{\lambda^2} = 10000 ; \quad (8)$$

This ruled out the possibility of using materials with very high absorption cross-sections. Instead, they chose aluminium and titanium for their differing chemical and nuclear properties, aluminium having a positive real scattering amplitude while titanium has a negative real scattering amplitude, though both have small absorption cross sections. The experiment, at the 10-MW University of Missouri Research Reactor, used thermal neutrons of wavelength $\lambda = 1.268 \text{ \AA}$ in a Bonse-Hart three crystal Laue-Laue-Laue interferometer (see [14] for a review of the techniques of neutron interferometry).

Kaiser's et al. result was that in changing the order of traversal of the Al and Ti barriers, no phase difference was observed to 1 part in 30000.

A theoretical analysis of the two potential barrier experiment was later undertaken by Davies and McKellar [15] using the quaternionic one dimensional Schrodinger equation of Adler [10]. This uses the symplectic decomposition of the wavefunction to rewrite the quaternionic problem into one of a pair of complex fields coupled through the hyper-complex components of the system Hamiltonian. In time independent form, this is the pair of equations

$$\frac{d^2 \psi_1}{dx^2} + V_1 \psi_1 - V_2 \psi_2 = E \psi_1 \quad (9)$$

$$\frac{d^2 \psi_2}{dx^2} - V_1 \psi_2 + V_2 \psi_1 = E \psi_2 \quad (10)$$

Davies and McKellar attempted to treat the problem exactly but found the resulting barrier transmission and reflection coefficients to be too unwieldy to analyse. Instead, they turned to numerical methods and concluded that the transmission coefficient for two quaternionic, square potential barriers of different heights will change by a phase when the system is traversed in the opposite sense. This then forces us to ask how this conclusion can be consistent with the null result (to one part in 30 000) of Kaiser et al.

Now the possibility remains that QQM modifies only some of the fundamental forces. The experiment of Kaiser et al. would appear to severely constrain any quaternionic component of the strong nuclear force, though this experiment does not rule out the possibility that there might be a quaternionic component to the underlying chromodynamics. Though QQM was at one time considered as a possible framework for QCD, Adler has proposed that QQM might provide a natural framework of preonic physics [10].

Alternatively, it might be that the experiment of Kaiser et al. is too simplistic, as it only involves a single fundamental force. One can conceive of a situation in which we associate a different complex algebra with each of the fundamental forces, so that within each class of interactions complex quantum theory succeeds to describe the experimental phenomena.

To rule out this possibility, Klein [16] has suggested that we should repeat the experiment with combinations of different fundamental forces¹. The simplest extension is to introduce a sequence of potential barriers and magnetic fields which interact with the neutron intrinsic spin. Additionally, there are known phenomena associated with the terrestrial gravitational field which are subject to investigation by neutron interferometric methods. The fact that such effects have been successfully taken into account in previous interferometric experiments means that we can already severely constrain the contribution of a quaternionic addition to the gravitational force, or else deduce that the gravitational force reached in the classical limit from a quaternionic theory of quantised gravity, involves observables and states constrained to lie in a complex subset of H coincident with that selected by the strong force.

An experiment of the type described in [16], if it were to produce a null result, would leave only the weak interaction as a possible force with a quaternionic component. We should then search for new effects in neutrino physics (already suggested by Peres [13]). The extreme weakness of these interactions, however, renders very unlikely the possibility of carrying out interferometric experiments of the type previously considered.

A positive outcome from Klein's experiment would give us some knowledge of the particular form of QQM preferred by Nature. If there was observed a pure phase shift introduced by swapping the order of potential barriers, this would imply that we live in the $Q\{$ at limit of QQM (or that $Q\{$ curvature is negligible at the scale of the experiment, and perhaps also in the vicinity of the Earth). Any more complicated observational result, in particular the production of qualitatively different interferometric patterns, would imply the active presence of a $Q\{$ curvature.

Before moving on to consider experiments on correlated few body systems, we mention that Peres [13] in fact suggested three different types of experiments. Besides the experimen-

¹Since this article first appeared, this experiment has been attempted. We refer the reader to a description of the preliminary results in [17].

tally realised interferometry test described above, he derived a universal relation between the scattering cross-sections of three coherent sources, taken singly and pairwise. Numerical violation of this relation would indicate the failure of the complex description. Additionally, he suggested the investigation of K_s meson regeneration by different media taken singly and pairwise. To the best of our knowledge, neither possibility has been taken up by experimental researchers.

III. EXPERIMENTS ON MULTI-PARTICLE CORRELATED SYSTEMS.

Note that in the previous discussion, the presence of quaternionic effects depends upon the existence of hyper-complex components in the interaction potentials of a physical system. Theoretical analysis of this situation requires the adoption of a particular form for the quaternionic dynamics, which in the non-relativistic QFT limit might be characterised by the assumption of QQM essentially reducing to CQM in potential free regions of space. Recently, Burnby, Joshi, and Anderson [18] have questioned the need to demand this integrability of the QFT curvature, instead proposing a different class of experiments.

To this end, consider the variant due to Greenberger et al. [19] of Bohm's gedankenexperiment test of the Einstein-Podolsky-Rosen programme. This envisages the preparation of an entangled four body state

$$|j_{GHZS}\rangle = \frac{1}{\sqrt{2}} (|j_{i_1} j_{i_2} j_{i_3} j_{i_4}\rangle + |j_{i_1} j_{i_2} \bar{j}_{i_3} \bar{j}_{i_4}\rangle); \quad (11)$$

whose constituents are permitted to propagate to space-like separations. We then carry out simultaneous measurements of a component of spin of each particle (using, say, Stern-Gerlach magnets) and consider the product of these values. The quantum mechanical expectation value is thus

$$E_{GHZS}(f_{n_k}g) = \hbar_{GHZS} \prod_{j=1}^4 n_j \cdot \prod_{j=1}^4 j_{GHZS} i; \quad (12)$$

where n_j gives the orientation of the j^{th} Stern-Gerlach apparatus, and σ_j^s is the s^{th} Pauli matrix at the position x_j of that apparatus (and so is a 2×2 , i_{x_j} -complex matrix).

Assuming only that in the nonrelativistic limit the intrinsic spin of a particle does not effect its propagation in free space, Burnby et al. now suggest that this expectation value will be sensitive to the QFT curvature throughout the 2d region whose boundary intersects the four observation points. Hence, geometric QQM predicts experimental results unexplainable within the structure of standard CQM. Our major difficulty lies in the adoption of a tensor product of single particle states which adequately handles the noncommutation of quaternionic wavefunctions and also permits us to return to an equivalent complex quantum mechanical system through some natural limiting process (we use the tensor product of quaternionic Hilbert modules developed by Horwitz and Razon [7]). While this question remains open, investigation of alternative structures has suggested composition of quaternionic Hilbert spaces could be viewed as a lattice theoretic problem [6].

It is potentially significant that we show geometric QQM to agree with CQM in the case of a two body correlated system, essentially due to the dimensionality of the space within which we align our experimental apparatus, and the natural requirement that we use only relative angles when constructing the operator corresponding to the simultaneous observation of components of intrinsic spin. This special "hiding" effect in two body systems is lost in $N = 3$ body systems, and hints that the apparent lack of evidence for QQM may be due to the subtlety of the theory.

Experimental investigations of violations of the Bell inequalities have almost always used photons rather than electrons (see [20,21]). Because photons possess a single quantum of angular momentum, the operator corresponding to a polarisation filter has real components and QQM has no opportunity to manifest itself. This situation is changed by the use of circularly polarised photons, in which case the quantum mechanical prescription for calculating the expectation values necessarily uses complex numbers, giving the quaternionic entangled state a probability distribution sensitive to a changing field (hence, not predicted by CQM). Our prediction of quaternionic terms in multiparticle correlation experiments provides a further motivation to the reasons given by Greenberger et al. to undertake such experiments.

With this type of experiment, we are probing the structure of geometric QQM's bundle of quaternionic fields over spacetime (equivalent to determining the integrability of the almost complex structure). This raises the possibility that topological defects in the spacetime manifold will have a new way of affecting the predictions of quantum mechanics in their vicinity.

Work is in progress on a quaternionic quantum field theory which will allow us to begin to investigate the consequences of the inclusion of such objects [23].

ACKNOWLEDGMENTS

One of the authors, S.P.B., acknowledges the support of an Australian Postgraduate Research Award. G.C.J. was supported by the Australian Research Council and the University of Melbourne.

REFERENCES

- [1] G. Birkhoff and J. von Neumann, *Ann. Math.* 37, 823 (1936).
- [2] E. C. G. Stueckelberg, *Helv. Phys. Acta* 33, 727 (1960).
- [3] C. Lanczos, *Variational Principles of Mechanics* (U. of Toronto Press, Toronto, 1949).
- [4] C. Itzykson and J-B. Zuber, *Quantum Field Theory* (McGraw Hill, New York, 1980).
- [5] F. Gursey, in *Symmetries in Physics (1600-1980) : Proceedings of the 1st International Meeting on the History of Scientific Ideas (Seminari d'Historia de les Ciències, Barcelona, 1983)*, pp. 557-589.
- [6] C. G. Nash and G. C. Joshi, *Int. J. Theor. Phys.* 31, 965 (1992); *J. Math. Phys.* 28, 2883 (1987); 28, 2886 (1987).
- [7] A. Razon and L. P. Horowitz, *Acta Appl. Math.* 24, 141 (1991); 24, 179 (1991); *J. Math. Phys.* 33, 3098 (1992).
- [8] D. Finkelstein, J. M. Jauch, S. Schiminovich, and D. Speiser, *J. Math. Phys.* 3, 207 (1962); 4, 788 (1963).
- [9] M. Nakahara, *Geometry, Topology and Physics* (Institute of Physics, Bristol, 1990), Chap. 8, Sec. 7.
- [10] S. L. Adler, *Phys. Rev. D* 17, 3212 (1978); *Phys. Lett.* 86B, 203 (1979) (ERRATUM – *ibid.* 87B, 406 (1979)); *Phys. Rev. Lett.* 57, 167 (1986); *Phys. Rev. D* 37, 3654 (1988); *Phys. Lett.* B 332 (1994) 358; *Quantum mechanics and quantum fields* (Oxford, New York, 1995), and references therein.
- [11] L. P. Horowitz, *J. Math. Phys.* 34, 3405 (1993); L. P. Horowitz and L. C. Biedenharn, *Ann. Phys.* 157, 432 (1984).
- [12] H. Kaiser, E. A. George and S. A. Werner, *Phys. Rev. A* 29, R2276 (1984).
- [13] A. Peres, *Phys. Rev. Lett.* 42, 683 (1979); quant-ph/9605024.
- [14] A. G. Klein and S. A. Werner, *Rep. Prog. Phys.* 46, 259 (1983).
- [15] A. J. Davies and B. H. J. McKellar, *Phys. Rev. A* 40, 4209 (1989); 46, 3671 (1992); A. J. Davies, *Phys. Rev. D* 41, 2628 (1990).
- [16] A. G. Klein, *Physica B* 151, 44 (1988).
- [17] B. Alman, et al., to be published in *J. Japan Phys. Soc.* (1996).
- [18] S. P. Brumby, G. C. Joshi and R. Anderson, *Phys. Rev. A* 51, 976 (1995).
- [19] D. M. Greenberger, M. Home, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, edited by M. Kafatos (Kluwer Academic, Dordrecht, The Netherlands, 1989), p. 73; D. M. Greenberger, M. A. Home, A. Shimony and A. Zeilinger, *Am. J. Phys.* 58, 1131 (1990).
- [20] J. F. Clauser and A. Shimony, *Rep. Prog. Phys.* 41, 1881 (1978).
- [21] A. Aspect and P. Grangier, in *Quantum Concepts in Space and Time*, edited by R. Penrose and C. J. Isham (Oxford Science Publications, Oxford, 1986).
- [22] P. M. Morse and H. Feshbach, *Methods of Theoretical Physics* (McGraw Hill, New York, 1953), Vol. 1.
- [23] S. P. Brumby and G. C. Joshi, hep-th/9610033; to be published in *Found. Phys.* (1996).