

Transfer formalism for quantum optics problems

V N Gorbachev, A I Zhilba

April 17, 2024

Abstract

Consistent quantum formalism based on the localized basis of the Wannier functions in Heisenberg and Schrodinger pictures to describe propagation of electromagnetic field in a three dimensional media including diffraction is presented. In the Schrodinger picture the Fokker-Planck equation for the Glauber-Sudarshan quasiprobability and corresponding Langevin equations are given. As result the space-time description is obtained by a simple changing variables in the temporal master equation of the field. Using this formalism it is shown that the existence of integrals of motion in the propagation of light in a medium under the condition of nondegenerated parametric and two-photon interactions results in amplification of modes when nonclassical properties of the light are conserved. Quantum propagation of light in a linear medium taking into account the diffraction is considered and its solution is found.

1 Introduction

When considering statistics of the light the usual approaches are based on the temporal evolution of the electromagnetic field. It is natural for the problems when the spatial behavior can be neglected as for the high-Q optical cavity scheme inside which one the temporal feature is important. By contrast the spatial behavior plays the key role in propagation of the light through media or distributed system that can be not considered as a point one. In these systems new states of the light named the spatially squeezed states arise [1]. The quantum formalism for distributed systems has been developed in a number of works. One-dimensional approaches were presented in [2]. Theory for continuous variables in the Heisenberg picture was given in [3], where the continual coherent states and squeezed states of the field have been introduced and theory of the light detection in free space has been presented. With the use of these approaches evolution of the light statistics has been examined for the three-photon [4] and four-photon [5] parametric interaction in transparent media. The multiphoton processes was discussed in [6].

In this paper we present a quantum formalism which enables to consider propagating of the light in a three-dimensional medium with diffraction to be taken into account. It is based on a localized basis of the Wannier functions, and allows the conversion of the temporal description to space-time description by a simple change of variables. Indeed it is the way used in solid state physics for transition from collective to local variables [7]. The main feature of the local description in the Wannier basis is that the problem turns out to be multiparticle when the local field oscillators interact among themselves even in free space. From the physical point of view this is a transfer of excitation or the light propagation process. In other hand the interaction of local oscillators produces the coupled equations which as we show have the form of the transfer equations. Our approach is formulated in both the Heisenberg and Schrodinger pictures where the Fokker-Planck equations for the field quasiprobability are derived together with the corresponding Langevin equations.

To describe interaction of light with atoms the adiabatic illumination of fast atomic variables is often used to obtain a closed equation of field or master equation [8], [9] that is the start point to analyze statistics of the light. The simple recipe of how to get the local description allows to obtain the transfer equation immediately from the temporal master equation missing all steps of the derivation procedure.

The potential of the presented formalism is illustrated by two problems. In the first one the statistics of the light is considered for the propagation in a one-dimensional medium with two-photon and parametric interaction. Here within the framework of local description an interesting peculiarity due to the integrals of motion arises. The existence of the integrals makes it possible to establish the main features of the statistics transformation immediately without going into the solution of the dynamic problem [10]. It has been shown by this way that enhancement of light and the conservation of nonclassical properties can be possible. In the second problem the propagation of the light in a linear three-dimensional medium with diffraction is considered and the solutions have been found.

2 Wannier basis. Local operators

In one-dimensional normalization space L plane waves $\psi_k(x) = \frac{1}{\sqrt{L}} \exp(ikx)$, where $k = 2\pi/L$, are forming a complete orthonormal basis. It is used for a standard representation of the electromagnetic field strength operator, where the operators of photon creation and annihilation $a_k^\dagger; a_k$ are arising with

commutational relations

$$[a_k; a_{k^0}^y] = \delta_{kk^0} \quad (2.1)$$

Operators $a_k^y; a_k$ describe the creation and annihilation of photons of a wave vector k in the whole space L . These operators may be named collective, because they are responsible for the excitation of the whole volume. For the local description of the electromagnetic field instead of plane waves we use Wannier functions known in solid state physics [7], which are packets of plane waves

$$w_m(x, l) = \frac{1}{\sqrt{N}} \sum_{k \in m} \exp(ikl) e_k(x) \quad (2.2)$$

the wave vectors $k \in m$ are lying in a band $m = a \leq k < m + a$. Here each band m , or zone, corresponds to the partition of one-dimensional space L into $N = L/a$ cells, which positions are determined as $l = na$. In k -space the centers of so defined bands are separated by an interval $\Delta k = 2\pi/a$. The terms in (2.2) are forming a geometrical progression, so summation is possible. Then it easily verified that for large N the Wannier functions are localized in a cell with a coordinate l in region $la \leq x < (l+1)a$. The Wannier functions defined in accordance with (2.2) form the complete orthonormal set

$$\sum_{m=1}^X w_m(x, l) w_m(x^0, l) = \delta(x, x^0) \quad (2.3)$$

$$(w_m(x, l); w_{m^0}(x, l)) = \delta_{mm^0} \delta_{ll^0} \quad (2.4)$$

where the scalar product is defined by the integration over the whole space L .

Wannier basis serves as a framework for the introduction of local operators of creation and annihilation $a_{m, l}^y; a_{m, l}$:

$$a_k = \sum_{m=1}^X a_{m, l} w_m(x, l) \quad (2.5)$$

From (2.5) it follows that the operators are coupled by a unitary transformation

$$a_{m, l} = \sum_k C_{m, k}(l) a_k \quad (2.6)$$

$$a_k = \sum_{m=1}^X C_{m, k}(l) a_{m, l} \quad (2.7)$$

where

$$C_{m, k}(l) = \frac{1}{\sqrt{N}} \sum_{k^0 \in m} \exp(ik^0 l) e_{k^0 k} \quad (2.8)$$

For local operators the following commutational relations are valid:

$$[a_{m, l}; a_{m^0, l^0}^y] = \delta_{mm^0} \delta_{ll^0} \quad (2.9)$$

The relations (2.5) allow to interpret them as operators of creation and annihilation of a photon in a point l in the vicinity $la \leq x < (l+1)a$. As a result local field oscillators described by the introduced operators are defined in the space L . From here on we will refer the packets of plane waves forming the local operators as local modes with wave number m and with a width $\Delta x_m = ca$. Strictly speaking these

packets are not modes, because there are no physical reasons for their distinctions, however the term seems to be convenient.

The equation (2.5) is a starting point in the formalism of quantum transfer theory, which may be formulated in different pictures. From the operational point of view the unitary transformations (2.6) and (2.7) following from (2.5) mean that the transition from the nonlocal to local description and vice versa is accomplished by the change of variables.

The specific feature of local description is its many-particle character, where the local oscillators are interacting already in the free space. This interaction describes the excitation transfer or light propagation process, which determine the structure of equations of motion, where chains of the BBGKI type (Bogolyubov, Born, Green, Kirkwood, Ivon) are arising resulting finally in the propagation equations.

Consider the case of free space, for which the field evolution is defined only by a Hamiltonian $H_0 = \sum_k \hbar \omega_k a_k^\dagger a_k$, $\omega_k = ck$. Let the operator a_k in (2.5) be defined in the Heisenberg representation, i.e. satisfying the equation $\partial a_k / \partial t = i\hbar^{-1} [H_0; a_k]$. To find the equation for the local Heisenberg operator

$$a_m(l;t) = \frac{1}{N} \sum_k a_k(t) \exp(ikl) \quad (2.10)$$

let us make the differentiation with respect to time. Assuming the size of the space cell a is small, $a \ll L$, and $a \neq 0$, l may be considered as a continuous space coordinate. In this approximation

$$k \exp(ikl) = i \frac{\partial}{\partial l} \exp(ikl) \quad (2.11)$$

as result it follows that the transfer equation for the local operator is

$$\frac{\partial}{\partial t} a_m(l;t) + c \frac{\partial}{\partial l} a_m(l;t) = 0 \quad (2.12)$$

This equation is valid in a coarse space scale with a characteristic size a providing the fulfillment of (2.11), in this case the replacement $\sum_l \rightarrow \int dl$ is possible.

For a many-particle problem (2.12) may be represented as set of coupled equations due to the interaction of local operators. The Hamiltonian H_0 appears to be nondiagonal with respect to the indices l^0 . The equation of motion takes the form

$$\frac{\partial}{\partial t} a_m(l) = i\hbar^{-1} [H_0; a_m(l)] = i \sum_{l'} a_m(l;l') a_m(l') \quad (2.13)$$

the coupling constant

$$a_m(l;l') = \frac{c}{N} \sum_k \exp(ik(l-l')) \quad (2.14)$$

links the oscillator in the point l with all neighbors. However, due to its δ -shape, the coupling appears to be significant only for two adjacent oscillators. In other words, in the free space the interaction between local oscillators is described in a coarse scale by a derivative with respect to l :

$$\frac{\partial}{\partial l} a_m(l) = \frac{i}{c} \sum_{l'} a_m(l;l') a_m(l') \quad (2.15)$$

the equation (2.15) is obtained by differentiation of (2.10) with respect to l with taking into account (2.11). Consider the commutator of local operators taking for different times. For the free evolution, when the dynamics of operators is determined by the Hamiltonian H_0

$$[a_m(l;t); a_m^y(l;t+\tau)] = \frac{1}{m} \exp(i\omega_m \tau) a_m(l) \quad (2.16)$$

the function (2.17)

$$a_m(l) = \frac{1}{N} \frac{\sin(\frac{\omega_m l}{N})}{\sin(\frac{\omega_m}{N})} \quad (2.17)$$

N^{-1} has a sharp maximum for $l \neq 0$. Its weight is concentrated in the vicinity of the order of $a=c = \frac{1}{m}$, so it may be considered as a delta-function when $\tau = a=c$. Notice that $a(0) = \frac{1}{m}$. The occurrence of the scale temporal delta-function is connected to the use of coarse space scale with a characteristic size a , where the time interval will be $\tau = a=c$.

Let us introduce the interaction picture. Consider the slowly changing part or envelope in the local operator $a_m(l)$:

$$a_m(l;t) = A_m(l;t) \exp(-i\omega_m t + i\omega_m l) \quad (2.18)$$

where $\omega_m = \omega_m$, Then the unitary transformations in (2.6) and (2.7) will take the form

$$A_m(l;t) = \frac{1}{N} \sum_{k,m} a_k(t) \exp(i(\omega_k - \omega_m)t + i(k - m)l) \quad (2.19)$$

$$a_k(t) = \frac{1}{N} \sum_{l} A_m(l;t) \exp(i(\omega_k - \omega_m)t - i(k - m)l) \quad (2.20)$$

the evolution of the operators a_k and $A_m(l;t)$ is defined only by the Hamiltonian of interaction V

$$\frac{\partial}{\partial t} a_k(t) = i h^{-1} [V(t); a_k(t)] \quad (2.21)$$

$$\frac{\partial}{\partial t} + c \frac{\partial}{\partial l} A_m(l;t) = i h^{-1} [V(t); A_m(l;t)] \quad (2.22)$$

3 Three-dimensional case. Quasioptical approximation

Consider a normalized volume L^3 with a number of cells $N = N_1 N_2 N_3, N_1 = N_2 = N_3 = L/a$, where for simplicity the cell was chosen to be of cubic shape. Then in the initial formulas the evident replacements will be: $k \rightarrow k, m \rightarrow m, l \rightarrow l, x \rightarrow r$. For the local operator in the interaction picture the following expression should be written instead of (2.19):

$$A_m(l;t) = \frac{1}{N} \sum_{k,m} a_k(t) \exp(i(\omega_k - \omega_m)t + (k - m)l) \quad (3.23)$$

where the dispersion law has the form $\omega_q = c \sqrt{q^2}$, $q = k - m$. Let us find the equation of motion for local operators. Similarly to the one-dimensional case let us differentiate (3.23) with respect to time. Assuming l to be a continuously changing variable, we will find instead of (2.11)

$$c \frac{\partial}{\partial l} \frac{1}{(q;q) \exp(iq l)} = \frac{\partial}{\partial l} \frac{1}{2q} + \frac{\partial^2}{\partial l^2} + \frac{\partial^2}{\partial l^2} \exp(iq l) \quad (3.24)$$

When summing over k we replace $k_z \approx m_z$. This approximation corresponds to a quasiplane wave of frequency $\omega_m = cm$, propagating along the z -axis. Inasmuch as a coarse spatial scale was introduced, for which vector l may be considered as a continuous function of the coordinates, let us make a replacement $l \rightarrow r(x, y, z)$. As a result the equation of motion takes the form

$$\frac{\partial}{\partial t} + c \frac{\partial}{\partial z} - \frac{c^2}{2\omega_m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) A_m(r) = -i\hbar^{-1} [V; A_m(r)] \quad (3.25)$$

where V is the Hamiltonian of interaction. The equation following from (3.25) for the free field ($V = 0$) is well known in the classical theory. It describes the light propagation with diffraction taken into account in a quasioptical approximation.

In the equation (3.25) the Hamiltonian V describing the electromagnetic field interactions with the medium should be expressed in terms of local operators. As an example consider the Hamiltonian of the light interaction with atoms in the dipole approximation, which is often a basis for a variety of problems. It has the form

$$V(t) = \sum_{A, k} \frac{\hbar \omega_k}{2\epsilon_0 L^3} a_k \exp(-i\omega_k t + ik r_A) d_A(t) + \text{h.c.} \quad (3.26)$$

The equation (3.26) is given in the interaction picture, where d_A is the operator of the dipole moment for an atom located at a point r_A . Changing to local operators with the aid of (2.20), where the three-dimensionality should be taken into account, we will use the following approximations. Let the packet or the local mode interacts with the atom as a whole. It means that inside the band ω_m all frequencies $\omega_k \approx \omega_m$. Replace the atom position r_A by the position of the cell where this atom is located. Then the summation over the atoms may be divided into a sum over the cells l containing atoms and a sum over the atoms inside the cell. Suppose d_l is the operator of the atomic dipole moment in the cell l , then the equation (3.26) will take the form¹

$$V(t) = \sum_{m, l} \frac{\hbar \omega_m}{2\epsilon_0 a^3} A_m(l) \exp(-i\omega_m t + im l) d_l + \text{h.c.} \quad (3.27)$$

Here a new normalization volume a^3 appears, while the summation is performed only over the cells containing atoms. The obtained Hamiltonian describes the elementary interactions of the local field oscillators or photons in cells l with atoms located inside. As a result changing to local oscillators in the interaction Hamiltonian (3.26) reduces to an ordinary replacement: $a_k \rightarrow A_m(l)$, $k \rightarrow m$, $r \rightarrow l$.

In a number of cases effective interaction operators obtained, for example, by unitary transformations of the starting Hamiltonian ((3.26), are used for the description of multi-photon processes. Given below are two effective interaction operators for two-photon and parametric interactions in the presentation of local operators:

$$H_2 = \sum_{l_1, l_2} f_{12} A_{m_1}(l_1) A_{m_2}(l_2) S_1 \exp(i(\omega_{l_1} + \omega_{l_2})t) + \text{h.c.} \quad (3.28)$$

$$H_3 = \sum_{l_1, l_2, l_3} G_{123} A_{m_1}(l_1) A_{m_2}(l_2) A_{m_3}^y(l_3) \exp(i(\omega_{m_1} + \omega_{m_2} - \omega_{m_3})t) + \text{h.c.} \quad (3.29)$$

¹details of these and following calculations are given in [11]

where $f_{12}, G_{123} = g_{123} \exp(i(m_1 + m_2 - m_3)l)$ - are the coupling constants. The Hamiltonian H_2 describes two-photon interaction of the modes having frequencies in the region of two-photon resonance: $\omega_{21} = \omega_{k_1} + \omega_{k_2}$. The operator $S_1 = \sum_{ij} h_{ij}$ corresponds to the transition of atoms located at the point l from the lower to the upper working level. The process of parametric interaction of three waves in a transparent medium conforms to the Hamiltonian H_3 .

4 Integral of motion and statistics of the light

The integrals of motion may appear in the problems of light propagation in a medium, for which it is natural to use local description. The existence of the integrals enables to examine some peculiarities of the light statistics transformation without going into the solution of dynamic equations. Consider as an example two nonlinear processes: two-photon and parametric interactions, which are described by the effective Hamiltonians (3.28) and (3.29). Both processes are multimode, involving all pairs and triplets of modes having frequencies related as

$$\omega_{m_1} + \omega_{m_2} = \omega_{21} \quad (4.30)$$

$$\omega_{m_1} + \omega_{m_2} = \omega_{m_3} \quad (4.31)$$

Hereafter we restrict ourselves by the case of one-dimensional medium assuming the light propagating along the z -axis.

The pairs of modes m_1 and m_2 connected by the conditions (4.30) and (4.31) will be referred to as conjugated. For them the difference of the photon number operators

$$I = n_1(l) - n_2(l) \quad (4.32)$$

where $n_j = A_{m_j}^\dagger(l) A_{m_j}(l)$, $j = 1, 2$ commute with the Hamiltonian

$$[H_{2,3}; I] = 0 \quad (4.33)$$

Consequently I and any function of the form $f(I)$ are integrals of motion. For a problem with boundary conditions l will be considered as a continuous coordinate replacing $l \rightarrow z$. Introducing the new variables and assuming the light propagating along the z -axis: $t^0 = t - z/c; z^0 = z$. Then by virtue of (4.33) the difference of the number of photons in conjugated modes is conserved for any point of the medium:

$$I(z;t) = I(0;t - z/c) \quad (4.34)$$

With the use of the obtained operator integrals of motion it is possible to find the peculiarities of the transformation of the mutual correlation of conjugated modes for the propagation in a medium. Thus all statistical properties described by the correlation functions of directed intensity, e.g. of the form $\langle h(I(z;t))^p I(z;t+\tau)^q \rangle = K^{(p,q)}(z; \tau)$ are conserved in virtue of (4.34). From the standpoint of observation of direct interest is the lowest order correlation function and its Fourier image

$$K(z; \tau) = \int_0^z hI(z;t) I(z; t+\tau) \exp(i\omega\tau) d\tau \quad (4.35)$$

It is possible to measure the function (4.35) in a scheme with two photodetectors (Fig.1), where the detection spectrum of difference photocurrent $i^2(\tau)$ is observed. For this scheme

$$i^2(\tau) = \langle n_1 + n_2 \rangle + 2K_N(\tau) \quad (4.36)$$




Figure 1:

Here the first independent of frequency term is the shot noise. Index N indicates normal ordering of the field operators to describe a detector that responds at the absorption of photon. The rate of registration $\dot{P} = q \dot{m}$, where q is the quantum efficiency of the detectors, which for simplicity are assumed to be equal. The presence of the local mode $\dot{m} = c a$ in the expression for the photocurrent spectrum follows from the formulas of detection given in the presentation of local operators. This may be elucidated as follows. A wide-band photodetector is needed to describe the light detection. Therefore to register the local mode in front of the detector we have to place an optical filter of a bandwidth \dot{m} . Then \dot{m} that determines the rate at which the photons fall on the detector is the bandwidth of the scheme.

Assuming that at detector the local operators commute as free field operators (2.16) let us to write the correlation function (4.35) in the normal and time-ordered form $K(\tau) = K_N(\tau) + \dot{m} n_1 + n_2 i = \dot{m}$. Then taking into account the motion integral one finds

$$K_N(z; \tau) + \frac{1}{\dot{m}} \dot{m} n_1 + n_2 i = K_N(0; \tau) + \frac{1}{\dot{m}} \dot{m} n_{10} + n_{20} i \quad (4.37)$$

where $\dot{m} n_{10} i$ and $\dot{m} n_{20} i$ are the input photon number. As result the expression for the photocurrent or noise spectrum (4.36) takes the form

$$\dot{P}^2(z; \tau) = \dot{m} \dot{m} n_1 + n_2 i q (1 - q) + \dot{m} \dot{m} n_{10} + n_{20} i q (q - 1) + \dot{P}^2(0; \tau) \quad (4.38)$$

where

$$\dot{P}^2(0; \tau) = \dot{m} q \dot{m} n_{10} + n_{20} i + (\dot{m} q)^2 K_N(0; \tau) \quad (4.39)$$

is the input noise.

It is seen from (4.38) that for an ideal photodetector ($q = 1$) the input and output noise spectra of the light are equal. This means that the correlation of conjugated modes is conserved during propagation in the medium. Suppose that the input state of the modes is nonclassical, so it is a quantum correlation for which the shot noise is suppressed in bandwidth \dot{m} by a factor of $1 - q$:

$$\dot{P}^2(0; \tau) = \dot{m} q \dot{m} n_{10} + n_{20} i (1 - q) \quad (4.40)$$

The state of the light resulting in (4.40) produces, for example, by optical parametric amplifier (OPA). Then the level of the shot noise suppression in the output is unchanged:

$$\dot{P}^2(z; \tau) = \dot{m} q \dot{m} n_1 + n_2 i (1 - q) \quad (4.41)$$

This means that the quantum property of the light in the medium is conserved. Moreover in the medium with two-photon or parametric interaction the conjugated modes may be amplified, at that the observed initial correlation, in particular the quantum correlation, is conserved. This specific feature determines the properties of spontaneous radiation. Thus, if input is the vacuum state, then $\langle \hat{z}; \hat{z} \rangle = 0$ at the output. This means that the conjugated modes arising in spontaneous radiation have nonclassical correlation, that results in a suppression of shot noise. Indeed all these properties are following only from the existence of integrals of motion in systems expanded in space.

5 Local quasiprobabilities

In previous items the Heisenberg picture was introduced for local operators. In the Schrodinger picture the evolution is determined by the density matrix of electromagnetic field, which may be connected to c-number functions $P(f_k; g; s)$ called s -ordered quasiprobabilities. They arise in the density matrix expansion over operators $(f_k; g; s)$, which are Fourier-images of s -ordered displacement operators forming the complete set:

$$(f_k; g; s) = \frac{1}{Z} \int d^2_k \exp \left[f_k \hat{J}_k + (g_k - a_k) \hat{h}_k \right] \quad (5.42)$$

$d^2_k = \int d^2_k$. Then the s -ordered quasiprobability is determined by the expression

$$P(f_k; g; s) = \text{Sp} \left((f_k; g; s) \right) \quad (5.43)$$

Using the replacements $a_k \rightarrow a_m(l)$ in the above formulas we obtain the local quasiprobability $P(f_m(l); g; s)$. In the following we will restrict ourselves by the case $s=1$ corresponding to the normal ordering of the field operators which is described by the Glauber-Sudarshan quasiprobability $P(s=1) = P$. This function arises in the density matrix expansion over coherent states or diagonal representation. From the given expressions it is possible to obtain the relation between local and non-local quasiprobabilities. Similarly the case of operators, it is possible to proceed from one distribution function to another by the change of variables:

$$P(f_m(l); g) \rightarrow P(f_k; g) \quad (5.44)$$

$$f_m(l); g \rightarrow f_k; g \quad (5.45)$$

where the variables are related by the expression of the (2.6) and (2.7).

In the Schrodinger representation the peculiarities of the local description specified by the many-particle character of the problem become apparent to the same extent as in the Heisenberg representation. Here the transfer equations occur, which are formally equivalent to the BBGKI chains for the partial distribution functions. Of all the hierarchy of the distribution functions in the present case the following two types of one-particle quasiprobabilities are of interest. They arise from the function $P(f_m(l); g)$, of all modes of all local oscillators. Averaging over all oscillators except for the chosen one we will find the distribution function $P(f_m(l); g)$ for all modes of one local oscillator located at a point l . The averaging of $P(f_m(l); g)$ over all modes except for one results in a function $P(f_m(l))$ describing one mode of one local oscillator.

For a free evolution the introduced one-particle distribution functions $P_1 = P(f_m(l); g), P(f_m(l))$ obey the following transfer equation:

$$\frac{\partial}{\partial t} + c \frac{\partial}{\partial l} P_1 = 0 \quad (5.46)$$

In (5.46) the derivative over l describes the interaction of local oscillators or excitation transfer. Similarly to (2.13) it may be presented in the form

$$\frac{\partial}{\partial t} P(m, l) = \sum_{l' \in l} h_m(l; l') P(m, l') + \sum_{l' \in l} d^2_{m, l'} h_m(l; l') P(m, l'; m, l') \quad (5.47)$$

where $h_m(l; l') = i_{P^m}(l; l') (\partial = \partial_{m, l})$ is the differential operator of free evolution with Hamiltonian $H_0 = \sum_{m, l} h_m(l; l')$.

6 Fokker - Planck and Langevin equations

An approach based on the master equation for the quasiprobability $P(f, g)$ in the Fokker-Planck approximation is often used to describe the statistical properties of light. In the problems of the light interaction with atoms the master equation for the electromagnetic field can be obtained by an adiabatic elimination of atomic variables. It is not necessary to derive the field equation once again in terms of the local description, because one may immediately use the change of variables (5.45).

As an example consider the interaction of light with a two-level system in the lowest approximation for which the processes of the linear amplification or absorption type occur. Such a medium is described by a linear susceptibility, $\chi(k) = \frac{1}{2} d^2 (N_2 - N_1) \frac{1}{\gamma - i(\omega - \omega_0)}$ where d and ω_0 is the dipole moment and the frequency of the atomic transition, γ is the decay rate or transverse relaxation. The form of the field equation obtained by the adiabatic elimination of the fast atomic variables is well known (see, [9]):

$$\frac{\partial}{\partial t} P(f, g) = \sum_k A(k) \frac{\partial}{\partial k} P + Q(k) \frac{\partial^2}{\partial k^2} P + \text{c.c.} \quad (6.48)$$

Here the coefficients are defined by the linear susceptibility and populations of the upper N_2 and lower N_1 levels:

$$\begin{aligned} A(k) &= \frac{1}{2} d^2 (N_2 - N_1) \text{Im} \chi(k) - \frac{1}{2} d^2 (N_1 + N_2) \text{Re} \chi(k) \\ Q(k) &= \frac{1}{2} d^2 N_2 \text{Im} \chi(k) \\ \gamma &= \hbar^{-2} (\hbar \omega_0 = 2\pi \nu_0 L^3) \end{aligned}$$

When changing the variables we will assume that the wave packet forming the local mode interacts as a single whole. Then inside the band $k \in m$ it is possible to neglect the dispersion of all harmonics. This means that $A(k) = A(m)$, $Q(k) = Q(m)$. This approximation enables us to write at once the transfer equation for one-particle quasiprobability, say $P(f, g, l) = P_1$ in the form

$$\frac{\partial}{\partial t} P_1 + \frac{\partial}{\partial l} P_1 = \sum_m A(m) \frac{\partial}{\partial m} P_1 + Q(m) \frac{\partial^2}{\partial m^2} P_1 + \text{c.c.} \quad (6.49)$$

Notice that the structure of differential operators in right-hand side is the same in (6.49) and (6.48), and the whole of the passage procedure to the local description reduces to an addition of a derivative over l to the left-hand side of the equation.

The equation of quasiprobability (6.48) is in agreement with the following Langevin equations:

$$\begin{aligned} \frac{\partial}{\partial t} k &= A(k) k + f_k \\ \frac{\partial}{\partial t} k &= A(k) k + f_k^y \end{aligned} \quad (6.50)$$

where the correlator of random forces is defined by a dispersion coefficient

$$\langle f_k^y(t) f_k^y(t+\tau) \rangle = 2Q(k) \delta_{kk} \delta(\tau) \quad (6.51)$$

Inasmuch as the Langevin variables here are associated with the diagonal representation they correspond to the normally ordered averages of the field operators $\langle a_k^y(t) a_k^y(t+\tau) \rangle = \langle a_k(t) a_k(t+\tau) \rangle$.

In the local description the main moment in the formulation of Langevin equations is the finding of the random force correlator. To this end we use changing of variables of the form (2.19), (2.20), where $A_m(l) = \sum_k A_k(l) a_k$, thereby we introduce the local Langevin variable $f_m(l)$. Then the approximation $A(k) \approx A(m)$, $Q(k) \approx Q(m)$ used in writing down the equation for local quasiprobability leads to Langevin equations

$$\begin{aligned} \frac{\partial}{\partial t} + c \frac{\partial}{\partial l} f_m(l;t) &= A(m) f_m(l;t) + f_m(l) \\ \frac{\partial}{\partial t} + c \frac{\partial}{\partial l} f_m^y(l;t) &= A(m) f_m^y(l;t) + f_m^y(l) \end{aligned} \quad (6.52)$$

For the used variables there follows an expression for a random source

$$f_m(l;t) = \frac{1}{N} \sum_k f_k \exp(i(k-m)l + i(k-m)t) \quad (6.53)$$

From this we will find that the correlator of random sources as well as in (6.51) is defined by the dispersion coefficient

$$\langle f_m^y(l;t) f_m^y(l;t+\tau) \rangle = 2Q(m) \delta_{mm} \delta(\tau) \quad (6.54)$$

As to (6.54) it should be noted that during the transition to a local description with a coarse space scale a scaling time-function should be used defined as in (2.17) and the magnitude should be considered as a continuously coordinate. This implies that in (6.54) the replacements take place $\delta(k) \rightarrow \delta_a(k)$, $\delta(l) \rightarrow \delta_a(l)$. As a result the correlator of random sources in local Langevin equations has the form

$$\langle f_{m_1}^y(l_1;t) f_{m_2}^y(l_2;t+\tau) \rangle = 2Q(m) \delta_{m_1 m_2} \delta_a(l_1 - l_2) \delta_a(\tau) \quad (6.55)$$

In comparison with the ordinary description in (6.55) there arises a space delta-function due to the local correlation of oscillators in space.

7 Diffraction in linear medium

The differential operator in (3.25) becomes non-Hermitian due to diffraction effects, so the corresponding equation for quasiprobability becomes very complicated. However it is possible to use the Langevin formulation. In this way for the case of a linear medium considered above the change of variables in nonlocal Langevin equations (6.52) leads to Langevin equations which in the three-dimensional case have the form

$$\begin{aligned} \frac{\partial}{\partial t} + c \frac{\partial}{\partial z} + \frac{c^2}{2!_m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) f_m &= A(m) f_m + f_m(r;t) \\ \frac{\partial}{\partial t} + c \frac{\partial}{\partial z} + \frac{c^2}{2!_m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) f_m^y &= A(m) f_m^y + f_m^y(r;t) \end{aligned} \quad (7.56)$$

where the random force correlator is defined as

$$\langle f_m^x(r^0; t) f_m^y(r; t + \tau) \rangle = 2Q(m) m m a^3 \langle (r - r')_a \rangle \quad (7.57)$$

Here $Q(m)$ is the diffusion coefficient, which may be obtained from the corresponding (7.56) but nonlocal Fokker-Planck equation (6.48): $Q(m) = Q(k)$.

The equations (7.56) are linear in field amplitudes, so they are easily integrated. Let us denote the transverse radius-vector $s = (x; y)$ and introduce a travelling coordinate system $t^0 = t - z/c, dz^0 = dz$. Then for the given conditions at the boundary one finds

$$f_m(z; s; t) = \exp\left(\frac{A}{c}z\right) \int_{z_1}^z d^2s_1 f_m(0; s_1; t - z/c) U(0; s_1; z) + W(z; s; t - z/c) \quad (7.58)$$

$$W = \frac{1}{c} \int_0^z dz_1 \exp\left(\frac{A}{c}(z - z_1)\right) \int d^2s_1 f_m(z_1; s_1; t - z/c) U(z_1; s_1; z) \quad (7.59)$$

where the Green function is defined as

$$U(z_1; s_1; z) = \frac{m}{2(z - z_1)} \exp\left[-im \frac{(s - s_1)^2}{z - z_1}\right] \quad (7.60)$$

Here $f_m(0; s_1; t - z/c)$ is the transverse field distribution in the input, $A = A(m)$

Consider the properties of a random source W , defining its correlation function

$$\langle W(z; 0; t) W^y(z; s; t + \tau) \rangle = \frac{2}{c^2} Q a^3 \int_0^z \exp\left(\frac{A + A}{c}(z - z_1)\right) D(z - z_1) dz_1 \quad (7.61)$$

where the value

$$D = \frac{m}{2(z - z_1)} \exp\left[-im \frac{s^2}{z - z_1}\right] \int d^2s_1 \exp\left[-im \frac{s_1^2}{z - z_1}\right] \quad (7.62)$$

describes the diffraction effects. Since the diffusion coefficient Q is determined by the number of atoms at the upper level N_2 the transverse sizes of the medium where the light propagates should be taken into account. In this way the limits of integration over s_1 in D are defined. Thus $D = \delta(s)$ if the medium is not limited in the transverse direction. This means that the noise source is δ -correlated and the noise is white. Taking into consideration the finiteness of the size in transverse direction changes the situation. Thus in a medium confined by a cylinder with a radius R

$$D = \frac{mR}{2(z - z_1)s} J_1\left(\frac{m s R}{z - z_1}\right) \exp\left[-im \frac{s^2}{2(z - z_1)}\right] \quad (7.63)$$

where J_1 is the Bessel function.

References

- [1] I.V. Sokolov, M.I. Kolobov. Zh. Eksp. Teor. Fiz. 96, 1945, (1989)
M.I. Kolobov, I.V. Sokolov. Phys. Lett., 140, 101, (1989)
- [2] Yu.M. Golubev Zh. Eksp. Teor. Fiz. 65, 466, (1973)
I. Abram. Rhys. Rev. A 35, 4661, (1987)
C.M. Caves, D.D. Crouch. J. Opt. Soc. Am. B 4, 1553, (1987)
T.A. Kennedy, E.M. Wright. Phys. Rev. A 38, 212, (1988)

- [3] K. J. Blow, R. Loudon, S. J. D. Phoenix. *Phys. Rev. A* 42, 4102 (1990)
- [4] Yu. M. Golubev, V. N. Gorbachev. *Zh. Eksp. Teor. Fiz.*, 95, 475, (1989)
 V. N. Gorbachev, A. I. Trubilko. *Zh. Eksp. Teor. Fiz.*, 103, 1931, (1993)
 V. N. Gorbachev, A. I. Trubilko. *Opt. and Spektrosk.*, 80, 301, (1996)
- [5] K. J. Blow, R. Loudon, S. J. D. Phoenix. *J. Opt. Soc. Am. B* 8, 1750, (1991)
 K. J. Blow, R. Loudon, S. J. D. Phoenix. *Phys. Rev. A* 42, 8064 (1992)
- [6] V. N. Gorbachev, A. I. Trubilko. *Zh. Eksp. Teor. Fiz.*, 102, 1441, (1992)
 J. R. Jeers, N. Imoto, R. Loudon. *Phys. Rev. A* 47, 3346, (1993)
- [7] C. Kittel. *Quantum Theory of Solids*. New-York - London, 1963
 A. S. Davydov. *The solid state theory* Moskva Nauka. 1976
- [8] M. Scully, W. E. Lamb. *Phys. Rev.* 159, 208, (1967)
 H. Haken. *Laser Theory*. Encyclopedia of Physics 2nd edn vol. XXV/2c (Berlin: Springer). 1984.
 L. A. Lugiato, F. Casagrande, I. Pizzuto. *Phys. Rev. A* 26, 3438, (1982).
- [9] V. N. Gorbachev, A. I. Zhilba. *Quant. Opt.* 5, 193, (1993).
- [10] V. N. Gorbachev, A. I. Trubilko. *Optics and Spektroskopy*, 84, 879, (1998)
- [11] V. N. Gorbachev, A. I. Zhilba. *Transfer equation in statistical nonlinear optics*. Reprint-98, Tver state University. 1998.