

## Qubit and Entanglement assisted Optimal Entanglement Concentration

Somshubhro Bandyopadhyay

January 17, 2022

Department of Physics, Bose Institute, 93/1 A.P.C Road, Calcutta - 700009, India

## Abstract

We present two methods for optimal entanglement concentration from pure entangled states by local actions only. However a prior knowledge of the Schmidt coefficients is required. The first method is optimally efficient only when a finite ensemble of pure entangled states are available whereas the second method realizes the single pair optimal concentration probability. We also propose an entanglement assisted method which is again optimally efficient even for a single pair. We also discuss concentrating entanglement from  $N$ -partite cat like states.

## 1 Introduction

Quantum superposition principle gives rise to what is known as quantum entanglement [1], a non classical property exhibited by composite systems. By virtue of this property, subsystems of a composite system show nonlocal correlations between them and had been studied extensively in the context of EPR problem [2] and Bell's inequality [3]. However rapid developments in the last few years changed the scenario altogether. Now it is well understood that entanglement serves as an useful physical resource for information processing [4], and quantum computation [5] and allows manipulation like any other physical resources. Some key applications of entanglement include, quantum teleportation [6], dense coding [7], secure key distribution [8] and reduction of communication complexity [9]. Here one may note that maximally entangled states (Bell states) are essential for faithful quantum communication, for example teleportation [6] and secure quantum key distribution [8]. Therefore, protocols have been developed for obtaining a better entangled state from a less entangled one by local operations and classical communications. These processes are suitably termed as entanglement concentration [10, 11, 12, 13] when one extracts maximally entangled states (henceforth MES) from pure entangled states and purification or distillation [11, 14, 15] when MES are obtained from mixed entangled states.

The basic idea of entanglement concentration is the following: Two distant observers, Alice and Bob, are supplied with a finite ensemble of pure states from which they wish to extract maximum possible MES, where they are only allowed to perform local actions, e.g., unitary

---

dhom@bosemain.boseinst.emet.in

transformations and measurements, on their respective subsystems along with any auxiliary system (ancilla) they might prepare and classical communication.

The aim of this contribution is to present two methods for optimal entanglement concentration from pure entangled states using only local actions. For our methods to be successful Alice and Bob should know the Schmidt coefficients of the given entangled state(s). The first method becomes optimally efficient when a finite (not necessarily large) ensemble of pure states are available. The second method that we suggest however produces the optimal single pair concentration probability and possibly powerful than the first method. Besides we also propose an entanglement assisted concentration protocol. We show that if Alice uses an entangled state as an ancillary resource (for example, as in the protocol of Bose et. al. [12]) then one can obtain the optimal single pair concentration probability by opting for a different measurement scheme. We also discuss how the methods developed for entanglement concentration for bipartite systems can also be successfully applied in case of multipartite cat-like states.

**Qubit assisted methods:** The first method that we suggest requires Alice to prepare a qubit (ancilla: an auxiliary two level quantum system) in a state, say,  $|j\rangle$  (the coefficients of this state are initially chosen to be the Schmidt coefficients of the supplied entangled state). The procedure needs to be carried out separately on each member of the given ensemble. Thus, the ancilla qubit after being used once to purify a single pair, is brought back to the desired state by passing it through a polarizer for further application. Here we would like to point out that in order to obtain the optimal fraction of MES the method should be continued in an iterative fashion, in principle, indefinitely. Let us explain what we mean by this. Suppose Alice and Bob are initially supplied with  $N$  (as we shall see need not be necessarily very large) pure entangled states. After carrying out the protocol over all the members of this ensemble they are left with say  $N_1$  number of MES and  $(N - N_1)$  of less entangled pairs whereby they select the members of this less entangled subensemble, repeat the protocol and so on. This iterative process if continued indefinitely, Alice and Bob finally end up with the optimal fraction of MES. It may be worth mentioning that the present method doesn't require the supplied ensemble to be infinite (i.e., the optimal fraction is not approached asymptotically), but in practice the iterative procedure makes sense only when the supplied ensemble is reasonably large. Our method can also be understood intuitively from conservation of entanglement. As will be shown later that at every step of this concentration procedure average entanglement remains conserved implying that as MES are being produced the remaining pairs turn less entangled. Finally when the optimal fraction of the Bell states is obtained in the limit of an infinite sequence, the remaining pairs become totally disentangled.

The second protocol goes like this: Let us assume that Bob takes the responsibility of performing the desired local operations for entanglement concentration. He now prepares an ancillary qubit in state  $|j\rangle$ . The procedure now works in two steps. The first step involves in performing a CNOT on the two qubits that Bob holds. The second step is to perform an optimal state discrimination measurement (an optimal POVM) on any one of the qubits belonging to Bob. Consequently a conclusive result of such a measurement generates a maximally entangled state between Alice and Bob.

**Entanglement assisted method:** Here any one of the parties, say Alice requires to prepare an entangled state to implement the protocol. In Ref. [12] the authors proposed an optimally efficient entanglement assisted concentration protocol using entanglement swapping

[16]. However the method [12] is not optimally efficient for concentrating entanglement from a single pair. We show that resorting to a different measurement scheme one can however obtain the optimal single pair concentration probability.

**Multipartite entanglement Concentration:** Bipartite pure entangled states have unique representation through their Schmidt decomposable property. This makes dealing with bipartite pure states relatively easier than multipartite states not because of the larger number of parties being involved in the later case but for the fact that there is no unique representation for pure multipartite entangled states analogous to Schmidt decomposition. In this paper we treat the problem of multipartite entanglement concentration only for a restricted class of states, viz. the  $N$ -partite cat-like states and one should note that these type of states are Schmidt decomposable. One advantage of the methods that we developed for treating bipartite systems is that they are equally applicable for multipartite systems without any modifications whatsoever. Using them we show that the probability of entanglement concentration for multipartite cat-like states is same as that in bipartite systems. Thus the obtained concentration probability is conjectured to be optimal for multipartite systems that are Schmidt decomposable.

**Tools required for entanglement concentration:** Local Operations and Classical communication: The local operations that are in general used for entanglement concentration and distillation procedures include projective Von Neumann measurement, generalized measurements, in particular the POVM required for optimal state discrimination between two non orthogonal states [17], incomplete Bell measurements (for example, see Ref. [12]) and the CNOT (or quantum XOR) gate (an unitary transformation acting on pairs of spin-1/2 that flips the second spin if and only if the first spin is  $\uparrow$  i.e., it changes the second bit if the first bit is  $\uparrow$ <sup>1</sup> and is defined by the following transformation rules:  $|\uparrow\uparrow\rangle \rightarrow |\uparrow\uparrow\rangle; |\uparrow\downarrow\rangle \rightarrow |\downarrow\uparrow\rangle; |\downarrow\uparrow\rangle \rightarrow |\downarrow\downarrow\rangle; |\downarrow\downarrow\rangle \rightarrow |\downarrow\downarrow\rangle$ ).

Besides these, classical communication is an integral part of all protocols. It can be either two way or one way depending on the respective protocol. This is necessary to inform the partners about the result of the local quantum operations in order to select the successful cases.

This paper is arranged as follows. In Sec. 2 we present the qubit assisted entanglement concentration methods. In Sec. 3 we discuss entanglement assisted entanglement concentration. We propose a measurement scheme that produces the optimal single pair concentration probability. Sec. 4 is devoted to discussions regarding the relative merits of our schemes compared to the existing protocols [10, 11, 12]. Experimental feasibility of the suggested and the existing methods is also discussed. In Sec. 5 entanglement concentration from multipartite cat-like states is discussed. Finally in Sec. 6 we summarize and conclude.

## 2 Qubit assisted Entanglement Concentration

### 2.1 Proposal one:

Suppose Alice and Bob share a pure entangled state of the form ,

---

<sup>1</sup> In our notation  $\uparrow$  is  $|\uparrow\rangle$  and  $\downarrow$  is  $|\downarrow\rangle$ .

$$|j\rangle_{AB} = \sqrt{p_0}|0\rangle_{AB} + \sqrt{p_1}|1\rangle_{AB} \quad (1)$$

where we take  $p_j$  to be real and  $p_0 < p_1$ .

Alice prepares a qubit in the state,

$$|j\rangle_A = \sqrt{p_0}|0\rangle + \sqrt{p_1}|1\rangle \quad (2)$$

The preparation of the qubit in state (2) is crucial. Note that Alice should know the Schmidt coefficients of the supplied pure entangled state in order to prepare her ancillary qubit. Thus the combined state of the three qubits is given by,

$$|j\rangle_{AB} = |j\rangle_A |j\rangle_{AB} = \sqrt{p_0^2}|00\rangle_{A_1A_2B} + \sqrt{p_1^2}|11\rangle_{A_1A_2B} + \sqrt{p_0p_1}|01\rangle_{A_1A_2B} + \sqrt{p_1p_0}|10\rangle_{A_1A_2B} \quad (3)$$

The first two qubits belongs to Alice (denoted by  $A_1$  and  $A_2$ ) and the last one belongs to Bob. The entanglement concentration procedure involves two steps.

Step 1: Alice performs a CNOT operation on her two qubits. Bob doesn't need to do anything. This is the most difficult stage because to carry out CNOT operation is in no sense a trivial job. The resulting state turns out to be

$$|0\rangle_{AB} = \sqrt{p_0^2}|00\rangle_{A_1A_2B} + \sqrt{p_1^2}|11\rangle_{A_1A_2B} + \sqrt{p_0p_1}|10\rangle_{A_1A_2B} + \sqrt{p_1p_0}|01\rangle_{A_1A_2B} \quad (4)$$

Interchanging the position of the first two qubits since both belong to Alice Eq. (4) can be written as,

$$|0\rangle_{AB} = \sqrt{p_0^2}|00\rangle_{A_2A_1B} + \sqrt{p_1^2}|11\rangle_{A_2A_1B} + \sqrt{p_0p_1}|10\rangle_{A_2A_1B} + \sqrt{p_1p_0}|01\rangle_{A_2A_1B} \quad (5)$$

Step 2: This is an easy part where Alice performs a Von Neumann projective measurement on the qubit  $A_2$ , she holds i.e., she measures the z-component of the spin of qubit  $A_2$ . This is brought about by writing Eq. (5) as,

$$|0\rangle_{AB}^E = \sqrt{p_0} |0\rangle_{A_2} [\sqrt{p_0}|0\rangle + \sqrt{p_1}|1\rangle]_{A_1B} + \sqrt{p_1} |1\rangle_{A_2} [\sqrt{p_1}|0\rangle + \sqrt{p_0}|1\rangle]_{A_1B} \quad (6)$$

Thus if the outcome of Alice's measurement is  $|0\rangle$  i.e.  $|0\rangle_{A_2}$ , the resulting pair shared by Alice and Bob gets maximally entangled. Otherwise they come up with a lesser entangled state than what they initially shared. So the question is performing the above operations how often they succeed in getting a maximally entangled state. This can easily be seen by noting that the probability with which outcome  $|0\rangle$  is obtained is  $2^{-2}$ . This is in fact the single pair concentration probability using this method. However this is not the optimal probability. We now show that given a finite number of entangled states one can implement an iterative procedure to obtain the optimal fraction of maximally entangled states.

Suppose Alice and Bob initially shared  $N$  (which we shall presently see need not necessarily be very large) pure entangled states. The basic steps are the following:

(1) Applying the protocol over the  $N$  members individually, they end up with  $2N^{-2/2}$  number of MES.

(2) Now they pick out the remaining  $N(1 - 2^{-2/2}) = N(1 - 2^{-1})$  number of pairs which are not maximally entangled. Note that now each member of this less entangled sub ensemble are in a state given by

$$j_{11}i_{AB} = \alpha j_{00}i_{AB} + \beta j_{11}i_{AB}; \quad (7)$$

where  $\alpha = \frac{2^{-2/2}}{1 + 2^{-2/2}}$  and  $\beta = \frac{2^{-2/2}}{1 + 2^{-2/2}}$ . Accordingly, Alice prepares her qubit in the state

$$j_{11}i_A = \alpha j_{00} + \beta j_{11}; \quad (8)$$

and the single pair concentration procedure is applied again.

(3) This iterative procedure is continued indefinitely.

Now we show that the above procedure, when continued indefinitely, in the limit of an infinite sequence, the final ensemble generated comprise  $2^{-2}$  fraction of MES.

The proof is as follows: If they begin with  $N$  pair of pure entangled states and finally end up with  $N_{ME}$  number of MES, then the fraction of MES produced is given by,

$$\frac{N_{ME}}{N} = 2^{-2/2} + \frac{2^{-4/4}}{(1 + 2^{-4/4})} + \frac{2^{-8/8}}{(1 + 2^{-4/4})(1 + 2^{-8/8})} + \frac{2^{-16/16}}{(1 + 2^{-4/4})(1 + 2^{-8/8})(1 + 2^{-16/16})} + \dots \quad (9)$$

which can be rewritten as,

$$\frac{N_{ME}}{N} = 2^{-2/2} + 2^{-4/4} \frac{1}{(1 + x^4)} + \frac{x^4}{(1 + x^4)(1 + x^8)} + \frac{x^{12}}{(1 + x^4)(1 + x^8)(1 + x^{16})} + \dots \quad (10)$$

where  $0 < x = 2^{-1} < 1$ .

It is straightforward to show that the following infinite series

$$I = \frac{1}{(1 + x^4)} + \frac{x^4}{(1 + x^4)(1 + x^8)} + \frac{x^{12}}{(1 + x^4)(1 + x^8)(1 + x^{16})} + \frac{x^{28}}{(1 + x^4)(1 + x^8)(1 + x^{16})(1 + x^{32})} + \dots \quad (11)$$

uniformly converges to 1 for all  $x \in (0, 1)$ , whereby  $\frac{N_{ME}}{N} = 2^{-2}$ , known to be the optimal fraction of MES obtainable from pure entangled states. Hence our protocol indeed succeeds in extracting the optimal fraction of Bell states from an arbitrary number of pure entangled states. The efficiency of this method though optimal crucially depends on the rate of convergence of the series (11). However it is easy to see from (11) that the series converges very rapidly. From a practical point of view the optimal fraction is therefore approached very fast starting with a reasonable number of pure entangled states.

We now discuss the operational meaning of our protocol. We have seen that the optimal fraction is independent of the size of the ensemble. By this we mean that the optimal fraction of Bell states that can be obtained is not reached asymptotically i.e., it is not necessary to have an infinite ensemble. However, to achieve the optimal result the iteration procedure needs to be continued, in principle, indefinitely. However the rapid convergence of the series (11) ensures that, even in practice, to continue this iterative procedure in order to approach the optimal fraction we only need to have a reasonably sized ensemble. Note that for this method to be successful it is necessary to know  $\alpha$  and  $\beta$ , the Schmidt coefficients of the initially supplied pure entangled states. Classical communication is also required for Alice to convey her result to Bob in order to select the successful cases.

Now we show that a particular measure of entanglement viz. entanglement of single pair purification [12], is conserved on an average. We treat this conservation of entanglement in the same sense as discussed in Ref. [12]. We show that in our case also average entanglement is indeed conserved and therefore optimal in the sense that best combination of entangled states are obtained in the process. From the results of Lo and Popescu [13] it follows that initially the average values of entanglement shared between Alice and Bob is

$$E_{\text{before}} = 2^{-2} \quad (12)$$

where  $\alpha$  is the smaller Schmidt coefficient. After carrying out our protocol on a single pair the average entanglement shared by Alice and Bob is given by,

$$E_{\text{after}} = 2^{-4} + 2^{-2} \cdot 2^{-2} = 2^{-2} \quad (13)$$

Thus average entanglement is conserved at each step of the above procedure which implies that when the optimal fraction is reached, the remaining fraction becomes totally disentangled provided the process is continued indefinitely.

Now a few remarks regarding the efficiency of our method as compared to the other existing protocols [10, 11, 12]. As we have discussed earlier, to realize the optimal fraction of MES the iterative procedure needs to be continued indefinitely. But in practice the iterative procedure makes sense only when Alice and Bob have in their possession a reasonable number of pure entangled states to start with. Therefore we can only say that our method is as efficient as the other optimal ones [10, 11, 12]. As noted earlier the optimal fraction is approached very fast (see (11)) so any reasonably finite number of pure entangled states is required to implement this method successfully. However we note that a knowledge of Schmidt coefficients is necessary to implement our method and Procrustean method [10] whereas the Schmidt decomposition method, although works for any unknown ensemble of pure states but there the optimal fraction is approached asymptotically.

## 2.2 Proposal two:

Alice prepares an ancilla qubit in state  $|j\rangle_i$ . Thus the combined state is

$$|j\rangle_{AB} = |j00\rangle_{AB} + |j10\rangle_{AB} \quad (14)$$

where the first qubit belongs to Alice and the last two belong to Bob. Bob now subjects his two qubit to a CNOT operation whereby the new state given by

$$j_{AB} = |00\rangle_{AB} + |11\rangle_{AB} \quad (15)$$

can also be written as

$$j_{AB} = \frac{1}{\sqrt{2}} (|0\rangle_A + |1\rangle_A) \otimes (|0\rangle_B + |1\rangle_B) \quad (16)$$

where the states  $j_{AB}$  are defined by,

$$j_{AB} = \frac{1}{\sqrt{2}} (|00\rangle_{AB} + |11\rangle_{AB}) \quad (17)$$

From (16) it is clear that a state discrimination measurement which can conclusively distinguish between the two non orthogonal states  $(|0\rangle + |1\rangle)$  and  $(|0\rangle - |1\rangle)$  will give the desired result. Now, this optimal state discrimination measurement which is an optimal POVM measurement can be carried out on any one of the two qubits that Bob holds and let us assume that it is the second qubit on which such a measurement is performed. Note that the scalar product of these two nonorthogonal states is  $\frac{1}{2}$ . The respective positive operators that form an optimal POVM [17] are:

$$A_1 = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; A_2 = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}; A_3 = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (18)$$

The optimal probability of obtaining a conclusive result from such a generalized measurement (POVM) is  $\frac{1}{2}$ . It is clear that this is also being the probability of obtaining a maximally entangled state shared by Alice and Bob because a conclusive outcome implies that the entangled state shared by Alice and Bob is now given by either  $|j\rangle_{AB}$  or  $|j\rangle_{AB}$  depending on the state of the second qubit of Bob. For example, suppose Bob concludes that the state of his second qubit after the POVM measurement is  $(|0\rangle + |1\rangle)$ , then with certainty he also concludes that the maximally entangled state that he now shares with Alice is  $|j\rangle$ . Thus this method produces the optimal probability of entanglement concentration for a single pure entangled state.

### 3 Entanglement assisted Entanglement Concentration:

In the method that we now discuss Alice needs to prepare a similar entangled state locally. In Ref. [12] Bose et. al. proposed an optimally efficient protocol for entanglement concentration via entanglement swapping [16] where an ancillary entangled state is prepared beforehand to carry out the protocol. We note that the single pair concentration probability for a state of the form (1) as discussed in Ref. [12] is  $\frac{1}{2}$  and this is not the optimal value. Here we would like to point out that the first proposal of ours (see Sec. 2.1) succeeds in realizing the same single pair concentration probability using only a single qubit as an additional

resource. Since an entanglement is a more powerful resource than a qubit it is not unusual to suspect that a better measurement scheme might be devised which can improve the single pair concentration probability. This is what we suggest here. Of course the feasibility to realize our method experimentally is not very certain taking into account the present day technology. The advantage of the protocol of Bose et. al [12] is that their method can be successfully implemented in the laboratory with the present day technology.

We begin with the following facts. Alice prepares Alice and Bob share a pure entangled state of the form (1). Alice also locally prepares another entangled pair in the same state. Thus the combined state may be written as,

$$|j\rangle_{AB} = |j\rangle_A |j\rangle_{AB} = \frac{1}{\sqrt{2}}(|0\rangle_{A_1A_2} + |1\rangle_{A_1A_2}) \frac{1}{\sqrt{2}}(|0\rangle_{A_3B} + |1\rangle_{A_3B}) \quad (19)$$

where the subscripts  $A_1, A_2$  denote the qubits of the auxiliary entangled pair and the subscript  $A_3$  denotes the qubit that is the part of the entangled pair shared by Alice and Bob. Now we note that (19) can also be written as

$$|j\rangle_{AB} = \frac{1}{2}(|0\rangle_{A_1A_2} + |1\rangle_{A_1A_2}) \frac{1}{\sqrt{2}}(|0\rangle_{A_3B} + |1\rangle_{A_3B}) + \frac{1}{2}(|0\rangle_{A_1A_2} - |1\rangle_{A_1A_2}) \frac{1}{\sqrt{2}}(|0\rangle_{A_3B} - |1\rangle_{A_3B}) \quad (20)$$

Now the measurement part of Alice takes place in two steps:

Step 1: A measurement that projects the state onto either of the subspaces span by  $|0\rangle; |1\rangle$  or  $|1\rangle; |0\rangle$ .

Step 2: An appropriate measurement depending on the outcome of step 1 that generates a maximally entangled state between Alice and Bob.

First note that there are two possible outcomes of the measurement done in step 1 and consequently, measurement part of step 2 is to be defined accordingly.

Outcome one: Alice's measurement projects the state onto the subspace spanned by  $|0\rangle; |1\rangle$ . This happens with probability  $\frac{1}{4} + \frac{1}{4}$ . At this point Alice needs to perform a state discrimination procedure to discriminate between the two non orthogonal states (after normalization)

$$|j\rangle_{A_2A_3} = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \text{ and } |j\rangle_{A_2A_3} = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \text{ where } \frac{1}{\sqrt{2}} = \sqrt{\frac{2}{4+4}} \text{ and } \frac{1}{\sqrt{2}} = \sqrt{\frac{2}{4+4}}.$$

This generalized measurements are performed on the qubits  $A_2$  and  $A_3$  jointly. This is an important point to note. The optimal probability with which a conclusive result is obtained by performing an optimal POVM measurement is

$$p(\text{conclusive}) = 1 - \frac{1}{4} = \frac{3}{4} \quad (21)$$

It is clear from (20) that a conclusive result immediately implies that Alice and Bob share a maximally entangled state. For example if Alice concludes with certainty after the state discrimination measurement that the state is, say  $|j\rangle_{A_2A_3}$ , then it immediately follows that Alice and Bob now share the maximally entangled state  $|j\rangle_{AB}$ . We keep the subscripts  $A_i, B$  etc. in order to avoid any confusion. Note that, given "outcome 1" has occurred the single pair purification probability is just  $p(\text{conclusive})$ .



Outcome two: After performing the measurement defined in step 1 the other possible outcome is: the state is projected onto the subspace spanned by  $\{|01\rangle; |10\rangle\}$ . This outcome occurs with probability  $2^{-2}$ . This result when occurs actually simplifies the measurement part in step 2. Since now there is now no need to perform a POVM measurement. The measurement that needs to be performed in this case is an incomplete Bell measurement on the qubits  $A_2$  and  $A_3$ . It is clear from (22) that such a measurement always results in a maximally entangled state ( $|j+i\rangle$  or  $|j-i\rangle$ ) between Alice and Bob. Therefore, given "outcome 2" has occurred the single pair purification probability is 1.

Now the question is: What is the efficiency of the above scheme? Or, in other words what is the single pair concentration probability?

It is easy to obtain that the probability of single pair purification by implementing the above method denoted by  $p_{\text{SPC}}$  (SPC stands for single pair concentration) is:

$$p_{\text{SPC}} = 2^{-4} + 2^{-2} \cdot 2^{-2} = 2^{-2} \quad (22)$$

Thus the present method produces the optimal single pair concentration probability. In this method the additional resource required is an entangled state. However as we have seen in the previous section (Sec. 2.2) that to obtain the optimal probability, an ancillary qubit is sufficient. This implies that the qubit assisted method is a better one than the entanglement assisted method although both are able to convert a pure entangled state to a MES optimally.

## 4 Entanglement Concentration for N-partite Cat like states

We now proceed to show how our scheme works for multipartite entangled states. The method used above relied strongly on the existence of Schmidt decomposition for bipartite states. The difficulty in treating multipartite entangled states is that there are many possible forms of entanglement and there is no analogue to the Schmidt decomposition of bipartite systems. We therefore deal in particular with N-partite cat like pure entangled states. For simplicity let us first consider the following three partite state,

$$|j\rangle_{ABC} = |j00\rangle_{ABC} + |j11\rangle_{ABC} \quad (23)$$

Here our task becomes easier because the two proposals discussed in Sec. (2) can also be successfully applied for concentrating entanglement from these multipartite cat like states. Thus the methods for entanglement concentration from the state (23) proceeds exactly the same way as discussed in Sec. 2.1 and Sec. 2.2.

If we follow the scheme of Sec. 2.1 then Alice needs to prepare a qubit in the state defined by Eq. (2). She then performs a CNOT operation on her two particles and finally a Von Neumann projective measurement in the  $\{|0\rangle, |1\rangle\}$  basis. If the result of her measurement is  $|1\rangle$  which occurs with probability  $2^{-2}$ , Alice, Bob and Carol then end up with a GHZ state of the form,

$$|j\rangle_{ABC}^{\text{GHZ}} = \frac{1}{\sqrt{2}} (|j11\rangle_{ABC} + |j00\rangle_{ABC}) \quad (24)$$

Thus it turns out given a finite ensemble of the three partite entangled states of the form defined by (25) the maximum fraction of GHZ states obtainable is  $2^{-2}$ .

We can also follow the method discussed in Sec. 2.2. and the result is the same. The usefulness of the second proposal is that it does not require an ensemble to become successful. Thus given a single multipartite entangled state of the form (23), the probability with which one can successfully generate a GHZ state is  $2^{-2}$ .

It is clear that our scheme is trivially generalized to purify  $N$ -partite states of the form ,

$$|j\rangle_{1,2,\dots,N} = (|j0\dots0\rangle_{1,2,\dots,N} + |j1\dots1\rangle_{1,2,\dots,N}) \quad (25)$$

where, the maximum fractional yield for a finite ensemble remain the same as noted in case of bipartite systems. The probability that we obtain for concentrating entanglement for  $N$ -partite cat like states having the form (25) is  $2^{-2}$  and is conjectured to be optimal.

## 5 Discussion

Entanglement Concentrating procedures generate maximally entangled states which can be used for quantum communication with highest efficiency. The protocols that we discussed are state dependent in the sense that knowledge of the Schmidt coefficients is required.

It should be noted that the qubit assisted method discussed in Sec. 2.2 is better than the entanglement assisted method although both the protocols are optimal for a single pair. The advantage is two fold: First is it is easier to prepare a qubit in any desired state (pass it through a Stern-Gerlach apparatus appropriately oriented) than to prepare an entangled state. The second advantage is more important. The qubit can be reused once the operation is over for one pair. But in case of entanglement assisted process the auxiliary entangled state needs to be prepared for every individual pair because after a single operation the state gets destroyed.

One important issue is how many pure states are available to carry out the concentration protocols. It may so happen that only a limited number of entangled states are available. In that case one has to resort to the single pair concentration protocols and apply the methods on the members individually. However when an ensemble of pure entangled states are available one may apply single pair protocols on individual pairs or may use protocols that are not efficient for a single pair but becomes optimally efficient for a large number of supplied states, for example, the method suggested in Sec. 2.1. In this context an important issue is the experimental feasibility of the protocols.

For our methods to be successful we need a CNOT between one particle of the entangled pair and ancilla. But this is not something that can be implemented with photons as the technology stands today. On the other hand Procrustean method [10], though it involves a POVM, can be implemented with a polarization dependent beam splitter for photons. The scheme in Ref. [12], only needs incomplete Bell state measurements. However for ions entangled in distant traps, it is difficult to have a polarization dependent filter for the procrustean method. To purify by entanglement swapping would mean involving two more trapped ions. In such cases a scheme with only one ancilla ion on which only a CNOT is to be made will be very helpful.

## 6 Conclusions

In this paper we have described two optimal protocols for concentrating entanglement from pure bipartite entangled states. The first method becomes optimally efficient only when a reasonable number of pure states are made available whereas the second method is optimally efficient even for a single pair. We would like to stress that, although in principle, using the first qubit assisted method, one can extract the optimal fraction of MES from a finite ensemble of pure states provided the iterative procedure is carried on indefinitely but this iterative procedure makes sense in practice, only when Alice and Bob shares a reasonably sized ensemble of pure states. We also suggested an entanglement assisted concentration scheme which is also optimally efficient for a single pair. We also discussed why a qubit assisted method is better than the entanglement assisted one. Finally we have shown how these methods can be successfully used to concentrate entanglement from multipartite cat like states. The concentration probability thus obtained for N-party cat like states is found to be the same as that in bipartite systems and is conjectured to be optimal.

## Acknowledgments

I wish to thank Ujjwal Sen and Debasis Sarkar for useful comments and careful reading of the manuscript. I am also thankful to an anonymous referee for pointing out the experimental feasibility of different protocols.

## References

- [1] E. Schrodinger, *Naturwissenschaften* 23, 807 (1935); 23, 823 (1935); 23, 844 (1935); For a review see: M. B. Plenio and V. Vedral, *Cont. Phys.* 39, 431 (1998); Lecture notes of Lucien Hardy available at <http://www.qubit.org>.
- [2] A. Einstein, B. Podolski, and N. Rosen, *Phys. Rev.* 47, 777 (1935) [reprinted in *Quantum theory and Measurement*, edited by J. A. Wheeler and W. Z. Zurek (Princeton University Press, Princeton, 1983)].
- [3] J. S. Bell, *Physics* 1, 195 (1964) [reprinted in J. S. Bell, *Speakable and Unsayable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987), p. 14].
- [4] C. H. Bennett, *Physics Today* 48, 24 (1995); J. Preskill, *Proc. Roy. Soc. A: Math., Phys. and Eng.* 454, 469 (1998).
- [5] A brief but excellent article is by R. Jozsa, [quant-ph/9707034](http://arxiv.org/abs/quant-ph/9707034)
- [6] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres and W. K. Wootters, *Phys. Rev. Lett.* 70, 1895 (1993).
- [7] C. H. Bennett and S. J. Wiesner, *Phys. Rev. Lett.* 69, 2882 (1992).
- [8] A. K. Ekert, *Phys. Rev. Lett.* 67, 661 (1991).
- [9] H. Barnum, R. Cleve, and W. van Dam, [quant-ph/9705033](http://arxiv.org/abs/quant-ph/9705033).

- [10] C . H . Bennett, H . J . Bernstein, S . Popescu, and B . Schumacher, Phys. Rev. A 53, 2046 (1996).
- [11] M . M urao, M . B . Plenio, S . Popescu, V . Vedral and P . L . Knight, Phys. Rev. A 57, 4075 (1998).
- [12] S . Bose, V . Vedral and P . L . Knight, Phys. Rev. A 60, 194 (1999). Also at LANL archive, quant-ph/9812013.
- [13] H . K . Lo and S . Popescu, quant-ph/9707038.
- [14] C . H . Bennett, D . P . Di Vincenzo, J . A . Smolin and W . K . Wootters, Phys. Rev. A 54, 3824 (1996).
- [15] C . H . Bennett, G . Brassard, S . Popescu, B . Schumacher, J . A . Smolin and W . K . Wootters, Phys. Rev. Lett. 76, 722 (1996); D . Deutsch, A . Ekert, R . Jozsa, C . M acchiavello, S . Popescu and A . Sanpera, Phys. Rev. Lett. 77, 2818 (1996).
- [16] M . Zukowski, A . Zeilinger, M . A . Home, and A . K . Ekert, Phys. Rev. Lett. 71, 4287 (1993); For multiparticle generalizations see, S . Bose, V . Vedral and P . L . Knight, Phys. Rev. A 57, 822 (1998).
- [17] K . Kraus, States, Effects and Operations: Fundamental Notions of Quantum theory (Springer-Verlag, Berlin/New York, 1983); A . Peres, Quantum Theory: Concepts and Methods (Kluwer, Dordrecht, 1993), Ch. 9.